





Mauritii Antonii  
Bruno  
Medicinae Doctori  
1800.



VERMORE D. 46

VERR. E. 45

~~1003~~

~~N. d' inventario~~

371

~~371~~

X-K-22





Atque per in Valle (Circumscriptum)  
in (Circumscriptum) (Circumscriptum)  
in (Circumscriptum) (Circumscriptum)  
in (Circumscriptum) (Circumscriptum)

in (Circumscriptum) (Circumscriptum)  
in (Circumscriptum) (Circumscriptum)  
in (Circumscriptum) (Circumscriptum)

in (Circumscriptum) (Circumscriptum)  
in (Circumscriptum) (Circumscriptum)  
in (Circumscriptum) (Circumscriptum)

in (Circumscriptum) (Circumscriptum)  
in (Circumscriptum) (Circumscriptum)

in (Circumscriptum) (Circumscriptum)

in (Circumscriptum) (Circumscriptum)

in (Circumscriptum) (Circumscriptum)

in (Circumscriptum) (Circumscriptum)

in (Circumscriptum) (Circumscriptum)

in (Circumscriptum) (Circumscriptum)

in (Circumscriptum) (Circumscriptum)

in (Circumscriptum) (Circumscriptum)  
in (Circumscriptum) (Circumscriptum)

in (Circumscriptum) (Circumscriptum)  
in (Circumscriptum) (Circumscriptum)

in (Circumscriptum) (Circumscriptum)



*Stirpes, quas in Valle Stura Cuneensi,  
Thermis Vinadiensibus, locisque Circumvicinis  
Oviri tradidit Cl. Floræ sped. Auctor enumerantur;  
Ordine ipsius Auctoris servato.*

1. *Valeriana angustifolia.*  
Coppione Circathermas Vinadii in sacris frequent in Valle Ma-  
cra et Stura. Incolit loca lapidea et meridiano sole percutta.
2. ——— *montana.*  
In alpiibus Vinadiensibus
3. *Galium boreale.*  
G. foliis quaternis lanceolatis trinerviis glabris caule erecto semin. hispida.  
Circathermas Vinadii in sacris Sylvatum.
4. ——— *rotundifolium*  
prope thermas Vinadii Cl. Bellardi.
5. *Salvia verticillata.*  
Circathermas Vinadii.
6. *Thymus alpinus.*  
Circathermas Vinadii frequent est.
7. *Satureja montana.*  
In Valle Stura frequent est.
8. *Lavandula spica*  
ad Montes subalpinos Vinadii.
9. *Laminium ovata*  
Circathermas Vinadii.
10. *Betonica Monierii*  
In alpiibus Vinadiensibus.
11. *Melissa grandiflora.*  
Circathermas Vinadii.
12. *Buglossum Barrelieri.*  
In Valle Stura prope Demont.
13. *Spingicula lusitanica*  
Sp. nectaris apice incrassato  
Ex alpiibus Vinadiensibus Cl. Giavelli.
14. ——— *Villosa*  
Sp. scapo villosa flore minore calneo  
Ex alpiibus Vinadiensibus Cl. Giavelli.
15. *Toxaria Alpina.*  
Circathermas Vinadii in montibus le Vallon di S. Anna.



16. *Melampyrum Sylvaticum*  
In alpiibus Vinadiensibus.

17. *Euphrasia officinalis*  
C. L. alpina flava luteis floribus  
In alpiibus Circa Vinadii thermas

18. ——— *tricuspidata*  
C. fol. linear. tricuspidatis  
In alpiibus Vinadiensibus Cl. Gravelli.

19. *Pedicularis hibernica*  
P. caule simpliciter fl. spic. rostratis calyc. glabris aphodeli radice.  
In alpiibus Vinadiensibus prope lacellum J. Annae.

20. *Scrophularia Vernalis*  
In montibus Vinadii prope balnea.

21. *Linus alpinus* Calyc. 5-phyllus corol. pene infundibuliform. 5-fida vix  
bilabiata segmentis emarginatis capsula 4-valvis polysperma  
Didynamia angiospermia.  
Circa thermas Vinadii.

22. ~~*Primula*~~ *Primula viscosa*  
P. fol. viscosis odoratis linguiformibus sinuatis integerrimis  
folia 6-nervi olentia palmaria subtiliter villosa Cynogloss. Vulgari  
fol. similia Scapus palmariis flores coruleo pallidi 6. aut 7.  
Calyc. communiter 5-lobis 5-partitis foliolis quot sunt radii  
fructus maturus eximendus.  
Hae Thierper per errorem hic enumerata non ad Vinadii sed ad  
Valderi thermas pertinet.

22. *Primula integrifolia*  
P. fol. integerr. glabr. oblong. calycibus obtusis  
In alpiibus Vinadii.

23. *Gentiana punctata*  
G. coroll. subsidis campanul. punctat. Calycibus 5-dentatis  
pallidum florem habet punctis distinctum fol. ovato lanceol. nervos.  
Circa Vinadii thermas.

24. ——— *palustris*  
In Spongioris lac de la Maddalene in extrema Vinadii Valle  
Cl. Gravelli.

25. *Campanula latifolia*  
Circa Vinadii thermas.

26. *Phytolacca Halleri*  
P. foliis radicalibus cordatis acutis incisis tubis longissimis  
hirsutis emarginatis spica ovata obesa fl. purpureo coruleo  
Circa thermas Vinadii in pratis umbrosis.



Radice

Stem  
radice

radice  
radice

radice

radice

radice

radice  
radice

1. *Adiantum*  
2. *Asplenium*  
3. *Polypodium*  
4. *Marattia*  
5. *Psaronius*  
6. *Phlegmaria*  
7. *Phlegmaria*  
8. *Phlegmaria*  
9. *Phlegmaria*  
10. *Phlegmaria*  
11. *Phlegmaria*  
12. *Phlegmaria*  
13. *Phlegmaria*  
14. *Phlegmaria*  
15. *Phlegmaria*  
16. *Phlegmaria*  
17. *Phlegmaria*  
18. *Phlegmaria*  
19. *Phlegmaria*  
20. *Phlegmaria*  
21. *Phlegmaria*  
22. *Phlegmaria*  
23. *Phlegmaria*  
24. *Phlegmaria*  
25. *Phlegmaria*  
26. *Phlegmaria*  
27. *Phlegmaria*  
28. *Phlegmaria*  
29. *Phlegmaria*  
30. *Phlegmaria*  
31. *Phlegmaria*  
32. *Phlegmaria*  
33. *Phlegmaria*  
34. *Phlegmaria*  
35. *Phlegmaria*  
36. *Phlegmaria*  
37. *Phlegmaria*  
38. *Phlegmaria*  
39. *Phlegmaria*  
40. *Phlegmaria*  
41. *Phlegmaria*  
42. *Phlegmaria*  
43. *Phlegmaria*  
44. *Phlegmaria*  
45. *Phlegmaria*  
46. *Phlegmaria*  
47. *Phlegmaria*  
48. *Phlegmaria*  
49. *Phlegmaria*  
50. *Phlegmaria*



*[Faint, illegible handwritten text, likely bleed-through from the reverse side of the page.]*

27. *Scabiosa alp*  
*J. fol.*  
*prop*  
 28. *Cirsium*  
*hott*  
*Abn*  
 29. — *h*  
*fol.*  
*hinn*  
*Abn*  
 30. — *C. fol.*  
*caly.*  
*3n*  
 31. *Centaurea*  
*C. cal*  
*tuy to*  
*Supra*  
*J. fol*  
 32. *Artemisi*  
*Ar. fol*  
*Ar. rup*  
 33. *Achillea*  
*3n*  
 34. *Chamaem*  
*C. fol*  
*Ar. fol*  
*longi*  
*Ar. rup*  
 35. *Artemi*  
*3n*  
 36. *Artemi*  
*3n*  
 37. — *od*  
*3 fol.*  
*rande*  
*Cirsia, h*



27. *Scabiosa alpina*  
f. foliis pinnatis pinnis serratis capitulis globosis hispidis floribus cernuis  
prope Demente loco dicti le Courmale

28. *Cirsium pyrenaicum*  
C. foliis decurrentibus oblongo lanceolatis margine ciliatis et spi-  
natis utrinque tomentosis floribus tuberculatis.  
Abunde circa thermales Vinadii.

29. — *helenoides*  
fol. ciliat. subtus toment. radical. petiol. ovatis lanceolatis aut semi-  
pinnatis  
Abunde circa thermales Vinadii.

30. — *heterophyllum*  
C. fol. amplexicaulis lanceolatis ciliatis integris laciniatis. caul. subnatis  
caly. inermi.  
In jugo Col de la Madalene.

31. *Centaurea rhaphanifolia*  
C. calycibus scariosis fol. ovatis-oblongis denticulatis integris petiolatis sub-  
tus tomentosis  
Supra thermales Vinadienses edithora alpina parva Verne sanctuaria  
J. Anna inhabitat.

32. *Artemisia vuprestis*  
A. fol. pinnatis caul. adnatis fol. globosis cernuis recept. pappo.  
Abrupta lammatum alpinum incolit et frequens est in Valle Vinadii.

33. *Achillea herbacea*  
In montibus Vinadii.

34. *Chamaemelum montanum*  
C. foliis late pinnatis pinnis palmatis caule folioso unifloro Hall.  
A. fol. pinnatis multifidis planis lacinis linearibus acutis. W. p. pedunculo  
longissimo.  
Abunde in alpinis Vinadii.

35. *Aster amellus*  
In loco Vinadii dicto Nivajsa.

36. *Aster montanus*  
In alpinis Vinadiensibus

37. — *odoratus*  
3 fol. amplexicaulis dentatis hirsutis radical. ovatis caulinis lanceolatis  
caule paucifloro radice odora.  
Circum thermales Vinadii.



38. *Hieracium aurantiacum* H. fol. integr. caule tubundo simpliciss. ppter. longin-  
Circum thetmas Vinadii. b. fer.

39. *Scorzonera laciniata*  
S. fol. linear. dentat. acut. caul. erect. squam. calyc. spatulo-erectis  
In montibus di. Demonte.

40. *Lunaria rediviva*  
L. foliis alternis siliquis longioribus ovatis lanceolatis oblongis  
Circum thetmas Vinadii.

41. *Statice hectoria*  
Circum thetmas Vinadii: ad rupes prope fontes thetmarum.

42. *Cardamine alavisfolia*  
Vinadiensis: alpis: fol. fasciculatis.

43. *Sinapis maritima*  
Circum thetmas Vinadii  
fol. radical. long. petiolat. caulina parvis dentata sine triangulatis  
summa hastata succosa crassula glaberrima caul. glaberrimus  
calyc. patens

44. ——— *recurvata.*  
S. fol. radicalibus subuncinatis caulinis pinnatifidis siliquis re-  
curvatis  
Ad thetmas Vinadii: locis sterilibus.

45. *Silybum hieracifolium* fol. lanceolatis serratis  
In montibus Vinadiensis: frequens locis apertis

46. *Silybrium tanacetifolium*  
In pascuis. umbratis circum thetmas Vinadii: abunde.

47. *Epilobium dyneri*  
Jecus aquar. ad thetmas Vinadiensis.

48. *Anthyllis montana*  
Ad rupes Berzel in Valle Vinadii.

49. *Ononis Columna*  
O. fl. sessilibus fol. omn. ternat. stipul. lanceol. dentatis.  
Vinadii e viciis Sarcorum La Brece

50. *Hedysarum alpinum*  
Inter Vinay & le plance.



Handwritten text from the adjacent left page, including fragments like "Vest", "ingul", and "pust".

Main body of the page containing extremely faint, illegible handwritten text, possibly bleed-through from the reverse side.



Orobanchaceae  
O. fol. p.  
Citra

Hippocrepis  
Bn

Lathyrus  
L. fol. p.  
Bn

Ligustrum  
Bn

Bn

Juniperus  
Abund

Abund

Myrica  
Bn

Geranium  
Citra the  
B. p. d. b.

Ranunculus  
R. fol.  
R. p. d. b.  
Bn

Potentilla  
L. p. d. b.

Bn



51. *Orobanchus luteus*  
O. fol. pinnat. ovat. oblong. stipul. rotundato lunatis, caule simplici  
Circa thermas Vinadii Cl. Gravelli.

52. *Hippocrepis unisiliquata*  
In apertis secus Ichiator supra Vinadii thermas

53. *Lathyrus pinnatus*  
L. foliis cuneiformibus furcatis.  
In Valle fl. Thura prope Demont.

54. *Ligusticum meum*  
In pratis alpinis Vinadiensibus

55. — *ferulaeum*  
In alpiibus Vinadiensibus loco dicto le Barricade.

56. *Imyrenium olusatrum*  
Abunde circa thermas Vinadii.

57. — *nodiflorum*  
Abunde in sylvis proximisque pratis Vinadii.

58. *Myrrhis odorata*  
In pinguis pratis alpinis Vinadii.

59. *Geranium sylvaticum*  
Circa thermas Vinadienses in Valle di Lemie.  
G. ped. biflor. fol. subpeltat. 5-lob. inci. serrat. caul. erect. petal. emargin.

60. *Ranunculus thora*  
R. fol. inferior. subrotund. reniform. crenat. floral. ~~exsert.~~ lanceolat. in-  
tegrum. caul. infera. nud. caul. subflor.  
In alpiibus Vinadiensibus.

61. *Potentilla caulescens*  
e scopulis Vinadiensibus prope la Madonna del Vallon praevalit.

62. — *Valderia*  
In saxosis alpinis Vinadii.



63. *Potentilla grandiflora*  
Circum thermas Vinadii.

64. *Delphinium elatum*  
Circum Thermas Vinadienses in umbratis lacosis non rarum.

65. *Aquilegia virgata*  
In alpinis Vinadii: Cl. Bellardi.  
A. caule subnudo subunifloro foliis ternatis foliolis subtrilobis sessilibus.

66. *Helleborus foliosus*  
In alta Valle Thuræ Cuneensis.

67. *Saxifraga corymbosa*  
S. folior. ora cartilaginea caule triplicato variegato petalis immaculatis  
In Valle Thuræ loco dicto le Batticade.

68. *Dianthus superbus*  
Loco dicto Col d'Y mul. D. Giavelli.

69. *Silene cordifolia*  
Circum thermas Vinadii

70. *Viola alpina*  
Folia dentata uniformia ovata hanc ab aliis affinis distinguunt  
Cl. Bellardi secus torrentem Bschiator prope thermas Vinadii legit.

71. *Cistus serpyllifolius*  
C. suffruticosus stipulatus fol. herbac. venos. marginibus parvis ciliatis  
Circum thermas Vinadii in rupibus

72. *Dictamnus albus* Decandria Monogynia L.  
Cal. 5-phyllus deciduus corollae 5-petala subpapilionacea l. petal. descendente capsulae 5. margin. inter. Coarctae superiores deliscentes  
Cornutae polyspermae semina nigra calyptra elastica involuta.  
D. fol. pinn. caul. simplicis: odor naturalis vultu communis  
petala magna patentissima ex albo purpurascens.  
In alpinis Vinadiensibus Cl. Giavelli.

73. *Sedum rupestre*  
Circum Vinadii thermas locis siccis et apertis  
S. foliis subulatis 5-fariis confertis basi solutis floribus cymosis.

74. *Ribes alpinum*  
In dumetis et montanis proximis pratis, circum Vinadii thermas.



senilibus

calahj

ingruat  
legit.

calahj

et al de  
centu

munj

fil.

sum.



75. Ribes glo.  
R. var.  
Grossul.  
Ab.

76. Allium  
A. sea  
Bn.

77. Tulipa  
T. fl.  
Bn.  
High

78. Salix m.  
S. fo.  
Bn.

79. Alavum  
Circ.

80. Poa sea  
P. sp.  
Bn.

81. Festuca  
F. fac.  
much.  
C. A.

82. Pteris  
P. fo.  
legu.  
Bn.

83. Polyp.  
P. sp.  
Bn.

84. Ulva  
U. c.  
Bn.

85. Cacti.  
Loc.



75. *Ribes grossularia*

A. ramis aculeatis, petiolorum ciliis pilosis, bacis hirsutis  
*Grossularia* fuchs minimo hirsute margaritarum fere colore.  
 Abunde circa Vinadii Thermas.

76. *Allium triquetrum*

A. scapulo undato fol. triquetris stamin. simpl. caule triangulo.  
 In alpidibus Vinadii Cl. Giavelli.

77. *Tulipa sylvestris*. Corolla campanulata petal. oblongo cuspidatis, concavis.

T. flor. subnata fol. lanceolatis

In pratis alpidibus Vinadiensibus.

Stigma triangulare angulis divisis.

78. *Salix myrsinites*

S. foliis serratis glabris ovato lanceolatis venis julis tomentosis

In alpidibus Vinadii.

79. *Alnus europaea*

Circa Thermas Vinadienses.

80. *Poa setosoides*.

P. spica ovata locustis subserilibus seefloris obtusis.

In pratis elatioribus Vinadii Cl. Bellardi.

81. *Festuca phaeocoides*

F. racemo indiviso spicul. altera. n. brevilibus teretibus foliis involutis  
 mucronato pungentibus

C. Bellardi in Valle Sturae prope Dement.

82. *Pteris Crispa*

F. fol. duplicato pinnatis sterilibus obscure dentatis floralibus ovatis in-  
 tegerrimis.

In montibus Vinadii frequentissima.

83. *Polypodium Pontederæ*

P. pinnis paucis semipinnatis brevissimis acute serratis

In Sylvis umbratis ad Thermas Vinadii

84. *Ulva labyrinthiformis* fructificationes habet in membrana diaphana

U. cellulis labyrinthiformibus prominentibusque clavatis.

In thesauris Vinadii grad. lat. 50.

In Auctario Cl. Adlonis nullis plantis sponte natali  
 loco assignat Thermas Vinadienses nec Vallem Sturae

In Appendice

85. *Cacalia albifrons*

Locis humidioribus Vinadii.

Cunei 1800  
 Die 25. Maji.



Addenda

86. *Ranunculus palustris*  
in foliis Circa Demont.  
planta graminifolia ramosa aquatica:

87. *Cyananthemum alpinum*  
singularum varietatem vel distinctam speciem refulit D. Balby ex  
propositum per alpes Vinatii instituit, quod videtur  
*Leucanthemum pyrenaicum* minimum multifido incano folio.

---



77

lio.



Betonica Monieri hirtuta L.

B. spica majore hirtior labio superiore emarginato plano staminibus saepe  
longiuscula Chaire

B. spica bati foliola, corollae galea integra L.

obl. floribus purpureis differt a B. alopecuroides L. quod flores habet luteos.

Veronica austriaca

Caulis ferre desquipedatus villis folia varie ludunt plerumque  
abrotani similia quandoque Coronopifolia amulantur glabra racemi  
longi Calyces 5fid: Corolla pallide vel saepe caerulea

Phytneuma Haller

Rapunculus foliis radicalibus cordiformibus acute incisis, tubis longis.  
simis hirtis emarginatis spica ovata obesa. Haller.

Artemisia rupestris

A. foliis pinnatis caulibus adscendentibus floribus globosis cernuis  
receptaculo papposo Linn.

Abrutivum etc alio flodum foliis angustis sericeis pinnatis pinnis

3. fidis & 5. fidis Haller

Artemisia mutellina foliis petiolatis multifidis floribus pedunculatis  
axillaribus terminalibusq. receptaculis hirtis calyce anguloso villis.



Enumeratio plantarum, quas in Valle Gessii,  
prope Valdevii thermas, Oppida Envaque, Roascia,  
et Vicina loca Crevere adnotavit Cl. Allionius.

1. *Lavandula spica.*  
ad montes subalpinos Valdevii
2. *Lonicum ovula*  
Circa thermas Valdevii in saxosis umbratis.
3. *Betonica menierii*  
In alpibus Valdevianis
4. *Melissa grandiflora*  
Circa thermas Valdevii copiosissime.
5. *Melampyrum Sylvaticum*  
in summa sylva umbrosa thermis tepidi Valdevianis imminente  
abunde.
6. *Veronica austriaca* ~~LL~~  
V. racem. lateral. fol. linear-lanceol. pinnat. dent. tenuis. lacin.  
Cl. Bellardi in Montibus Roascia  
Mac V. verna varietas est All. auch.
7. *Jucculenta*  
V. erecta spicata, fol. succul. caulibus pinnato-dent. floral. integerrim.  
In saxosis appicis circa thermas Valdevianas
8. *Samolus Valevandi.* Pentandria Monogynia.  
Calyx 5 fidus Corolla 5-fida rotata squama muniens 5. Capsula  
unilocularis intera polysperma. 5 Valvis.  
Caul. ped- cubital. etiam ramulos fusco-virens fol. petiol. obovata  
subcrenata nitida fl. racemosi albi Caps. fusca glabra.  
Circa fontes calidiores Valdevii.
9. *Primula Vircola.* Vide plantas Vinadienses.  
Circa thermas Valdevii ad rupes.
10. *Gentiana asclepiadea*  
Circa Valdevii thermas
11. *Campanula ulicifolia*  
in umbratis sylvis Circa Valdevii thermas
12. *Phitheimma Halleri*  
Circa thermas Valdevii in pratis umbratis
13. *Artemisia vulgaris* abrupta summorum alpinum incolit &  
est frequens in valle Valdevii. & Vinadii



14. *Achillea herbacea*  
In montibus Valdevii.

15. *Chamaemelum montanum*  
In alpi-bus Valdevii.

16. *Matricaria inodora*  
M. receptac. hamisphaer. radii patentibus seminibus coronato mar-  
ginalis squamis calycinis margine obsolete.  
Differt. a M. chamomilla habitu firmiori defectu odori  
majori flore  
In montibus Valdevii.

17. *Linaria halmifolia* Vide Stirpes Vallis Maure.  
Alyrium fol. lanceol-lineur. acut. integerr. caul. procumb.  
perennant. flores racemosi primum umbellati albiissimi.  
Siliqua ovata compressa.  
In Montibus di. Roasca Cl. Bellard.

18. *Cardamine alarifolia*  
familiavit alpi-bus Valdevianis.

19. *Sisymbrium tanacetifolium*  
Circa thesmas Valdevii Coppioze.

20. *Brachiolobos pyrenaicus*  
in alpi-bus Valdevii.

21. *Smiranium nodiflorum*  
abunde in sylvis proximisque praetis Valdevii.

22. *Potentilla valderia*  
frequens in sacris alpi-bus Valdevii.

23. *Saxifraga stellata*  
abunde circa thesmas Valdevianas.

24. *Saxifraga pedemontana*  
Hujus insignis Valichy narsiber circa Thesmas Valdevii  
foliis 5. lobis lobo medio saepe 2. aut 3. pido petalis albis ad  
ungues roseis

25. *Silene cordifolia*  
Circa thesmas Valdevii.



Chamaemelum montanum

C. foliis laevibus pinnatis pinnis palmatis caule folioso (-floro. Haller.  
Anthemis foliis pinnato-multifidis planis laciniis linearibus acutis  
bispidis pedunculo longissimo Linn.

Caul. palm-terquiped. ramulus incurvus Folia Carrociniana apice  
latiora, & tantum a medio ad apicem pinnata laciniis ovatis acutis non  
linearibus. Ad basin petiolorum pinnata bipartita ut in A. arvensi.  
Flores Chrysanthemi Leucanthemi. Vitman.

Saxifraga pedemontana All.

S. caule erecto subnudo foliis trilobis & bidentatis calicinis palmatis  
Ad radicem haec habet rotundam foliorum quae ex angustis principio  
triangulariter amplificatur & in tres lineares lobos profunde secantur qui  
lobi bidentati sunt raro 4. dentati. Caulis inter ea erigitur folio uno circa  
medium donatus quod manus instar in 5. Lobos dividitur umbellatim  
quasi flores ex summo caule nascuntur ad pedunculorum ostium quin-  
que digito aut tridigito foliolo opposito. Sed singuliflores videntur  
pedunculo ter floribus longiore, & nudo. Flor. albus spatens petalis ex  
ungue ovatis etiam ellipticis calyce duplo longioribus. Calyx lobulatus  
profunde sectus in laciniis lanceolatas. Stamina cum antheris luteo  
virentibus calyce non altiora. Tota planta viscida & hirsuta est  
Circum Thermas Valderii eadem planta aliam faciem induit  
atque folia minime gerit profunde hispida sed pinnata 5-loba  
lobo medio saepe bispido aut 3-fido atque petala alba ad  
ungues rosea sunt. Allion



*Viola nummularifolia*  
*V. foliis cordato-acuminatis glabris integerrimis.*  
*V. fol. subrotund. obiter crenatis stipulis lanceolatis serratis caule bre.*  
*vicino bifloro.*

Obi. Folia in hac viola sunt cordato acuminata glabra integerrima  
 flori violæ marisæ æquali stipulæ tridentata. Alion.

Cette plante est très-petite elle n'a que 2. ou 3. pouces de haut  
 & les pedoncules sont les deux tiers de sa longueur, la racine est  
 vivace noirâtre & noueuse les feuilles sont presque rondes avec  
 une pointe presque insensible & une échancrure encore plus  
 petite à leur base elles sont toutes pétiolées n'ont que 3. à 4.  
 lignes de diamètre & autant de longueur elles ont à leur base  
 2. stipules lanceolés & dentés à dents aiguës & sensibles qui  
 regardent le sommet la fleur est petite penchée, bleu lavé ou  
 blanchâtre, le nectar est obfus la moitié plus long que le calice

Willard

*Viola Valdenia*

*V. erecta herbacea foliis ellipticis sinuatis non dentatis stipulis inte-*  
*gerrimis*

Radix hujus violæ est parva & albicans etc quæ cauliculus erectus vix uncin-  
 vum duarum altitudinem habens striatus & angulosus. Folia alternatim &  
 dense nascuntur. Radicatia aut inferiora pene orbiculata sunt reliqua ellip-  
 tica, aut longe elliptica petiolo pene longitudine folii & intus sulcata.  
 Foliorum aversa facies venosa est ambitus autem sinuatus & integerrimus Flores  
 etc alii foliorum nascuntur pedunculis plantæ altitudinem superantibus fere bi-  
 uncialibus Flor paullo minor mihi videtur et quam in *V. Calcarata*. Tubus apen-  
 crassus est & perforatus & stipulas præstant duo foliola sæpius inæqualia.  
 Obiecto habet folia sinuata non dentata stipulas etiam integras.

Alion



26. *Viola nummularifolia*  
Inter diffractos lapides summi et ediculi jugi Col de la femamorta  
ad alpes Valderianas pertinentis enascitur. Locis summi & alpe-  
ribus quae pertinent ad alpes d'entraive lantolla Tenda  
& Madonna delle Finestre.

27. *Viola Valdevia*  
Circa Valderii thermas in Saxosis torrentis ad radices montis. Boanni.  
Copiose circa la Madonna delle Finestre.

28. *Stellaria hypericifolia* Vide ~~Montes Valderii~~ *Macros*  
Copiosissime circa thermas Valderii.

29. *Sedum alpinifolium* Vide *Stirpes* in Valle Macros valcentis  
Abbellardi in montibus di Roalchia.

30. *Rubus cossii*  
Circa thermas Valderii.

31. *Tulipa sylvestris*  
in pratis alpiniis Valderianis.

32. *Asarum europaeum*  
Circa thermas Valderianas.

33. *Spheria crispata*  
in montibus Valderii frequentissima.

34. *Polypodium dryopteris*  
filiis ramolaminos. pinnulis dentatis  
frequentissima circa thermas Valderii.

35. *Uva labyrinthiformis*  
in thermis Valderii gradu Caloris 50.

### Auctarii

36. *Buglossum Barvelleri*  
in Vallibus di Vaudier et Entraque

37. *Phytolacca Carnetii* Vill.  
prope Entraque in regione dicta la Lanza.

38. *Hieracium ruppertii*  
In montibus d'entraive locis Saxosis et Calidis dictis la Lanza  
H. fol. ovatis. pilos. dentatis. laciniatis. caul. sub 1. floro non plio  
calycibus subnatis.

39. *Hieracium Anthybaceum*  
Circa thermas Valderianas.



40. *Turris stricta*  
T. glabra lucens caule simplicissimo erecto siliquis planis margine  
attenuatis  
prope thevma Valderii

41. *Selinum Seguietii* L.  
S. involucri universi nullo spiritibus diuaticatissimi  
prope entwaque.

42. *Geranium phaeum* prope entwaque.

43. *Cucubalus alpestri*  
In alpidibus di entwaive e rupium fissuris prodit locis umbris  
et dascosis

44. *Viola nummularia* folia  
locis humilis et alpidibus quae pertinent ad alpes di entwaive.

45. *Epipactis atrorubens*  
L. caule aphyllis flore resupinato labello ovato lanceolato calcar ovato  
turgido.  
labium bulbis compressis dent. caul. vagin. corollae resupinatis ne-  
ctarii labio indiviso  
In fageis prope entwaque locis umbris sub humidi.

46. *Corallorhiza neottia*. Genus Valde affine *Epipactis*: differt an-  
theris ymaternis  
Ophrys bulb. vacuol. flescos. caul. vagin. aphyll. nectar. lab. 3. fid.  
Orchance radice coralloide vuberrima.  
In fageis molineti in fageis prope entwaque.

47. *Zunera filiformis*  
locis paludosis di entwaive.

48. *Festuca altissima*.  
F. fol. plan. vilinearibus albo nervo divitis. Panícula contracta pedunc.  
communibus geminatis bulbis trifloris.  
locis laesidosis & umbris prope thevma Valderii et in Valle di  
entwaque prope lacum S. Joannis.



*Cucubalus Alpestris* All.

C. dichotomus foliis linearibus calycibus pelvisimilibus inflatis striatis  
petalis bipartitis.

Radice parum fibrosa inter rumpum fissuras oblique repit perenni, atque  
plures caules porrigit qui: primum geniculati sunt deinde se erigunt  
ad palmarem altitudinem rotundi glabri simplices: tota etiam planta glaber-  
rime est nihil habens viscosi aut villati. Folia linearia acuminata  
conglabra mollia ad 2. fere pollices longa de more gentis opposita & con-  
nata caulis summus præbet perennium aut alteram dichotomum  
flores singulares pedunculatos: pedunculus lateralis est brevior non bra-  
cteatus: calyce ovato live polius pelvisimiles sursum ampliores inflati  
glabri altera parte rubentes ore breviter 5-dentato dentibus opheliis.  
petala esse solent introflexa profundissime bifida ex his sulco rubentia  
intus sordide alba. Stylis squamæ nulla coronatice sed earum loco  
(non quidem semper duo parvi tumores.

Exhibet habitum suo J. Alpestrum Jacq. sed hæc & squamæ habet et  
petala bifida calycis etiam diversitatem. Aliden

*Festuca altissima*

Radix biennis & ultra. Radicis primarias truncas plures teretes fibras demittit. Ge-  
summa radice plures culmi teretes prodeunt ad 5. & ultra pedum altitudinem.  
los vertunt undique vagina foliorum usque ad prostratum folium. Hæ  
asperæ sunt & asperitudine sua superant foliorum superficiem & ambitum  
calamus vaginis denudatus lavis & striatus est. 4. circiter folia in unoquoque  
culmo esse solent. Sed præter hæc ad culmi basim 2. folia adiunt & recta  
multum reliquis breviora. Folia caulina recta ex vagina discedunt fere  
horizontalia 3. linear. & ultra latitudin. subus nervo albo divisa plana in  
aristam denique acuminata. Ad latera culmorum solet oriri fasciculus  
foliorum altero anno novam plantam daturus quæ pedalis & ultra al-  
titudinis sunt Culmi sunt simplices & ferunt paniculam non patentem  
sed collectam pedunculis erectis. pro planta altitudine brevis panicula  
est quæ scilicet 5. uncias fere non superet. Pedunculi primi:



¶ secundarii geminati nascuntur secundi pedunculi alternatim pro-  
ferunt locustas tres aut 4, quarum prima pedunculata est aliis  
sensilibus. Haec locusta interdum biflora saepe 3-flora glabra acumi-  
nata. Calyx flore brevior & angustior ex glumis angustis acuminatis  
persistenteribus, altera gluma multum brevior. Corolla florum ex glu-  
mis aequalibus glabris exteriori sub antherata facile caduca Semen  
gracile ——— Allion Auctar

9. Lichen  
f. loc.  
July




49. Lichen Cornucopioides.  
L. Scyphifer simplex folio brevior tuberculis coccineis  
Zutylui prope thesaurum Valderii locis umbratis et rupibus.

---

Cunei 27. Maji.  
1800.

---







11. ... in Nalle chavos







Itinerario in Valle d'Aosta



Observationes.

Valeriana morandea Vill.

Valeriana angustifolia Ses feuilles sont linéaires très étroites souvent par paquets aux aisselles des autres feuilles. Elle vient dans les endroits pierreux, sur les graviers, le long des torrents des montagnes. Villars.

Lunaria biolimyfolia

Cespitem facit cauliculorum vix erectorum robustior. Semipalmatis aut etiam altiorum. folia longe elliptica crassata concava glandulosa granulata cinereo virentia sessilia flores racemosi primum umbelati fl. albissimi patentissimi petal. 7 ovatis sep. levit. emargin. siliqua ovata compressa semina marginata. — All.

Scorzonera luciniata

S. fol. linear. dentatis pinnatifidis acutis caule erecto squamis calycinis paulo mucronatis  
La tige est perpendiculaire droite & ramifiée les fleurs sont jaunes petites dans un calyce anguleux dont les écailles sont terminées par un bouton coronné. Elle vient dans les lieux escarpés exposés au midi & paron: les bleds. Villars

Lythrum hiernifolium L.

La tige est haute de huit ponce environ simple & non ramifiée les feuilles inférieures sont dentées un peu blanchâtres ou cendrées celles de la tige deviennent linéaires. Les fleurs d'un jaune pâle terminent la plante. Les siliques sont dures un peu velues & rapprochées de la tige. Elle vient dans les montagnes pierreuses Villars



*Stirpes quae in Valle Maerae  
Crescere videntur. Cl. Allionius.*

1. *Valesiana angustifolia*  
V. fol. integr. longissim. et angustiss. flor. calcar. denato.  
In Valle Maerae.
2. *Satureja montana*  
In Valle Maerae legit Terraneus.
3. *Primula longiflora* / similis *primulae farinosa*.  
sp. fol. glabr. elliptic. rugos. serrat. flor. umbell. tubo longissimo.  
In Valle Maerae prope la Brusà & in alpidi Vraita.  
Segmenta flor. habet purpureo-violacea - A. A.
4. *Erica herbacea*  
Frequent in Valle Maerae.
5. *Atropa belladonna*.  
In Valle Maerae et tot.
6. *Santolina chamaecyparissus*  
S. pedunculis unifloris, foliis quadrifariis dentatis.  
Franciscus Peyroletti in Valle Maerae Collegit.
7. *Senecio parviflorus*.  
In alpidi Vallis Maerae.
8. *Cineraria alpina*  
in Valle Maerae.
9. *Scrophularia laciniata*  
In Valle Maerae locis siccis ad Viar prope Strop.
10. *Lunaria halimifolia*  
In aridioribus locis Vallis Maerae.
11. *Geranium Garreianum*.  
In Valle Maerae circa Strop.
12. *Oxyrimum hieracifolium*.  
In montibus Vallis Maerae frequenter locis apricis.  
L. foliis lanceolatis serratis.



13. *Trifolium pannonicum*.  
In hum. dir. alpiibus di. Skopj. et la Chianale.

14. *Ononis Cenisia*  
In summa Valle Maera loco dicto la Ciappa frequens est.  
O. glabra inermis, stipulis, dentato-vaginantibus, corollam vespillo reflexo  
purpureo, alis & carina n. v. vill.

15. *Astragalus Uvalensis*  
A. scapis, apophyllis, fol. ovato lanceolatis, sessilibus, siliquis hirsutis,  
hirsutis erectis ~~styliferis~~ styliferis. Haller.  
In alpiibus la Chianale.

16. *Pimpinella pumila* Genus involucri nulli. Semina intus plana, paulisper  
acuminata, dorso gibbo incistato.  
Frequens in Valle Maera  
Tragoselinum caul. crass. sulcat. divaricato fol. multif. capill.  
Sp. umbell. 7 numeroisimis simplicissimis, compositisque

17. *Cichus pilosus*.  
In Valle Maera nascitur.

18. *Sedum alpinefolium*. (a)  
Circa S. Damian in Valle Maera.

19. *Acorus Calamus*. *Hexandria monogynia* ?  
Gen. Clava ex sinu folii oriunda digitiformis, flor. sessil. un-  
digmetecta calice 6phyllis, foliolis obtusis, superne crassioribus  
pervirgatis, stylus brevissimus, capsula triangulatis, trilobatis,  
seminibus 3. feta.

Locis spongiosis in Valle Maera lectus est a Peyroletti.  
Radice horizontal. articulat. fol. radical. vaginantibus, ensiform. integris.  
odorata. Scapus apice foliosus. folii lobes sulcatum proferit spadicem  
nudum digitiform. cylindric. subnudentem fl. in spica attenuata.  
Variat clavis ternis, quaternis.

Cunei die 27. Maji 1800.

(a) *Sedum alpinefolium*  
S. erectum ramorum foliis planis, floribus albis, longe pe-  
dunculatis Allior.



Observationes

*Trifolium pannonicum*

*T. spicis villis longis corollis monopetalis calyce erecto*

villis nigris jaceq.

Cette espèce s'élève à un pied environ: les feuilles sont pointues très-velues ainsi que la tige: celle-ci se termine par un ou deux épis arrondis, que des corolles très-relevées blanches, ou jaunâtres beaucoup plus grandes que dans les autres espèces font paraître allongés —

Villars

*Ononis laevis.*

*O. pedunculis unifloris multicijs folijs ternatis cuneatis ligulatis serratis caulibus prostratis*

Cette espèce est petite quoique ligneuse elle est entièrement glabre excepté le calyce les folioles sont allongues avec des dentelures aiguës les fleurs naissent avec les feuilles supérieures: la capsule est glabre courte ventrée très-peu allongée. Villars

*Asragalus luteus*

*As. acutis scapo erecto folijs longiore leguminibus subulatis inflatis villis erectis. Vill.*

La racine est grosse mais simple ne produisant qu'une plante pour l'ordinaire les feuilles ne sont pas couchées par terre mais relevées le long du scape les scapes sont longs de six pouces & un peu plus droits & terminés par un épi de fleurs jaunâtres (ochroleucis). Les gousses sont droites rapprochées de la tige oblongues velues & pointues. Elle vient parmi les prés fertiles des Alpes dans les endroits herbueux & humides Vill.

*Pimpinella pumila* — *Dioica* L.

*Caulis divaricatus palmatis laevissimus parvis foliatis folia ternata live bignata linearia. Umbellae terminalis & laterales 2. oppositae plerumque compositae paucis simplicibus petala longe lanceolata incurva non emarginata Stamina alba longa involucrum nullum*

Dans la plante mâle & hermaphrodite — G. libert. Sp. plant.



*[Faint, mostly illegible handwritten text, likely bleed-through from the reverse side of the page. The text appears to be organized into several paragraphs or sections, with some lines being more legible than others.]*

Qles  
 et ad ja  
 hocavi  
 De  
 dium boreale  
 prope Ten  
 Operula horop  
 et foliis sen  
 Ad vup  
 Hymus alpinu  
 Inter Lino  
 piperu  
 ad oppida  
 Sideritis Rou  
 J. herbacea  
 riri ovate.  
 Cl. Bellard  
 Lemium orval  
 supra oppi  
 gargar  
 Ocher ci  
 Buglossum B  
 inter Ro  
 Scrophularia  
 Circa limon  
 Urtica alpe  
 Inter Te  
 Androsace v  
 in albiobu  
 Urtica herba  
 in subalpi  
 Alropana de  
 In Vall  
 Thymale  
 supra Te



*Clenebus* stirpium, quæ in Montibus Tenda,  
et adjacentibus Territoriis Carlin, Briga, Limon,  
Roccavione, Bovey, Valperio &c. nascuntur.  
Æthimonio Cl. Fl. Piedemontaneæ auctoris.

1. *Galium boreale*  
prope Tenda in herbidiis ad Mineræ fodinam.

2. *Asperula hirsutifolia*  
A. foliis densis lineariibus floribus umbellatis terminalibus subsessilibus  
Ad rupes prope Tendam nascitur.

3. *Thymus alpinus*  
Inter Limon et Tenda copiose.

4. *Piperella*  
ad oppida Carlin la Briga etc. e rupibus prodit.

5. *Sideritis Romana*  
S. herbacea decumbens ebracteata calycibus spinosis labiolis  
riori ovato.  
Cl. Bellardi in monte Tenda.

6. *Lamium ovata*  
supra oppidum Limon

7. *Garganicum*  
oritur circa Limon.

8. *Buglossum Battellieri*  
inter Roccavione & vobitant.

9. *Scrophularia vernalis*  
Circâ Limonem frequent

10. *Erinus alpinus*. Vide Plant. Vinadii.  
Inter Tenda et Savorio.

11. *Androsace villosa*  
in altioribus montibus supra Tenda.

12. *Erica herbacea*  
in subalpinis sterilibus supra Tenda.

13. *Atropa belladonna*  
In Valle di Perio prope monasterium charthusianorum.

14. *Thymus alpinus*  
supra Tendam et Limonem ad rupes Ignatii et Molineri.



15. *Thymalæa cheorum*  
In montibus di Tenda.
16. *Scabiosa alpina* Vid. Pl. Vindob.  
Plurima provenit in Valle di Perio & montibus di Tenda e la Buga
17. — *graminifolia*  
Erysimum fissurij prodit inter ulmeta et Carlino observ. Cl. Dana.  
Juxta Limone etiam occurrat.
18. *Globularia nudicaulis*  
G. fol. oblonge ovatis integerrimis caule nudo unifloro  
Cl. Bellardi in Montibus di Tenda.
19. *Lappula tomentosa*  
Prope Tenda frequens est.
20. *Carduus acanthoides*  
abunda prope Tenda.  
C. alis caulinis latiss. fol. semipinnat. pinn. angulis spinosis flor.  
longis fasciculatis
21. — *Collinoideus*  
C. fol. decurrentibus decussatis pinnatis pinnulis palmatis bipartitis  
dentatis lanatis caul. corymb. multiflor. flor. glomeratis.  
Circum Tenda
22. *Cirsium medium*.  
Carduus fol. decurr. pinnatis marg. spinos. caul. 1-floro peduncato  
longiss. calycem squamis internodiis patentibus setaceis.  
In montibus di Tenda
23. — *Ochroleucum*.  
C. fol. pinnatis pinnis ciliatis squam. calycin. recurvis.  
Indescentia di Col di Tenda Cl. Bellardi.
24. — *canum*  
Carduus fol. decurrent. lanceol. exsert. dentatis ciliatoaculeatis  
utrinque arachnoideo-subvillosis.  
in montibus Tenda.
25. — *helenioides*  
C. fol. ciliatis subtus tomentosis radiculis petiolatis ovato lanceol.  
aut semipinnatis  
Ignobis Molinari circa Tenda.



elabig

Dana

on/lor

4/12/1

duncalo

leah/

anuel



Cicuta fevora  
foliis decur  
Abunde in

causa tha  
supra find

Attemilia lo  
Jactu Vian

Achillea ha  
nullis

— m  
supra J

Cyanthem  
in montu

Matricaria  
inter li

Cineraria  
In mon

— la  
inter J

Arnica C  
A. uniflo  
inter

Hieraciu  
H. fol. inte  
In me

Prenant  
Gen. Cal  
quod m  
Vimplex  
P. folio  
prop.

Scorzonera  
In mon

Tragopo  
T. calyc  
Non inh



26. *Cnicus fevoris*  
C. foliis decurrentibus singulatis dentatis spinosis (aule ramo erecto.  
Abunde in montibus circa Tenda.

27. *Centauria thapsus* V. J. Vinadiensis.  
Supra Tendam loco dicto ripidis.

28. *Attemilia lobelia*  
Juncus Viam inter Tendam & Nicoram.

29. *Achillea herbacea*  
nullibi copiosius quam supra Tenda.

30. ——— magna  
Supra Tenda copiose nascitur.

31. *Gysanthemum Ceratophylloides*  
in montibus Tenda et Briga

32. *Matricaria inodora* Vide plant. Valder.  
Inter limen et Robilant.

33. *Cineraria alpina*  
In montibus Tenda Briga, limone

34. ——— longifolia  
Inter Tenda et la Madonna delle fenestre.

35. *Arnica Clusii*  
A. uniflor. fol. altern. ov-lanceol. hirs. integer. calyc. hirt.  
Inter Tenda & limone.

36. *Hieracium clavuliacum*  
H. fol. integr. caul. subnud. simpliciss. pilos. corymbifero.  
In montibus Tenda et altioribus.

37. *Prenanthes viminea*.  
Gen. Calyce cylindraceo ex paucis squamis, fere parallelis cum ali-  
quot minoribus ad basim. Semiflosculi in normis sepius 5. Pappus  
simplex stipitatus aut sessilis.  
Sp. foliorum ramentis, caul. adnatis, flor. sessil. planta villosa  
prope Tenda.

38. *Scorzonera laciniata* Vide pl. Vinad.  
In montibus di Tenda

39. *Tragopogon corymbosus*  
T. calycibus coroll. rad. longior. fol. integr. radical. pecunculatis. bas. villis.  
Non infrequens a jugo Col di Tenda ad Nicoram.



40. *Bilantella auriculata*  
B. Calycibus necario untrique gibbis siliculis in stylum coeuntibus.  
Circa la Briga Tenda.

41. *Thlaspi. lanceolite*  
Supra la Briga

42. *Thlaspi garreoliana*  
Supratendam in montibus editis inter Briga & Callia.

43. *Cardamine alavisolia*  
Secus torrentes supra Tendam.

44. *Sinapis maritima* Vid. plant. Vinad.  
in decembris di. Col di Tenda.

45. — *Jounefortii*  
Brassica fol. runcin. hispida. caul. hispida. siliq. torulosa. glabra. patent.  
rostra longissimo.  
locis rupestribus propeta Madonna delle fenestre.

46. *Heperis Sylvestris*  
In sylvis caducis circa Limone

47. — *laciniata*  
H. caul. ramosus. fol. inaeq. laciniatis. flor. sulphureo.

48. *Anthyllis montana*  
Circa Cerane supra Tenda

49. *Ononis villosa*  
Abunde provenit Tenda locis aridis  
O. pedunc. uniflor. avertat. fol. simpl. infim. ternat.

50. *Cytisus lupinus*  
in montibus ad Tendam pertinentibus  
C. fl. umbellat. terminal. ram. decumbent. foliol. ovatis.  
C. fl. axillar. ram. decumb. siliq. moll. lanug. pubescentibus.

51. *Genista pilosa*  
Inter habitant et occasione  
G. fol. lanceol. obtus. caul. tubercul. decumb. inermis.

52. *Asragalus depressus*  
A. acaulis. scap. fol. brevior. legum. cernuis foliol. subemarginatis.  
Cl. Bellardi: Copiose crescere vidit in jugo Col di Tenda dicto.  
Eucalyptus di Petris Petrus Cornalia.



104.

105.

106.



63 Pencedanu  
p. fol. 14  
3n comit

64 p. fol. pin  
Supra

65 Caraghitu  
l. fol. lip  
inter res

66 Jungerium  
ligusticum  
verh. 1/2  
Abund

67 Myrtis  
Cider

68 Geranium  
Sup

69 G. pedunc

70 Malva m  
3nter

71 Ranunc  
Ranunc  
nodoso  
ad petio  
pov. 1/2  
3n m  
pide Ca

72 A. fol.  
ceolato  
Cl

73 Potent  
p. fo  
Locit



53. *Pencedanum officinale*  
p. fol. ~~trifloris~~ quinquies 3 partitis filiformibus linearibus  
In comitatu nicæensi supra Tenda.

54. ——— minus  
p. fol. pinnat. foliol. pinnatifid. lacin. linear. oppos. caul. v. amovissim. patulo.  
Supra Tenda.

55. *Lathyrum gallicum*  
l. foliol. cuneiformibus parcat  
inter Tenda & Melicavina

56. *Juniperum nodiflorum*  
ligusticum nodiflor. fol. cordat. profunde serrat. acut. caul. v. amov.  
verticillato umbell. inordinat. numerosiss.  
Abunde Circa Tenda, Limone.

57. *Myrtus odorata*  
Circa Limone

58. *Geranium macrorrhizon*  
Supra Tendam observavit Ignatius Molineri.

59. ——— *pyrenaicum*  
g. pedunculis 2. floribus fol. multipid. lacin. obtus. inaequal. pet. 2. fid.

60. *Malva moschata*  
Inter Tenda & Limon.

61. *Ranunculus colummar.*  
*Ranunculus* fol. orbiculatum multipid. caule prostrato ad genicula  
nodoso fol. circumspersione robusta a breviter hinc glauca splend.  
ad petiol. usque 5. fida caulibus brevibus brevibus pedunculis 1. floribus  
porrigit flores habent amplius antheras luteas.  
In montibus di Limone loco dicti Vermelina ex diffraeto la.  
pide Calcaneo supra Castin. Cl. Dana.

62. ——— *alpestris*  
R. fol. radic. sulcordat. obtus. 3. partitis lobis 3 lobatis caulibus lan-  
ceolato integerrimo caule subuni floro.  
Cl. Dana in monte Relegà versus Tendam.

63. *Potentilla fruticosa*  
p. fol. pinnat. caul. fruticosa.  
Locis apricis in montibus 1. In perno dicti ad Tendam pertinentibus.



64. *Potentilla caulescens*.  
Supra Tendam occurrat.
65. *Potentilla Valdensis*  
Supra Tendam
66. *Aquilegia Virgata*  
In alpibus Tendae.
67. *Jasione pedemontana*  
J. caule erecto subnovo fol. tr. lob. et trident, caulibus palmatis  
In editis jugis di. Tenda et Limone.
68. *Silene Cordifolia*  
Supra Tenda.
69. *Viola palmata*  
V. caulis fol. reniform. glabris.  
Copiose ad fontem Sabunigenae circa Tendam.
70. *Arenaria montana* idest. *Capillacea*.  
A. fol. linear. lanceolat. scabr. calibus steril. longior. proeminent.  
Alpine fol. linear. acuminat. petal. flor. integr. calyc. duplo longioribus.  
In humo jugo Tendae la. Cola dicto
71. ~~*Potentilla*~~ *Petraea*  
A. fol. ovatis calicibus recurvis quadrifloriam imbricatis. flor. aggregat.  
In montibus Tendae locis siccis.  
Fl. in capitulis congestis 5. vel 6.
72. *Sedum alpine folium*.  
Sedum vancouver fol. plan. flor. alb. longe pedunculatis  
Inter Tendam & Robilant.
73. *Allium arenarium*  
A. caule planifolio bulbifero vaginis ferebilibus spatula multica staminibus brevissimis  
A. fol. ciliat. umbella bulbifera stamin. alterne bifida.  
Secus torrentes proximamque prata circa Tendam.
74. *Allium nigrum*  
in monte Vermelina prope Limon.
75. *Fritillaria meleagris*  
in pratis alpibus Tendae.
76. *Tulipa Sylvatica*  
In pratis alpibus Tendae.
77. *Semprevivum hirtum*.  
S. fol. caule petalorumq. apicibus hirsutis  
Uberrime in montibus ad Tendam spectantibus.
78. *Buscus sempervirens*  
inter Robilant & Rocavione.







*[Faint handwritten text, likely bleed-through from the reverse side of the page. The text is mostly illegible due to fading.]*

*[Handwritten notes on the right margin, likely from an adjacent page or a separate list.]*  
 Scleroidy  
 praktelation  
 cur molli  
 glum. 2flori  
 hiculata.  
 Bu palcu  
 usc. Baum  
 lypsa Cadu  
 fve albero  
 b. aphylla  
 luvul 2-3li  
 uli Capr. 3-  
 Bu Corticib  
 glosium Bo  
 Bu Vallid  
 Antisthinu  
 A. fol. plevi  
 inhabitat  
 Arimula glub  
 p. fol. Jov  
 Cicula e  
 Campanula  
 Thymelaea d  
 in Kap. Bay  
 Dicoic  
 Daphne floo  
 Supra Teni  
 Chamaemel  
 Bu jingo c  
 Doronicum  
 in montib  
 Hieracium  
 in pratib  
 in  
 Supra la  
 Arabis as  
 A. caule  
 silig. fere  
 ad oval ly



79. *Poa Selivoides* V. P. Vinadii  
in pratensatioribus Tenda.

80. *Holcus mollis*  
H. glum. 2 flori; nudiuscul. flolent. hermaphrodit. Anis. malculo arista  
geniculata.  
In pascuis elatioribus di Tenda Cl. Bellardi.

81. *Buscbaumia aphylla*. Gen. Anthera operculata hinc membrana rea  
Calyptra caduca inter operculum lacentis pollinis. Capsul. Ovali  
latere altero oblique plana orificio denticulato.  
B. aphylla pedunculo elongato capitulo l'allo bivalvi.  
Inscul 2-3 lineari. alter durius radice minimi ad basim rhipatus.  
tali caps. 3-lineari viridi.  
In corticibus Circa Tendam lecta est et D. Allionio communis.

in Auctario notata.

82. *Buglossum Barbatieri*  
In Valli di Simone ubique.

83. *Antirrhinum origanifolium*  
A. fol. pleriq. opposit. oblong. fl. r. albu.  
inhabitat rupes meridiem spectantes supra Tenda.

84. *Primula glutinosa*  
p. fol. serrat. glutinos. calycum laciniis capsula longioribus.  
Circa la Madonna delle finestre.

85. *Campanula latifolia* in pinguibus pratis di Petio.

86. *Thymelaea lanamunda*  
in rupibus meridiem spectantibus supra Tenda.

87. — dioica  
*Daphne* flor. axillar. geminis fol. linear-lanceol.  
supra Tenda in rupibus meridianum solem spectantibus.

88. *Chamaemelum montanum*  
In jago Madonna delle finestre.

89. *Doronicum austriacum*  
in montibus di Petio.

90. *Hieracium praemodum*  
in pratis, edilibus, della Madonna delle finestre

91. — intybaceum  
supra la Madonna delle finestre

92. *Arabis aspera*  
A. caule erecto flosculo ramoso fol. ovat. oppos. semiamplexicaulibus  
siliq. teretibus.  
Ad orosylvanum caduam supra Tenda.



93. *Arabis scabra*  
In alpidibus di Tenda.
94. *Turritia stricta*  
Supra Limone
95. *Vicia onobrychioides*  
abunde locis montanis & alpidibus di Tenda atque montium del Rifredo.
96. *Bypleurum petraeum*  
B. caule nudo fol. gramin. involueto et involuclis distinctis polyphyllis.  
In rupibus supra Limone loco dicto Armettin et Circa la Madonna delle finestre.
97. *Chorophyllum Sylvestris*  
prope castellum spiriti.
98. *Geranium phaeum*  
prope Robilant & Limone B. Molineri.
99. *Saxifraga mutata*  
plurima est locis umbratis humidis rupibus supra la Madonna delle finestre.
100. *Dianthus glaucus*  
in pascuis apricis et alpidibus di Tenda & Madonna delle finestre.
101. *Cucubalus alpestris*  
C. dichotomus fol. linear. calyc. pelvitorm. inflat. striat. petal. bipartit.  
In alpidibus di Tenda & rupibus firmissimis prodit locis umbratis & saxosis.
102. *Pyrola uniflora*  
Supra Tenda.
103. *Viola nummularia folia*  
locis summis alpidibus quae pertinent ad Alpes Tenda & Madonna delle finestre.
104. — *Valderia*  
copiose circa la Madonna delle finestre.
105. *Cirrus humulatus*  
C. exstipulatus suffrutescens. flor. subumbel. petal. lunula crocea notat.  
fol. oval.  
In extrema Valle spiriti, in montibus supra Limone.
106. *Fritillaria involuclata*  
F. fol. opposit. sum. teru. supra Tenda & il Rifredo locis saxosis  
& rupibus mare spectantibus.







17. unent filif  
Loc. 7 pal

18. Feghica herb  
F. fol. rad  
Lub. hirsut  
Loc. 1 m

19. Avena  
A. pan  
A.  
ti foli  
Abund

20. Avena  
Avena lu  
varicat  
In pa

21. Hordeum  
Circa Ca

22. Carex di  
Zucolit  
tem di

23. —  
frequen  
Lectum

24. Lichen  
L. umbilic  
Lapidib

25. Lichen  
L. fruti  
Cantibus Sc  
Rup  
Donna de

26. Lycopu  
hujus

27. Poa  
P. pa  
volu  
Zuc



107. *Juncus filiformis*  
locis paludosis alpinis di Tenda.

108. *Festuca heterophylla*  
F. fol. radical. setaceis, caulinis planis, panicula erecta glumis  
subhirsutis, margine scariosis subaristatis.  
Locis montanis siccis montium di Tenda.

109. *Avena sempervirens*  
~~A. paniculata patens locis siccis montibus floris albis.~~  
~~A. paniculata calycibus trifloris hermaphroditis, barilana-~~  
tis foliis viridibus acutis involutis sempervirentibus.  
Abund. in montibus di Castell. la Briga.

110. *Avena bromoides*  
*Avena subspicata*: spiculis binatis altera pedunculata arctis di  
varicatis calycibus 3-floris.  
In palen. siccis inter il Rifredo & Valpelio.

111. *Hordeum europaeum*  
Circa Carthesium di pelio locis siccis 3. Molineri.

112. *Carex dioica*  
Incolit prata montana di Rifredo quae est Tenda itur in Val-  
lem di pelio ad fontium frigidorum scaturigines.

113. — Bellardi  
frequens locis alpestribus solique patentibus alpinum Montega-  
lerium versus Castell.

114. *Lichen polyphyllus*.  
L. umbilicatus polyphyll. undique laevi, atro-virens venatus.  
Lapidibus adhaerens lectus est prope la Madonna delle Finestre.

115. *Lichen trisporus*  
L. reticulatus solidus ramulis compressis ramosis apicibus nigri-  
cantibus scutellis terminalibus.  
Rupes inhabitat, locis edicis & vento perflatis prope la Ma-  
donna delle Finestre.

Plantae quae pertinent in Appendice.

116. *Lycopus exaltatus*  
hujus varietas incana nascitur serus v. vultus circa Novesium

117. *Poa violacea*  
P. panicula erecta stricta spiculis compressis sub 4-floris co-  
volutis basi hirsutis.  
In montibus limoni locis siccis



118. *Poa stolonifera*  
*P. panicula expansa spiculis compressis seris flori, culmo stolonifero*  
*Mascitur in montibus Limoni et Vallispirii.*
119. *Festuca flavescens*  
*Non in frequen, locis sylvaticis subalpini, Vallispirii Limoni*  
*F. panicul. secunda erecta flavescens spiculis oblongis acutis*  
*in 65 flori fol. Capillari.*
120. *Avena setacea*  
*A. fol. setae. panicula purpurascens calycibus trifloris*  
*aviliis iniquis recurvis.*  
*In montibus Limoni locis calidioribus invenit D. Viale*
121. *Galium pusillum*  
*in Valle spirii sed planta major.*
122. ——— *montanum*  
*G. fol. tenuis linearis acuminatis glabris caule prostrato*  
*pedunculis trichotomis floribus subumbellatis.*  
*locis montanis Vallispirii Limoni.*
123. *Asperula cyanantha*  
*Stujus varietas vel distincta species crevit in rupibus*  
*Vallispirii Limoni Tendae.*
124. *Plantago argentea*  
*In montibus Vallispirii elatioribus parum a lanceolata diversa.*
125. *Supplementum incurvum.*  
*D. Viale in montibus limonenis.*
126. *Hyacinthus botryoides*  
*Fr. ligo prope Carthusiam.*
127. *Juncifraga lingulata*  
*J. fol. radicalibus longissimis lineariibus sulcat. cartilagineis integris.*  
*caule foliis paniculato.*  
*In Valle spirii, & montibus Limoni.*
128. *Juncifraga diaphanoides.*  
*In Valle spirii Fr. ligo in alpiibus limoni D. Viale. supra*  
*Tendam de Juffren.*
129. *Dianthus hirsutus*  
*D. flor. subaggregat. squam. calyc. vix aequant. corolla tonatim*  
*impressa caule foliisque laevibus.*  
*D. caule fol. scabri flor. panicul. squam. avit. calyce*  
*brevioribus*  
*In Monte Brimanda supra Bovitium locis sterilibus*
130. *Acora pimpinellifolia*  
*A. glom. globos. glab. peduncul. spat. rect. petal. scabr. foliol. obtus*  
*in montibus Tendae D. de Juffren.*



luna (luna)

in (in)  
gij (gij)

flor (flor)  
D. (D.)

rate

capit (capit)

divin (divin)

integ (integ)

supra

notum

calyre

y

o. o. o. o.



Rubus glau  
A. fol. quin  
mirch/ Hu  
Zn mon  
te Brin

Fr. lga

Anemo  
Zn al

Ranunc  
A. fol.  
fem. amp  
in pra  
D. Vial

Gerani  
Zn

Genit  
G. fol. r  
Zn

Heij  
Variet

Cacali  
inter 2

Artem  
prop

Erige  
L. cau  
in m

Ant  
A. e  
Non  
Pisid

Cen  
C. ca  
caul  
Zn

Caren  
Zn  
Non  
C. lpi  
mus



131. *Rubus glandulosus*  
 R. fol. pinnat. ternatis simplicibusque caulium spinis rubris villosis  
 mircis fructibus cymosis hirsutis  
 In montibus Vallis Pissii supra & circa Carthusiam. Copiose in mon-  
 te Brimanda supra Bovitium.
132. — *triphyllus*  
 Fr. ugo in Valle Pissii locis asperioribus.
133. *Anemone dubia*  
 In alpidibus Limonensibus D. Viale
134. *Ranunculus lacerus*  
 R. fol. radical. petiolat. cuneiform. laceris apice multifidis caulinis  
 semiamplexicaulis laciniatis summit. densibus linear. caul. laev. ramis  
 in pratis summorum alpidum limonensium locis humidioribus  
 D. Viale.
135. *Geranium fuscum*  
 In pratis Vallis Pissii.
136. *Genista scoparia*  
 G. fol. simpl. hirsut. ram. 10. angulat. cinereis.  
 In rupibus limoni a Sole percutitis
137. *Hedysarum onobrichis*  
 Varietas incana fol. longior nascitur in montibus Vallis Pissii.
138. *Cacalia albifrons*  
 inter distractos lapides Vallis Pissii limoni
139. *Artemisia camphorata*  
 prope Rocca di Vico.
140. *Erigeron Villarii*  
 E. caule erecto fol. oblong. lanceolat. binerviis fl. corymbos.  
 in montibus Carthusiae Vall. Pissii Fr. ugo. raro.
141. *Anthemis altissima*  
 A. erecta fol. pinnat. pinnar. 6-8 denticul. reflexis asperis  
 non infrequenter circa Bovitium inter segetes huminagris Vallis  
 Pissii prope Carthusiam
142. *Centauria nudicaulis*  
 C. calyc. setac. - spinos. squamul. fol. indivis. superior. subdent.  
 caule nudiusculo simpliciter  
 In montibus Vallis Pissii supra Carthusiam locis sylvest. Fr. ugo.
143. *Carex foliacea*  
 In Valle Pissii prope Carthusiam locis humidioribus D. Viale ex  
 montibus limoni  
 C. spical. subobov. sessil. remot. aedrogya caput. ovata. ferebimur  
 mutic. divaricat.



144. *Salsola pouteredera*.  
*Sals. fol. dur. oblong-lanceolat serrat. glandulos intern. glaucis jatis*  
*oblongis basi foliosis.*  
*In altitudinis limoni D. Viale.*
145. — *Daphnoides* !  
*S. fol. elliptico acuminatis subseratis inferne glaucis superne nitidis*  
*julis brevibus dense tomentosis.*  
*In Valle Spisii et in montibus limoni. 147.*  
*Stevia cretica*  
*in Sylvaticis et umbratis Jendoo.*
146. *Polypodium limborum*  
*In monte Bissimanda supra Bovitium. in Valle Spisii Fr. Ugo.*
147. *Picea minima* frondibus glabris bipartitis acutis  
*in Valle Spisii supra Castanetum locis humidis.*
148. *Lichen peltiphyllus*  
*L. subimbricat. foliol. orbiculat. lobatis. rub. marg. alb. tuberc. nigri.*  
*In altitudinis Vallis Spisii Fr. Ugo.*
149. — *Leucigerus*  
*L. crustaceus albicans sublobat. scutell. albo-margin. asate flavescens.*  
*In Valle Spisii*
150. — *Carpitum*  
*L. imbricat. cinereo-virescent sublobato margine albo tuberculis carneis*  
*lenitissimis saepe congestis*  
*Ex montibus limoni D. Viale*
151. *Lichen velleiformis*  
*L. umbilicatus subtus hirsutus peltis imbricatis subtus saeculis*  
*supra Jacea in montibus limoni D. Viale.*
152. *Boletus hirsutus*  
*Clavarium squamosum superne hirsutum & obscurum subtus etc pulvis*  
*aureum dentissime et tenuissime perforatum.*  
*In Valle Spisii ad caudices Castaneorum.*
153. — *Cinnabarinus*  
*B. acutis pulvinatis tomentis et foris ruber*  
*Fr. Ugo reperit supra truncos emaroidos in Valle Spisii.*
154. — *Uredula sparis*  
*B. crustaceus effusus farinosus albus in montibus Bovirii & Valle Spisii.*
155. *Clavaria cinerea* ramis rimosis apic. multis.  
*ad caudices arborum sprucecentium prope Bovitium.*
156. *Lycoperdon epidendrum* cortice farinaceo purpureo  
*supra truncos emaroidos fagi Sylvaticae in Sylvis prope Ch. Vall. Spisii.*
157. *Sphaeria vagiformis*  
*Ex montibus Castanetis Fr. Ugo. plantarum recent. in adultis nigra.*

Cunei die 29. Maji 1800.



19 July

18 July

17 July

16 July

15 July

14 July

13 July

12 July

11 July

10 July

9 July

8 July

7 July



123.

5



Handwritten text at the top of the page, possibly a title or header.

Handwritten text block, likely the beginning of a paragraph.

Handwritten text block, continuing the narrative or list.

Handwritten text block, continuing the narrative or list.

Handwritten text block, continuing the narrative or list.

Handwritten text block, continuing the narrative or list.

Handwritten text block, continuing the narrative or list.

Handwritten text block, continuing the narrative or list.

Handwritten text block, continuing the narrative or list.

Handwritten text block, continuing the narrative or list.

Handwritten text block, continuing the narrative or list.

Handwritten text block, continuing the narrative or list.

Handwritten text block, continuing the narrative or list.

Handwritten text block, continuing the narrative or list.

Handwritten text block at the bottom of the page, possibly a conclusion or signature.



Ceva, l.

1. *Valeriana*  
in *Flora*  
in *Monte*  
in *Monte*

2. *Thymus*  
in *Monte*  
in *Monte*

3. *Thymus*  
in *Monte*

4. *Thymus*  
in *Monte*  
in *Monte*  
in *Monte*  
in *Monte*  
in *Monte*  
in *Monte*

5. *Thymus*  
in *Monte*

6. *Thymus*  
in *Monte*

7. *Thymus*  
in *Monte*

8. *Thymus*  
in *Monte*  
in *Monte*  
in *Monte*  
in *Monte*  
in *Monte*

9. *Thymus*  
in *Monte*

10. *Thymus*  
in *Monte*  
in *Monte*  
in *Monte*  
in *Monte*  
in *Monte*  
in *Monte*



Plantae quae in Provincia Monvegalensi et circa  
Ceva, Garretto, Ormea crucere adnotavit Cl. Almonius  
in Flora sua Pedemontana.

1. *Valeriana tuberosa*  
*N. foliis triandris fol. radical. lanceolatis integerrimis, reliquis pinnatifidis.*  
*Montes montana radice olivari.*  
*Inter Garretto & Ormea loco dicti la fana dei Ovi*
2. *Thymus piperella*  
*3 natis pibus maritimis quae pertinent ad Monvegale. provinc. ad oppid.*  
*Garretto.*
3. *Hyssopus officinalis*  
*Oritur inter Garretto & Ormea.*
4. *Thymus spicata* Gen. Cal. subcylindric. 2. labiat. utrinq. lineavil-  
*losa excavatus. Corol. galea obtusa 2-fida barba 3-fida, laciniis fere*  
*aequalibus stylis semibifidis.*  
*Montes inter Garretto & Ormea*  
*Caul. dodrant. lignori, densiculati in can. Fol. hyssop. rigid. atroviscent.*  
*Minoribus intermixtis foliol. in sum. Caulib. margin. laevigi-*  
*nato. Fl. ex axillis bractear. terni purpur.*
5. *Laminium ovula*  
*in montibus umbratis Monvegalensi provinciae.*
6. *Pentstemon lucidum*  
*in sylvis Garretianis copiose.*
7. *Symphytum tuberosum*  
*J. foliis summis oppositis*  
*in montibus Monvegalensibus copiosissime.*
8. *Coris monspeliensis* Gen. Calyc. 10. dent. Coroll. parum 5-fida  
*capitata. Natu. Pentandria monogynia.*  
*Circa Garretium.*  
*Caul. strigosus palmis. ruber fol. altern. spars. linear. patentia*  
*crassiuscula succulenta ciliata Fl. capitata spicati corollae ex*  
*purpur. caerule.*
9. *Scrophularia veruialis*  
*copiose in montibus Monvegalensibus*
10. *Monarda fontana* Triandria 3-gynia  
*Calyc. 2. 3-phyllus corollae 1. petal. irregul. parva. 5-part. lacin.*  
*3. altern. min. Capsul. 1. locula 3. valu. sapient 3-lyseria.*  
*ad fontem scaturigine, circa Garretium.*  
*Caul. procumb. ramis. articul. in part. vix semiped. in gla-*  
*veol. subune. Fol. petiol. ovat. vel oblong. Flor. racemos. pendul.*  
*exiguu. albi. raro se se apperientes.*



11. *Vimula longiflora*  
In alpidibus Garrescio le Carente
12. *Lyrimachia nemorum*  
ad Torrentem oppidi: Monastero Monti, regali
13. *Campanula bononiensis* — petra in alpidibus di Cusotto.  
C. fol. ovato-lanceol. subtus scabris sessilibus caule paniculato.  
In aridis locis Grappiol dicitur supra Garrescio.
14. *Phytolacca Halleri*  
in stratis humosis alpidum Monregalerensium
15. *Thymelea alpina*  
Inter ormeam et Garrescium Zynatini Molineri  
Daphne flor. sessil. aggregat. lateral. foliis lanceolatis ob.  
tusiusculis subtus tomentosis.
16. *Poterium spinosum*  
sp. spinis ramosis  
In montibus circa Garrescium non in frequent.
17. *Onopordum rotundi folium*  
In alpidibus Monregalerensibus
18. *Carduus acanthoides*  
Circum Garrescio.
19. *Carduus acanthoides*  
C. fol. decurrentibus decurvis pinnatis pinnulis ~~graculatis~~ 4. pidi  
aculeatis lanatis. Caul. corymb. multiflor. floribus glomeratis.  
Circum Garrescio.
20. *Cirsium Carniolicum* *Carduus rivularis* L.  
Carduus fol. inferne pinnatis pidi in lacinias oblongas superne integris  
& serratis inermi-spinulosis sessilibus caule inermi stipitibus.  
ad fluentes aquas supra Garrescium.
21. *Artemisia lobelia*  
Supra Garrescium loco dicto il Garbo della luna.
22. *ponthica*  
Supra Garrescium cum precedenti eodem loco crescit.
23. *Chrysocoma linosyris* Gen. Calyce hemisphaerico imbricato. Fl. her.  
maphrod. uniformes. 5. pidi Stylus sine florentis longior Receptac.  
hudsonii seminum spargium simplex.  
C. herbacea fol. lineari. glabr. calyc. lac. squarrosis.  
Planta digitis hirsuta odorem aromaticum spirat.  
In provincia Monti regali saepe nascitur.



Ho.

706.

17

reg.

Fl. ker.  
h.c.

*[Faint, illegible handwritten text, likely bleed-through from the reverse side of the page.]*



*[Faint, illegible handwritten text, likely bleed-through from the reverse side of the page.]*

24. *Deva*  
30. *Alati*  
26. *Alati*  
27. *Alati*  
28. *Alati*  
29. *Alati*  
30. *Alati*  
31. *Alati*  
32. *Alati*  
33. *Alati*  
34. *Alati*  
35. *Alati*  
36. *Alati*



24. *Xanthemum annuum*  
In aridit montanis & collinis Monti Regali sat frequens
25. *Matricaria parthenium*  
In vineis Monti Regali frequentissima.
26. *Aster officinalis*  
In agro S. Albani Legit Cl. Bonelli.
27. — *montanus*  
In alpebus Garresei, & territoris Monti Regali.
28. — *odoratus*  
Bisula fol. amplexicaul. dentat. hirsutis. radical. ovatis. caul. in lan-  
ceolatis. caul. pauciflor.  
Circa S. Michele Monti Regali.
29. *Cineraria cordifolia*  
In montibus Garresei Villar Souran dicti.
30. — *longifolia*  
Supra Garressio
31. *Eupimedium alpinum*  
Gen. Nectaria 4. cyathiform. petalis incumbentia Calyce caducis  
4. phyllis corol. 4. petal. lappata acuminata 1. locul. 2. Valv.  
polytr.  
In montibus Garresei  
Caul. orbital. internod. juveniles ditiore fuscis pallidus. Fol. long. petiol.  
filiigata 3. ternata foliol. cordiform. dentat. Saepe virid. glabra  
venosa rigida Fl. purpureo flavo variegati emarginati.  
In montibus Garresei
32. *Draba nemorosa*  
In montibus Monti Regali frequens.
33. *Lunaria holymifolia*  
Alyssum fol. lanceolat. linear. acut. integerr. caul. prostratis. peren.  
In collibus editioribus Garresei.
34. *Thlaspi alliaceum*  
T. siliculis subovatis ventricosis fol. oblong. obtus. dentat. glabr.  
In vineis Circa Mondovi & Garressio.  
Planta allium redolens Caul. pedat. fol. infer. petiol. caul. in-  
terit. vel amplexicaul. fl. fasciculati subumbellati longe pedun-  
culati cor. alba.
35. *3bevi garreseiana*  
Supra Frabosa
36. — *jascaulis*  
3. Infrutices fol. lanceolat. linear. cavis, acut. integerr. ciliat.  
Inter Jasca alpinum Capotto Vicinarum.



37. *Cytisus lupinus*  
Inter Ormea & Gavresio  
c. flor. umbell. & termin. ram. decumb. foliol. ova t.
38. *Genista pilosa*  
G. fol. lanceolat. obtus. cauf. tuberculat. decumb.  
Locis arid. montium Monti Regalis.
39. *Vicia Gerardii*  
In sylvestribus subalpini Gavresii dicti i pra di fior.  
V. pedunculat. multiflor. petiol. polyphyll. foliol. villos. stipul.  
acut. integ. appendiculat.
40. *Artragalus depressus*  
In alpiibus di Caroso.
41. *Malva moschata*  
In apricis sylvi Gavresii et praesque loco dicto i pra di  
fior.
42. *Fragaria sterilis*  
in montibus Monti Regalis.
43. *Aquilegia virgata*  
in umbros. alpinis Gavresii dicti i pra di Gale Cl. Dama
44. *Helleborus hyemalis*.  
H. flor. folio incidente fl. luteo. / Aconitum unipolium bulbosum  
Copiosissime nascitur in vineis et collibus Monti Regalis i pra di.
45. *Dianthus glaucus*  
in collibus Montis Regalis.  
D. fl. solitarii aut subolit. squam. calyc. lanceol. quaterni  
brevibus coroll. & crenatis.
46. *Cucubalus italicus*  
c. calycibus clavatis, petalis semibipidis, panicula dichotoma  
erecta genitalibus declinatis, caule erecto.  
Copiose ad Jesser circa Vico di Mondovi.
47. *Agrostemma Coronaria*  
A. tomentosa fol. ovato-lanceolatis petalis emarginatis coro-  
llatis serratis.  
lychnis tomentosa Calyc. coccatis floribus longissime pedicatis  
Locis siccis Monti Regalis ubi circa loburent Montale & Traboto.
48. *Viola Ruppii*  
V. caule erecto multiflora fol. ovato-lanceol. serratis  
in montibus Montis Regalis.
49. *Cistus pilosus*  
C. suffrutescens stipulatus erectiusculis fol. linear. subul. circulatis  
incaut. calyc. laevibus.  
passim inter Gavresio & Ormea.



1. *Alnus incana* (DuRoi) Koch  
Shrub, 10-15 dm. tall. Bark smooth, grey. Leaves alternate, ovate, 5-10 cm long, 2-4 cm wide, serrated margins. Flowers in loose racemes. Fruit a winged samara.

2. *Alnus crispa* (DuRoi) Koch  
Shrub, 10-15 dm. tall. Bark smooth, grey. Leaves alternate, ovate, 5-10 cm long, 2-4 cm wide, serrated margins. Flowers in loose racemes. Fruit a winged samara.

3. *Alnus crispa* (DuRoi) Koch  
Shrub, 10-15 dm. tall. Bark smooth, grey. Leaves alternate, ovate, 5-10 cm long, 2-4 cm wide, serrated margins. Flowers in loose racemes. Fruit a winged samara.

4. *Alnus crispa* (DuRoi) Koch  
Shrub, 10-15 dm. tall. Bark smooth, grey. Leaves alternate, ovate, 5-10 cm long, 2-4 cm wide, serrated margins. Flowers in loose racemes. Fruit a winged samara.

5. *Alnus crispa* (DuRoi) Koch  
Shrub, 10-15 dm. tall. Bark smooth, grey. Leaves alternate, ovate, 5-10 cm long, 2-4 cm wide, serrated margins. Flowers in loose racemes. Fruit a winged samara.

6. *Alnus crispa* (DuRoi) Koch  
Shrub, 10-15 dm. tall. Bark smooth, grey. Leaves alternate, ovate, 5-10 cm long, 2-4 cm wide, serrated margins. Flowers in loose racemes. Fruit a winged samara.

7. *Alnus crispa* (DuRoi) Koch  
Shrub, 10-15 dm. tall. Bark smooth, grey. Leaves alternate, ovate, 5-10 cm long, 2-4 cm wide, serrated margins. Flowers in loose racemes. Fruit a winged samara.

8. *Alnus crispa* (DuRoi) Koch  
Shrub, 10-15 dm. tall. Bark smooth, grey. Leaves alternate, ovate, 5-10 cm long, 2-4 cm wide, serrated margins. Flowers in loose racemes. Fruit a winged samara.

9. *Alnus crispa* (DuRoi) Koch  
Shrub, 10-15 dm. tall. Bark smooth, grey. Leaves alternate, ovate, 5-10 cm long, 2-4 cm wide, serrated margins. Flowers in loose racemes. Fruit a winged samara.

10. *Alnus crispa* (DuRoi) Koch  
Shrub, 10-15 dm. tall. Bark smooth, grey. Leaves alternate, ovate, 5-10 cm long, 2-4 cm wide, serrated margins. Flowers in loose racemes. Fruit a winged samara.

11. *Alnus crispa* (DuRoi) Koch  
Shrub, 10-15 dm. tall. Bark smooth, grey. Leaves alternate, ovate, 5-10 cm long, 2-4 cm wide, serrated margins. Flowers in loose racemes. Fruit a winged samara.







50. *Sedum Cappa*  
*S. foliis planis caule ramo. floribus paniculatis.*  
*In ericetis umbratis in valle Tanari frequens.*
51. — *Galioides*  
*Similis G. cappa sed. nobis distinguitur non procumbens ut prior*  
*sed se erigit fol. inferior. 3. & 4. junct. deinde opposita prot.*  
*brima alterna vacui flor. etiam axillares sed proprios petal.*  
*purpurascunt et ariditate sunt teretifolia.*  
*Supra Garresio loco dicto il Garbo della Luna.*
52. *Ribes lva crispum*  
*R. ram. aculeat. bacc. glabr. pedicellis bractea monophylla*  
*in montibus Garresianis. Vidit Cl. Dana*
53. *Pyrus Cydonia*  
*ad septem in provincia Montiregali.*
54. *Orchis anthropophora*  
*ophrys bulbis subrotundis scapulo foliolo nectarii. labio lineari*  
*3 partito medio elongato bifido.*  
*Inter Garresium & Ormeam.*
55. *Fritillaria meleagris*  
*in pratis alpium Montiregalium.*
56. *Alnus europæum*  
*Vulgare in umbratis Montiregalis.*
57. *Buxus sempervirens*  
*Sylvatica facit inter Garresio & Ormea.*
58. *Arum Dracunculæ*  
*A. fol. pedat. foliolis lanceolatis integerr. superantibus spa-*  
*tham pedice breviorum.*  
*Circa Montaldo di Mondovi Ignatis Cholineri.*
59. *Agrostis miliacea*  
*A. petalo anteriore arista terminali recta stricta medioevi*  
*Circa Garresio et Ormea*
60. *Phromus inermis*  
*B. panicula erecta spiculis subteretibus subulatis nudis submutatis*  
*Circa Ormea lectus est.*
61. *Strophoptheris filicatum*  
*locus flumen Collatiana Montibus Regaleis.*
62. — *Spicant*  
*In montibus supra Roburent.*



- Auctarii.
63. *Campanula latifolia*  
Maxime frequent in pinguibus pratis di. Calotto.
  64. *Artemisia abrotanum*  
prope Ceva & Garreno nascitur.
  65. *Doronicum austriacum*  
locis pinguibus & humidis; secus torrentes in montibus di. Calotto,  
& Frabosa.
  66. *3. beris nana*  
3. herbacea fol. petiol. subrotund. dentatis flor. umbellatis.  
In alpinis della Marchiera Monti Regali.
  67. *Cardamine hirsuta*  
in sylvis ceduis subhumidis circa Casthus. di. Calotto paucim reperitur.
  68. *Arabis scabra*  
Crescit frequent in alpinis di. Mondovi ubi la Marchiera versus Casthus.
  69. *Turritis stricta*  
T. glabra lucens caule simplicissimo erecto silig. plani margine  
attenuatis.  
In pratis & pascuis subhumidis; supra Calotto.
  70. *Sisymbrium barbarea*  
S. fol. simpl. spatulato-ovatis; dentatis amplexicaulis nudis.  
In agro Monregalensi secus agros & locis siccis.
  71. *Papaver alpinum*  
Vulgare in agris Monregalensibus
  72. *Fumaria capreolata*  
Inter la Jrea & Rastello & prope Ormea.
  73. *Asplenium petraeum*  
M. Cant. nudo fol. gramineis involucri & involucellis distinct. polyphyllis.  
In rupibus supra Ormea.
  74. *Cherophyllum sylvestre*  
abunde locis umbratis & pinguibus secus Sylva prope Casthus. Calotto.
  75. *Dianthus deltoideus*  
D. flor. solitar. squam. calycin. lanceolat. binis corollis crenatis  
repens.  
In pratis montium Monti Regali.
  76. *Silene quadrifida*  
locis rupestribus ad fontes in alpinis Monregalensibus frequent.







*[Faint, illegible handwritten text, likely bleed-through from the reverse side of the page.]*

77. C...  
78. C...  
79. M...  
80. F...  
81. j...  
82. P...  
83. F...  
84. A...  
85. A...  
86. C...  
87. ...  
88. ...  
89. P...  
90. I...



77. *Cucubalus alpestris*  
*C. dichotomus* fol. linear. Calycibus pediformibus inflatis. Petal.  
 bispalatis.  
 In alpis di. Mondesi & supram fissuris locis umbris.
78. *Cirsium lunulatum*  
 In montibus supra Ormea in jugis la Rarchera.
79. *Allium nigrum*  
 Oritur et frequens est in alpis Montis Regalis.
80. *Fritillaria involucreta*  
 supra Ormea locis siccis & vespertinis mare spectantibus.
81. *Juncus filiformis*  
 passim locis profundis alpium Monregalensium.
82. *Phleum michelii*  
 locis aridis et siccis supra Ormea.
83. *Festuca heterophylla*  
 locis montanis siccis montium di. Mondesi.
84. *Avena festiva*  
*A. panicula* patente locis densis trifloris festivis floribus  
 duobus aristis.  
 In agris avenae Montis Regalis.
85. — *bromoides*  
*A. subspicata* spiculis binatis altera pedunculata arista divaricata ca-  
 lycibus octofloris.  
 In summis alpium Monregalensium soli expositis.
86. *Carex Bellardi*  
 locis alpestribus solique patentibus alpium Monregalensium versus  
 Carlin.
87. — *elongata*  
 in pratis spongiosis alpium di. Carotto.
88. — *carecens*  
*C. spiculis* subrotundis remotis sessilibus obtusis androgynis cap-  
 sulis ovatis obtusiusculis.  
 in pratis spongiosis supra Carotto.
89. *Polypodium limbo-permum*  
 in alpis Monregalensibus.
90. *Saxifraga lingulata* *Append: cit.*  
 frequens locis alpinis Monregalensibus.



91. *Boletus Frondosus*

St. sub caulis frondosus cespitosus sicut frondibus imbricatis planis  
lii reflexis poris albis.

ad candidos arborum vegetarum & hypnorum quandoque acquirit  
magnitudinem a Montegaleis. Barba. dicitur.

---

Cunei die 5. Junii 1800.

---



*Scutellaria*  
... ..  
... ..

*Scutellaria*  
... ..  
... ..

*Scutellaria*  
... ..  
... ..

*Scutellaria*  
... ..  
... ..

*Scutellaria*  
... ..  
... ..  
... ..  
... ..

*Scutellaria*  
... ..  
... ..  
... ..  
... ..

*Scutellaria*  
... ..  
... ..  
... ..  
... ..  
... ..

*Scutellaria*  
... ..  
... ..

*Scutellaria*  
... ..  
... ..

*Scutellaria*  
... ..  
... ..  
... ..



*[Faint, illegible handwriting at the top of the page]*

*[Faint, illegible handwriting in the upper middle section]*

*[Marginal notes on the right side of the page, including:]*  
Valeriana  
V. fol.  
3n mac  
V. fol.  
3n ma  
ce  
V. fol. in  
3n lig  
Rubia  
3n lig  
B. lutea  
J. fol. ob  
ex par  
videtur  
3n ar  
6. h  
L. fol.  
L. fol.  
pici in  
3n Ol  
V. lideni  
Geny G  
Pa lacin  
Micum  
J. herb  
ovato  
in Ag  
Mallubium  
in lacer  
Stachys an  
J. verticill  
Fragaria in  
av  
J. verticill.  
Unelia



*Plantae quae in Liguria Crescunt adnotante*  
*C. Allionio in sua Flora pedemontana.*

1. *Valeriana rubra.*  
*N. flor. monandr. caudat seu calcaratis foliis lanceolatis glaberrimis integerrimis.*  
*In maceris Ligurum.*
2. *\_\_\_\_\_ angustifolia*  
*N. fol. integr. longissimis & angustissimis flore calcarato donato.*  
*In maceris Ligurum ubi vocatur Covirella*
3. *\_\_\_\_\_ celtica*  
*N. fol. integerr. radical. ovatis caulinis linearibus obtus.*  
*In Liguria alpina sterilibus non procul a Savona urbe.*
4. *Rubia hirtorum*  
*In Liguria. etiam circa Uneliarum.*
5. *Salvia viridis*  
*S. fol. oblong. crenat. corollatum galea semioboiculata calycibus pa-*  
*chis reflexis bracteis cordato-acuminatis sensim decrescentibus*  
*viridibus in spicam tetragonam digestis.*  
*In arvis Uneliensibus*
6. *\_\_\_\_\_ hispanica.*  
~~*S. fol. ovatis crenato-dentatis verticillis subnatis stylo corollae ta-*~~  
~~*bro inferiore inaequilobato. S. fol. ovatis petiol. utriusque mucronatis*~~  
~~*spicis imbricatis calycibus hispida*~~  
*In Oliveis Uneliae.*
7. *Sideritis Romana*  
*Genus Calycis subspidus galea erecta plevumque bipartita barba hispida*  
*laciniis lateralibus acutis, media subrotunda stigma superius cylindricum*  
*inferiori membranaceo involutum.*  
*S. herbacea decumbens ebracteata calycibus spinosis labio superiori*  
*ovato.*  
*in Agvis Uneliae.*
8. *Marrubium vulgare*  
*in Sarcotis Liguria*
9. *Stachys annua*  
*S. verticill. 6. flor. fol. ovatis lanceolatis. hibern. levib. petiol. caule erecto.*  
*Frequens in Liguria*
10. *\_\_\_\_\_ avensis*  
*S. verticill. 6. flor. fol. obtus nudiusculis coroll. longitudine calyc. caul. debili.*  
*Uneliae secus arva.*



11. *Myosotis apula*  
M. coroll. bic. calyc. superant. semin. nud. latibus foliis lanceolatis hispido.  
racem. filicis  
In agro uneliensi locis sterilibus & secundaeva.
12. *Covis monipulensis* in Montibus Ligurum.
13. *Antirrhinum bellidifolium*  
A. fol. radical. lingulat. dentatis lineatis caulinis parvis integerrimis  
in Liguria Collibus Circa Oneliam
14. *Scrophularia Vernalis*  
Patrium in Liguria locis sylvestribus & umbratis
15. *Plantago psyllium*  
in aridis Sabulosis Liguria.
16. *Samolus Valerandi*  
ad iungua Circa Oneliam.
17. *Lythymachia tenella*  
L. fol. ovata acutiuscul. peduncul. foliis longioribus caule repente.  
unelias secus torrentes & ad fontium scaturigines.
18. *Hyoscyamus albus*  
H. foliis petiolatis sinuatis obtusis floribus sessilibus  
In Liguria
19. *Hyoscyamus aureus*  
H. fol. petiolatis dentatis acutis flor. peduncul. fructibus pendulis  
ad duos unelias.
20. *Verbascum phlomisoides*  
V. fol. ovata utriusque tomentosis inferioribus petiolatis ligulate bicollis  
in aridissimis & siccis Liguria.
21. *Verbascum sinuatum*  
V. fol. radical. pinnatifido-repandis tomentosis caulinis amplexicaulis  
nudiusculis ramis primito oppositis.  
ad Maris litora & Javona frequent.
22. *Convolvulus althaeoides*  
C. fol. cordat. sinuat. sericeis lobis repandis pedunculis bifloris  
in Liguria ad pagum Villatata.
23. *Convolvulus cheorum*  
C. fol. lanceol. tomentosis flor. umbellatis calycibus hirsutis caule erecto.  
Circa Oneliam.



col. b. light

Regardant

ate.

247

Feb. 6. 1891.

Electric

le evich.



12. *Carex monophylla* Gray. in *Stemmat. N. York.*  
 13. *Andropogon bell. difformis*  
 14. *Andropogon bell. difformis*  
 15. *Andropogon bell. difformis*  
 16. *Andropogon bell. difformis*  
 17. *Andropogon bell. difformis*  
 18. *Andropogon bell. difformis*  
 19. *Andropogon bell. difformis*  
 20. *Andropogon bell. difformis*  
 21. *Andropogon bell. difformis*  
 22. *Andropogon bell. difformis*  
 23. *Andropogon bell. difformis*  
 24. *Andropogon bell. difformis*  
 25. *Andropogon bell. difformis*  
 26. *Andropogon bell. difformis*  
 27. *Andropogon bell. difformis*  
 28. *Andropogon bell. difformis*  
 29. *Andropogon bell. difformis*  
 30. *Andropogon bell. difformis*  
 31. *Andropogon bell. difformis*  
 32. *Andropogon bell. difformis*  
 33. *Andropogon bell. difformis*  
 34. *Andropogon bell. difformis*  
 35. *Andropogon bell. difformis*  
 36. *Andropogon bell. difformis*  
 37. *Andropogon bell. difformis*  
 38. *Andropogon bell. difformis*  
 39. *Andropogon bell. difformis*  
 40. *Andropogon bell. difformis*  
 41. *Andropogon bell. difformis*  
 42. *Andropogon bell. difformis*  
 43. *Andropogon bell. difformis*  
 44. *Andropogon bell. difformis*  
 45. *Andropogon bell. difformis*  
 46. *Andropogon bell. difformis*  
 47. *Andropogon bell. difformis*  
 48. *Andropogon bell. difformis*  
 49. *Andropogon bell. difformis*  
 50. *Andropogon bell. difformis*  
 51. *Andropogon bell. difformis*  
 52. *Andropogon bell. difformis*  
 53. *Andropogon bell. difformis*  
 54. *Andropogon bell. difformis*  
 55. *Andropogon bell. difformis*  
 56. *Andropogon bell. difformis*  
 57. *Andropogon bell. difformis*  
 58. *Andropogon bell. difformis*  
 59. *Andropogon bell. difformis*  
 60. *Andropogon bell. difformis*  
 61. *Andropogon bell. difformis*  
 62. *Andropogon bell. difformis*  
 63. *Andropogon bell. difformis*  
 64. *Andropogon bell. difformis*  
 65. *Andropogon bell. difformis*  
 66. *Andropogon bell. difformis*  
 67. *Andropogon bell. difformis*  
 68. *Andropogon bell. difformis*  
 69. *Andropogon bell. difformis*  
 70. *Andropogon bell. difformis*  
 71. *Andropogon bell. difformis*  
 72. *Andropogon bell. difformis*  
 73. *Andropogon bell. difformis*  
 74. *Andropogon bell. difformis*  
 75. *Andropogon bell. difformis*  
 76. *Andropogon bell. difformis*  
 77. *Andropogon bell. difformis*  
 78. *Andropogon bell. difformis*  
 79. *Andropogon bell. difformis*  
 80. *Andropogon bell. difformis*  
 81. *Andropogon bell. difformis*  
 82. *Andropogon bell. difformis*  
 83. *Andropogon bell. difformis*  
 84. *Andropogon bell. difformis*  
 85. *Andropogon bell. difformis*  
 86. *Andropogon bell. difformis*  
 87. *Andropogon bell. difformis*  
 88. *Andropogon bell. difformis*  
 89. *Andropogon bell. difformis*  
 90. *Andropogon bell. difformis*  
 91. *Andropogon bell. difformis*  
 92. *Andropogon bell. difformis*  
 93. *Andropogon bell. difformis*  
 94. *Andropogon bell. difformis*  
 95. *Andropogon bell. difformis*  
 96. *Andropogon bell. difformis*  
 97. *Andropogon bell. difformis*  
 98. *Andropogon bell. difformis*  
 99. *Andropogon bell. difformis*  
 100. *Andropogon bell. difformis*

24. *Convolvulus*  
 C. fol. renif.  
 in marit.  
 25. *Erica Ciner.*  
 E. anther.  
 glomerat.  
 in mar.  
 26. ——— *Je.*  
 E. anther.  
 rev. pate.  
 Erica Al.  
 27. ———  
 E. anther.  
 papentini.  
 Zn Coll.  
 28. *Cotyledo.*  
 Gen. Caly.  
 mar. rectan.  
 C. fol. en.  
 in Ma.  
 Carl. semi.  
 alido sub p.  
 29. *Olea Europ.*  
 in lignis.  
 30. *Momordic.*  
 M. mom.  
 in lignis.  
 31. *Vitex ag.*  
 Gen. Did.  
 lab. super.  
 monophyll.  
 Zn Coll.  
 V. fol.  
 Arbor ro.  
 C. fol. ar.  
 32. *Globular.*  
 G. caule.  
 in mar.



24. *Convolvulus Solanella*  
C. fol. reniform. peduncul. uniflor.  
in maritimis Uneliae.

25. *Erica Cinerea*  
E. anther. (virid.) styloque inclusis corollis ovatis foliis ternis floribus  
glomeratis  
in montibus Uneliae.

26. ——— *Scorpiaria*  
E. anther. bicoloribus inclusis corollis campanulatis longioribus foliis  
ternis patentibus ramis albis.  
Circa Albenga.

27. ——— *arborescens*  
E. anther. bicolor. inclus. coroll. campanul. longioribus foliis quaternis  
patentissimis caule subarborescente tomentoso.  
In collibus maritimis. Liguria ut Circa Tagia.

28. *Cotyledon Umbilicus Veneris*. Decandria pentagynia  
Gen. Calyc. 5-fidus Corol. campaniform. limbo 4-vel 5-fido squa-  
ma nectariferis 5. ad basin germinis. Capsulae 5-fol. carnos. succulent.  
C. fol. cucullato-peltatis serratis-dentatis. caul. ramos. flor. erect.  
in Maceris Ligurum frequent.  
Caul. semiped. subnud. fol. altern. flor. terminal racemos. Corol.  
albido subflavesc. aut purpurea.

29. *Olea Europaea*  
in Liguria maritima.

30. *Momordica Elaterium*  
m. pomis hispida cirsidis nullis  
in Liguria calidiori.

31. *Vitex agnus Castus*  
Gen. Didynamia angiosperma Cal. 5-fidus corol. vixenti 2-labiata  
lab. super. emarginato infer. 3-fido fructus est bacca 4-locularis loculis  
monospermis.  
In collibus maritimis prope Albenga.  
V. fol. digit. serratis. typic. verticillatis  
Arbor ramis folisque subter incanis foliol. 5-7 lanceol. glabris.  
Corol. albo-purpurea vel tota alba.

32. *Globularia alypum*  
G. caule stricco fol. lanceol. 3. dentatis, integrisque  
in maritimis Uneliae.



33. ~~Glebularia~~ Xanthium spinosum  
X. spinis ternatis foliis trilobis.  
ad Charit. litus occurrit.
34. Echinops Sphaerocephalus.  
E. capitul. globos. fol. sinuato pinnatifidis.  
in Collibus liguaria non infrequens.
35. Carduus linchus  
Serratula fol. lyrate pinnatifidis pinnat terminali maxima flor-  
culis confinis.  
in Sylvis liguaria non infrequens est.
36. Cirsium medium X. p. Tenda.  
in montibus circa Unetiam.
37. Callina lanata  
C. caul. bifido calyc. sanguin. terminal. primo axillari sessili.  
ad maris litus locis siccis.
38. Centaurea Galachites  
C. calyc. setaceo-spinos. fol. decurrent. sinuato spinosis subter  
fomentis  
In Collibus Unetiae.
39. ——— elegant  
C. caul. simpl. fol. indivis. linear. axill. uniflor. & flor. terminal.  
in Vineis Unetiae.  
Varietas praecedentis. Alti Auct.
40. Artemisia abrotanum  
in pascuis & sterilibus locis collium Calidiorum liguaria.
41. ——— lobelii  
in Collibus liguaria non infrequens.
42. Chrysocoma linolyvis X. p. Monrey.  
in liguaria saepe nascitur.
43. Conyza sordida  
C. fol. linear. integerr. pedunc. long. 3 flor. caul. suffrutesco.  
ad rupes & veteros muros in Comitatu Unetensi.
44. Muphitalmum spinosum  
M. calyc. acute folios. fol. altern. lanceolat. amplexicaul. integer.  
caul. herbac.  
in liguaria parce invenitur.
45. ——— salicifolium  
M. calyc. ~~obtus. folios. pedunculat.~~ fol. altern. lanceolat. suberr. villor.  
herbac. Abunde in liguaria.



12/14

1941

1000

terminal.

o/o.

integer

6/err. Vill  
Caval.







46. *Achillea ligustica*  
in collinis supra oppidum: & Cairo.
47. *Aster acris*  
A. fol. lanceol. linear. strict. integerr. panic. floribus corymbosis  
paniculatis pedunculis foliosis.  
in praeputio Unelice.
48. — *bipennis*  
Bruta fol. oblong. decurrentibus denticulatis floribus corymbis ter-  
minalibus subsessilibus.  
in Liguria.
49. *Erigeron graveolens*  
l. ram. lateral. multiflor. fol. sublinear. integerr.  
in collibus maritimis Unelice.
50. *Senecio Sylvaticus*  
l. coroll. revolut. fol. pinnatif. denticulat. calic. corymb. erect.  
in Liguria.
51. *Cineraria maritima*  
c. flor. panicul. fol. pinnatif. tomentis. incan. sinuatis. caul. stratis  
ad Vitis Unelice.
52. *Catananche carulea*  
Gen. Cal. imbricatus. Receptac. paleaceum pappus aristatus  
calyculo 3-seto.  
C. squamis calyc. inferior. ovatis.  
abunde in sterilibus Liguria.  
Caul. 2. cubit. teret. subimpt. fol. altern. sessil. linear-lanceol.  
subdent. 3-nerv. pedunc. termin. l. flor. petal. coroll. erect.  
villos.
53. *Picris hibernica*  
Leontodon. fol. pinnatif. stratis. Calyc. acut. hirsut.  
in praeputio Unelice.
54. *Hieracium tomentosum* *Aurycula lanata* L.  
in Liguria.
55. *Crepis pulchra*.  
C. fol. sagitt. denticul. caul. panicul. flor. pyramid. glabris.  
in Vineis & Uliveti Unelice passim.
56. *Senecio tenuifolius*  
abunde in Liguria.
57. *Scorzonera graminifolia*  
l. fol. linear. ensiform. integerr. cavinatis  
secus agros Unelice.



58. *Tragopogon picroides*  
*T. calycibus monophyllis; corollis brevioribus aculeat. fol. runcinat.*  
*denticulatis.*  
*in Olivetis Uneliae.*
59. *Epimedium alpinum* V. plant. Monregal.  
*in Liguria.*
60. *Alyrium maritimum.*  
*Clypeola perennis silicul. bifocular. ovat. disperm. fol. linear.*  
*in arenosis maritimis Uneliae.*
61. *Linaria halmifolia* V. pl. Valder.  
*in collibus Uneliae editionibus non tamen ita quicquid sunt proceri.*
62. *Sinapis maritima* V. pl. Vinad.  
*Nascentur ad ora maritima Uneliae.*
63. *Hyperic. lychnitis*  
*H. caul. subhispido erecto valis latius. Aliguae.*  
*in Valle Uneliae.*
64. *Ruta graveolens*  
*in principatu Uneliae. Varietas minor frequentior est.*
65. *Euphorbia spinosa*  
*E. umbella subsida simplici involuelli ovatis primariis bi-*  
*phyllis fol. oblong. integerr. caul. pubescent.*  
*ad Ruper in Liguria nascitur.*
66. ——— *erythymoides*  
*E. umbella trifida, bifida involuelli. ovatis fol. lanceolatis. obtus. subhy-*  
*villos.*  
*Locis incultis & ad ora sylvarum Liguria non infrequens.*
67. ——— *dendroides.*  
*E. umbella multifida dichotoma involuelli subcordatis primariis*  
*triphyllis caule arboresc.*  
*ad scopulos mari proximos prope Uneliam & Alarciam.*
68. *Papaver hybridum*  
*sp. caputis subglobosis torosis hispidis caul. folior. multiflor.*  
*inter segetes. Uneliae.*
69. *Trifolium ochroleucum.*  
*T. caul. hirsut. scabr. fol. integerr. moll. spic. subulosis. ochroleuc.*  
*Locis apricis collibus, Liguria, frequent est.*
70. *Lobus edulis*  
*L. leguminibus subrotatis gibbis incurvis*  
*in pascuis sterilibus Uneliae.*







58. *Tragopogon microides*

*T. calycibus monophyllis; corollis brevioribus aculeatis. fol. runcinatis.*

11. 10/11

59

60

61

62

63

64

65

66

67

68

69

70



оп. *Colus ornithopodoides*.

*Colus ornithopodoides* a 116 2.0

h/.

*in Liguria humilis? planta.*



58. *Tragopogon picroides*

*T. c. l. lobul. monothyllis corollis brevioribus aculeat. fol. runcinat.*

59

60

61

62

63

64

65

66

67

68

69

70

71. *Lobul.*  
*l. legu.*  
*in l.*

72. *Dory.*  
*in l.*

73. *Medi.*  
*Ch. pa.*  
*3h. 1.*

74. *Onon.*  
*in l.*

75. *Cyhir.*  
*pa.*

76. —  
*in l.*

77. *Hedy.*  
*H. fol.*  
*3nag.*

78. *Onogry.*  
*Hedi.*  
*alisbre*  
*in l.*

79. *Vicia*  
*V. leg.*  
*vibur*  
*in l.*

80. —  
*in d.*

81. *Onithea*  
*O. fol.*  
*ad ag.*

82. *Sparg.*

83. —  
*J. fol.*  
*in*

84. *Cercis.*

85. *Athyra.*



71. *Lolium ornithopodioides*.  
L. legum. subternat. arcuat. compress. caulibus diffusis  
in Liguria.
72. *Dorychnium Monpelienicum*  
in Liguria abunde.
73. *Medicago marina*  
M. pedunc. racemos. legum. Cochl. spinos. caul. prostratis. fomentis.  
In sterilibus litoreis maris proximis.
74. *Ononis Columna*.  
in Liguria frequenter obviam venit.
75. *Cytisus densifolius*  
passim in Liguria.
76. ——— *argenteus*  
in collibus Liguriae.
77. *Hedysarum spinosissimum*  
H. fol. pinnat. legum. articulat. aculeat. fomentis. caul. diffus.  
In agro unelienis locis sabulosis maris proximis & secus arva.
78. *Onobrychis sarcocolla*.  
Hedys. fol. pinnat. legum. i. specim. sulcat. rubic. corollae  
altissimis, scapis subradicatis.  
in sacris apricis unelias.
79. *Vicia Seppium*  
V. legum. pedicell. subquatern. erect. foliol. ovata. integerr. escheno-  
ribus decrescentibus  
in editionibus Liguriae montibus A. Tana.
80. ——— *harbonensis*  
in dumetis Liguriae.
81. *Ornithopodium scorpioides*  
O. fol. ternatis subterribus impari maximo.  
ad agros macilentos in Liguria.
82. *Spartium juncum*  
in Liguria.
83. ——— *spinosum*  
S. fol. ternatis ramis angulatis spinosis  
in Olivetis unelias.
84. *Cercis siliquastrum*  
in Liguria maritima
85. *Athyrium moussetianum*  
in Liguria humilis planta.



96. *Eryngium maritimum*  
l. fol. radical. subrotund. plicat. spinos. capitul. pedunculat.  
paleis micropidatis  
in sabulo ad maris litus.

97. *Platinaca oppositifolia*  
p. fol. bipinnatis foliol. oblique cordatis petalis paleaceo rigori.  
Unelia locis umbrati & montanis sylvestribus.

98. *Ferula nodiflora*  
Gen. Semen magnum oblongo-ovatum 5. striatum Cinchumata magna  
solida non membranacea.  
F. foliol. appendiculatis umbellis sessilibus  
ad vineas in collibus Unelia.

99. *Pencedanum officinale*  
p. foliol. hieraciparitis filiformibus linearibus  
in montibus liguria.

90. *Anerthum graveolens* fructibus compressis  
ad maris litus prope Uneliam.

91. *Angelica Levithicum*  
Liqueticum fol. multiplic. foliol. superm. incis. flor. luteolis  
inter savonum et Genuum spontaneam habent liquet ejusque caule  
& folia edunt inter Apii graveolentis.

92. *Coriandrum terculatum* fructibus didymis  
passim in agris Unelia

93. *Oenanthe spinipinelloides*  
Gen. fl. radiati disco sterili petala inaequalia fructus ovatus calyce  
inligui 5. dentato persistente coronatus sem. ovat. <sup>non</sup> acuminat.  
thialum inter planum  
O. foliol. radicalibus cuneatis filis caulin. integr. linear. longioribus  
canaliculatis.  
In liguria praesens occurrit.

94. *Bupleurum rotundifolium*.  
p. involucl. univers. nulli fol. persiliat.  
In agris collium calidiorum liguria passim speciatim Unelia.

95. ———— <sup>juncum</sup>  
B. caule erecto paniculato fol. linear. involucl. 3phyll. involucl. 5phyll.  
Juncus agris in liguria.

96. *Faniculum officinale*  
in Provincia Unelia.

97. *Bunium minus*.  
p. fol. uniformibus involucl. polyphyll. fructibus subcylindricis apice crot-  
sioribus stylis reflexis deciduis  
In edioribus montibus liguria.



98. *Crithmum maritimum*

in lat. maritimo lat. luvulicis

folia lanceolata, serrata, margine  
denticulata, nervis distinctis

caulis erectus, teres, hirsutus, apice  
capitulo terminali, foliis oppositis

fructus baccatus, globosus,  
seminibus distinctis, foliis oppositis

caulis erectus, teres, hirsutus, apice  
capitulo terminali, foliis oppositis  
fructus baccatus, globosus, seminibus  
distinctis, foliis oppositis

caulis erectus, teres, hirsutus, apice  
capitulo terminali, foliis oppositis

fructus baccatus, globosus, seminibus  
distinctis, foliis oppositis

caulis erectus, teres, hirsutus, apice  
capitulo terminali, foliis oppositis

fructus baccatus, globosus, seminibus  
distinctis, foliis oppositis

caulis erectus, teres, hirsutus, apice  
capitulo terminali, foliis oppositis

caulis erectus, teres, hirsutus, apice  
capitulo terminali, foliis oppositis  
fructus baccatus, globosus, seminibus  
distinctis, foliis oppositis

caulis erectus, teres, hirsutus, apice  
capitulo terminali, foliis oppositis

caulis erectus, teres, hirsutus, apice  
capitulo terminali, foliis oppositis



98. *Crithmum* *Gen. Flo.*  
Coharent  
C. fol. la  
ad Rupa

99. *Caucalis*  
Scandix de  
in vineis

100. *Ronunculus*  
R. semini  
in arvis

101. *Corrigiola*  
Gen. Cal. 5 p  
Loc. aren  
Paul. pro  
Hyp. bin  
Congesti in

102. *Potentilla*

103. *Helioscopia*  
in

104. *Silene* *Servic*  
J. fol. acut  
ad Mavir

105. *Oxy* *cornic*  
in mo

106. *Reseda* *lu*  
in

107. *Tribulus*  
Cal. pro  
pithum C  
J. fol. sub  
in liguri  
Cant. pedat  
nata foliolis  
lavy falcicu

108. *Quonimus*  
e. fol. acut  
in Odibov

109. *Cirsium* *albid*  
C. arboresce  
fertilibus  
in incultis



98. *Crithmum maritimum*  
 Gen. Flores aequales fertiles. Semen ovale intus planum tota superficie  
 cohaerens dorso tribus obrenis costis percussio.  
 C. fol. lanceolat. carnosis.  
 ad Ruper mari proximis in muris Vetusli Uneliae.
99. *Caucalis nodosa*  
 Scandix seminibus subcylindricis hispida; caule hispido genicul. humidi;  
 in vineis & sterilibus figuris minime varia.
100. *Ranunculus parviflorus*  
 R. seminibus mucatis fol. simpl. laciniat. acut. hirsut. Caule diffuso  
 in arvis subhumidis Uneliae.
101. *Corrigiola litoralis*  
 Gen. Cal. 5 phyll. Stamina 5. petal. 5. Sem. 1. Wignebr. nud. calyc. content.  
 Locis arenosis mari proximis.  
 Caul. prostr. vix palin. Ramosi fol. altern. sessil. lanceol. integer.  
 bipul. binat. acutae argenteae. Flores racemos. in ultimis ramis  
 congesti interd. axillares petal. Oval. alba.
102. *Potentilla recta*  
 in Liguria valde frequens.
103. ——— rubens  
 in Calidioribus collibus Liguria.
104. *Helleborus hyemalis* V. pl. Monveg.  
 in vineis et collibus Liguria.
105. *Silene sericea*  
 S. fol. acuminat. Succulent. hirsut. lucid. flor. carnea  
 ad Mavis lites in Sabatoris inter Oneglia & porto Maurizio.
106. *Oxy corniculata*  
 in montibus Liguria ad iurigna frequens.
107. *Aerida luteola*  
 in Liguria ad vias muros & rudrata.
108. *Tribulus terrestris* Decand. monogyn.  
 Cal. profunde 5-fidus petal. 5. aequal. stili nullus. Stygma la-  
 pitulum capsulae plures congestae aculeatae.  
 T. fol. sub 6-jugis subaequalibus seminibus quadrangulis  
 in Liguria  
 Caul. pedat. & ultra procumb. Ramos. villosus rubens. fol. abrupte pin-  
 nata foliolis oblique lanceolatis ciliatis subtus villor. Flores axil-  
 lares fasciculati praeduncat. flavi.
109. *Quonimus latifolius*  
 e. fol. acute & argute serratis capsul. angul. acut. membranaceis  
 in edithodibus montibus Liguria.
110. *Cistus albidus*  
 C. arborescens exstipulatus fol. ovato lanceolatis tomentosis incanis  
 sessilibus subminuor.  
 in incultis & maritimis collibus Uneliae.



111. *Cistus gultatus*.  
*C. herbaceus erectissimus fol. oppos. lanceolatis. trineuro. racem. erect.*  
*C. flor. pallid. macula punicea in angulo*  
*in pascuis dicitur sterilibus Liguria.*
112. — *Pilosus*  
*in Ligurum montibus.*
113. — *Vosens*  
*Unelia nalcitur.*  
*a C. helianth. differt. caul. magis ramos. & fruticosa. fol. firmior.*  
*subtus subincanis corollis roseis.*
114. *Linum hirsutum*  
*L. calyc. hirsuti, acuminati, dent. altern. fol. alterni, variet. oppositi.*  
*non varum in Liguria.*
115. *Jedum Cyprea*  
*J. fol. plani, caule ramoso flor. paniculatis.*  
*in collibus Liguria.*
116. — *acre*  
*in dicitur pascuis Liguria.*
117. *Rhamnus infectoria.*  
*R. spinis terminalibus flor. 4-fidis dioici, caulibus procumbent.*  
*in Liguria lectus est.*
118. — *Lycioides*  
*R. spinis terminalibus foliis linearibus*  
*in Liguria habuit Cl. Alton.*
119. *Rhus cotinus*  
*frequent in montibus Liguria.*
120. *Prunus domestica*  
*pr. pedunculis solitariis fol. lanceolat-ovatis. convolut. serrat. ram. multi.*  
*in Sylvis Liguria.*
121. *Myrtus Communis*  
*Glen. calyce superius 5. vel 4. fidis petalis. vel 4. ovata stamina*  
*plura calyce inserta stigma obtusum bacca olivaceo. coronata*  
*umbilicata 2 vel 3. loculari.*  
*Myrtus minor Vulgaris*  
*savona reperitur.*  
*Planta pubesca fol. busco acutiora opposita subsessilia ovata-acu-*  
*minata laeva satore viridia odore petala alba.*
122. *Azota rubiginosa*  
*A. germ. globosis petiol. aculeis. aculeis recurvis fol. subt. rubiginosis.*  
*in collibus Liguria ad Vias, & sepes occurrunt.*
123. *Pyrus Cydonia*  
*ad saepe in Liguria.*







124. *Orchis*  
 in pascuis siccis, sub montibus  
 in pascuis siccis, sub montibus
125. *Orchis*  
 in pascuis siccis, sub montibus  
 in pascuis siccis, sub montibus
126. *Orchis*  
 in pascuis siccis, sub montibus  
 in pascuis siccis, sub montibus
127. *Orchis*  
 in pascuis siccis, sub montibus  
 in pascuis siccis, sub montibus
128. *Orchis*  
 in pascuis siccis, sub montibus  
 in pascuis siccis, sub montibus
129. *Orchis*  
 in pascuis siccis, sub montibus  
 in pascuis siccis, sub montibus
130. *Orchis*  
 in pascuis siccis, sub montibus  
 in pascuis siccis, sub montibus
131. *Orchis*  
 in pascuis siccis, sub montibus  
 in pascuis siccis, sub montibus
132. *Orchis*  
 in pascuis siccis, sub montibus  
 in pascuis siccis, sub montibus
133. *Orchis*  
 in pascuis siccis, sub montibus  
 in pascuis siccis, sub montibus

124. *Orchis*  
 in pascuis siccis, sub montibus
125. *Orchis*  
 in pascuis siccis, sub montibus
126. *Orchis*  
 in pascuis siccis, sub montibus
127. *Orchis*  
 in pascuis siccis, sub montibus
128. *Orchis*  
 in pascuis siccis, sub montibus
129. *Orchis*  
 in pascuis siccis, sub montibus
130. *Orchis*  
 in pascuis siccis, sub montibus
131. *Orchis*  
 in pascuis siccis, sub montibus
132. *Orchis*  
 in pascuis siccis, sub montibus
133. *Orchis*  
 in pascuis siccis, sub montibus



124. *Orchis Coriophora*  
*O. bulb. indivis. nectar. lab. 3. fid. reflexe. crenat. corum brevi.*  
*petal. connivent.*  
*in prati/ siccioribus collium Liguriae non infrequens.*
125. — *hircina*  
*O. rad. subrotund. label. longiss. 3. paulito plicato.*  
*Sylvos. et apricis locis collium Liguriae. sed non frequens.*
126. — *muscaria*  
*O. rad. subrot. label. sericeo bitulro brachiol. brev. subulat.*  
*in sterilibus graminosis collium Liguriae.*
127. — *avachnites.*  
*O. rad. subrotund. label. holosericeo emargin. appendiculato.*  
*abunde in collibus Liguriae quandoque compressed. (verit.)*
128. — *lingua*  
*several bulbis subrotundis nectaris lab. 3. fido acuminato petalis*  
*longiore glabro.*  
*in prati & pratis subhumidis Liguriae frequens.*
129. — *anthropophora* V. pl. Monvegal.  
*in dumetis umbrosis unctis quandoque occurrat frequenter ad*  
*oppidum monte d'Alto.*
130. — *fusca*  
*O. rad. subrotund. spica long. label. 4. fido brachiol. angust. ciliolatis labis*  
*several.*  
*In collibus Ligurum nascitur.*
131. *Epipactis aethiva.*  
*O. bulbis fasciculatis filiform. nectar. lab. ovat. integerrimo caul. apophyll.*  
*in dumetis et sterilibus Calidivum collium Liguriae.*
132. *Pancreaticum maritimum.*  
*Gen. Calycis est spatula Nectarinum in pediculis formae 12. fid.*  
*fructu impositum stamina sustinent stamina 6. & 6. fid. petala.*  
*sp. spatula multiflora petalis planis filis lingulatis.*  
*ad Vesper prope litus maris ubique frequens*  
*Fl. magni petal. lanceol. alba sub hyl. linea viridi.*
133. *Allium roseum*  
*A. caul. planifol. umbellifero umbella fastiginta, petalis emar-*  
*ginatis stamin. breviss. simplicibus.*  
*In Olychis et arvis unctis.*
134. *Asphodelus fistulosus*  
*A. caul. nudo. fol. strichis subulatis striatis subfistulosis.*  
*in Liguriae montibus saeva proxima*



135. *Scilla maritima*.  
*S. nudiflora* bracteis reflexis  
 ad maris litem arenosa & suppet inhabitat.
136. *Asparagus acutifolius*  
*A. caule inermi angulato fruticoso fol. acinis vix rigidiis perennantibus mucronatis aequalibus*  
 in principatu Unelica.
137. *Lythrum hyssopifolia*.  
*L. fol. alternis linearibus fl. descandis*  
 in humidis & salubris liguia.
138. *Adonis flammea*.  
*A. fl. octopetalis fructibus subcylindricis calyc. hirsutis*  
 in collibus calidioribus liguia.  
 Floret arate petala flammea macula fere nulla.
139. *Pinus sylvestris*  
 in levibus montibus & collibus liguia.
140. *Juniperus oxycedrus*  
*J. fol. ternis patentibus mucronatis bacca brevioribus.*  
 in sylvis appicis versus Uneliam.
141. ——— *phænicea*  
*J. foliis ternis oblitteratis imbricatis obtusis.*  
 frequens in collibus Unelica.
142. *Atriplex verea*.  
*A. caule herbaceo fol. incan. serratis fruct. 4. angul. dentato vereo*  
 compresso.  
 In liguia frequens est.
143. *Hedysarum viviparum*  
 in liguia frequens.
144. *Salsola Kali*  
*Calyc. profunde 5-fidus quodammodo coloratus pedicellus stylus lim-  
 pleris semibifidus thamina 5. Capsula globosa semen. 1. cochlearine*  
*J. decumbens herbacea fol. subulata spinol. scabris calycibus*  
*marginalis arcuatis.*  
 in maritimis prope Uneliam.  
 Planta spinosa fol. crassa calyc. cinch. margine foliaceo stylus 3fid.
145. *Juncus acutus*  
*J. culmo subundo teret. mucronato terminali involucri 2-phyllo*  
 spinoso.  
 in locis maritimis Unelica.
146. *Buncus sempervirens*  
 in montibus liguia ad suppet.



na figura do livro de...



*[Faint handwritten botanical notes, likely bleed-through from the reverse side of the page. The text is mostly illegible due to fading.]*

in molibus figura ad vasa.

147. *Ruscus*  
*R. fol.*  
 in col.  
 148. — *R. fol.*  
 in Jarcon.  
 149. *Vitcum*  
*Gen. Ch.*  
*Calyci a*  
*globosa*  
*V. fol.*  
*Unelia*  
*Fol. opp*  
*albo-vu*  
 150. *Atrum a*  
*A. a cov*  
 in co.  
 151. *Nardus an*  
*N. calyc.*  
 ad  
 152. *Alopecurus*  
*A. culmo*  
 in pua.  
 153. *Rottbolla*  
*Gen. locusta*  
*sed glumif*  
*lig. lumif.*  
*N. spica tere*  
*Unelia*  
 154. *Lolium tenu*  
*L. spica mu*  
 in pascuis.  
 155. *Aira Canesce*  
*A. fol. seta*  
 in tabulo.  
 156. *Poa rigida*  
*P. panicula*  
 abunde in.  
 157. *Bromus ligu*  
*Gramen Brom*  
 in liguria a



147. *Ruscus aculeatus*  
R. fol. supra florifer. nudis  
in collibus Liguriae.
148. — *hypoglossum*  
R. fol. supra florifer. sub foliato.  
in saxosis incultis Liguriae.
149. *Vitcum album*. Dioecia tetrandria.  
Gen. char. Cal. 4. part. segment. ovat. Coroll. 0. filam. 0. antherae  
calyci adnatae. Fam. Cal. 4. phyll. superius Coroll. 0. Styl. 0. Bacca  
globosa glutine plena 1. sperma semina cordata compressa.  
V. fol. lanceolatis obtusis caule dichotomo spicis axillaribus.  
Umelia et locis maritimis frequens et parasitica planta  
Fol. opposita sessilia integerrima crassa dura sub lente. Baccae  
albo-rubrae nitidae.
150. *Atrum avicatum*.  
A. acaule fol. cordat. oblongis spathae apertura ovata.  
in collibus Umelia.
151. *Nardus arvensis*  
N. calyc. arvensis.  
ad Vini in Liguria.
152. *Alopecurus bulbosus*  
A. culmo erecto spica cylindrica radice bulbosa  
in pratis latis Liguriae.
153. *Rottbolla incurvata*.  
Gen. locustae uniflorae flexuosi caulis scrobibus appressis. Calyce 2. glumis  
sed glumis junctis se positis ita ut 1. glumis calyce videatur. folliculis  
biglumis.  
A. spica tereti subulata gluma calycina subulata appressa 2. partita.  
Umelia in agris inveni proceris.
154. *Lolium tenue*.  
L. spica multica tereti spiculis trifloris  
in pascuis Liguriae.
155. *Aira canescens*.  
A. fol. setaceis summo spathaceo panicul. inferne obsovente.  
in sabulis sterilibus Liguriae.
156. *Poa rigida*  
P. panicula lanceolata subramosa secunda ramul. altern. secundis.  
abunde in Liguria.
157. *Bromus Ligusticus*  
Gramen bromoideae festucae tenuique panicula ardens  
in Liguria ad litora maris.



158. *Elymus arenarius*  
L. Spica erecta aerea calyc. tomentos. florulo longioribus  
in arena ad maritima.

159. *Agropyron ovata*  
L. Spica ovata calyc. omn. brevissimis  
locis siccis in Liguria ubique

160. *Andropogon hirtum*  
L. Panícula spici. conjugatis calyc. hirsutis.  
Frequens in lina locis aridis & pascuis.

161. *Scirpus mucronatus*  
L. Culmo triangulo nudo acuminat. spic. conglomer. sessil. lateralibus.  
in maritimis lina.

162. *Schoenus mucronatus*  
L. culm. teret. nud. spicul. ovat. fasciculat. involucri sub 6. phyllo fol.  
canaliculatis  
In maritimis lina.

163. *Adiantum Capillus Veneris*  
L. fol. decompositis foliol. altern. pinn. ~~diversis~~ cuneiformibus  
lobatis ped. cellatis.  
Vulgatum in pratis & rupibus Liguria Colliumq. maritimorum.

164. *Fucus serratus*  
Charc. vesiculae villis intertextae Form. vesiculae ad-  
spere granis immixtis apice prominulis semina soli-  
taria substantia coriacea vesiculatis.  
F. fronde plana dichotoma serrato dentata fructificatio-  
nibus terminalibus tuberculosis  
In mari lina.

165. ——— *vesiculosus*  
F. fronde plana dichotoma integerrima vesiculis  
axillaribus geminis. terminalibus tuberculatis  
in mari lina.

166. ——— *nodosus*  
F. fronde compressa dichotoma fol. dich. integerr.  
vesiculis immixtis solitariis dilatatis  
ad mare lina.

167. ——— *selaginoides*  
F. frond. filiform. ram. dichot. subulatis  
alternis basi vesiculosis  
ad lina lina.



*Phaeophyta*  
F. frons p. l. ...  
foliolum ...  
in ...

*abundant*  
F. ...  
in ...

*abundant*  
F. ...  
in ...

*abundant*  
F. ...  
in ...

*abundant*  
F. ...  
in ...

*abundant*  
F. ...  
in ...

*abundant*  
F. ...  
in ...

*abundant*  
F. ...  
in ...

*abundant*  
F. ...  
in ...

*abundant*  
F. ...  
in ...

*abundant*  
F. ...  
in ...

*abundant*  
F. ...  
in ...

in mari unctis.



168. *Fucus* *serotinus*  
F. *serotinus* (L.) Lamour.  
Common in shallow water.

169. *Fucus* *vesiculosus*  
F. *vesiculosus* (L.) Lamour.  
Common in shallow water.

170. *Fucus* *granulosus*  
F. *granulosus* (L.) Lamour.  
Common in shallow water.

171. *Fucus* *maritimus*  
F. *maritimus* (L.) Lamour.  
Common in shallow water.

172. *Fucus* *racematus*  
F. *racematus* (L.) Lamour.  
Common in shallow water.

173. *Fucus* *littoralis*  
F. *littoralis* (L.) Lamour.  
Common in shallow water.

174. *Fucus* *serotinus*  
F. *serotinus* (L.) Lamour.  
Common in shallow water.

175. *Fucus* *vesiculosus*  
F. *vesiculosus* (L.) Lamour.  
Common in shallow water.

176. *Fucus* *granulosus*  
F. *granulosus* (L.) Lamour.  
Common in shallow water.

177. *Fucus* *maritimus*  
F. *maritimus* (L.) Lamour.  
Common in shallow water.

68. *Fucus*  
169. —  
170. —  
171. —  
172. —  
173. —  
174. —  
175. —  
176. *Ulva*  
177. —  
178. —  
179. —  
180. —  
181. —  
182. —  
183. —  
184. —  
185. —  
186. —  
187. —  
188. —  
189. —  
190. —  
191. —  
192. —  
193. —  
194. —  
195. —  
196. —  
197. —  
198. —  
199. —  
200. —



168. *Fucus fasciculaceus*

*F. fronde filiformi ramorissima vericulis ovali terminati  
foliolis multipartitis obtusis apice fructificantibus  
in Rupibus litor. & unelia.*

169. — *abrotanoides.*

*F. fronde ramosa capillari multipida fasciculata vericulis glo-  
boris.  
in mari unelia.*

170. — *acinarius.*

*F. caule filiformi ramis fol. linear. integerr. fructificat. globosis  
pedunculatis  
in mari unelia.*

171. — *volundus.*

*F. fere subdichotomus unipolus ramis obtusis verrucosis  
in mari unelia.*

172. — *muscoideus.*

*F. fronde fere ramorissima ramis sparsis spinis mollibus alternis.  
in mari unelia.*

173. — *longissimus.*

*F. fere rubens minus ramosus in longum protensus  
in mari unelia.*

174. — *confervoides.*

*F. subdichotomus fere filiformis ramoriss. ram. in aq. setaceis.  
in mari unelia.*

175. — *plumosus.*

*F. caule filiformi subcompresso ramoso frondibus lanceolatis  
bipinnatis plumosis (purpureo) coloratis  
in mari unelia.*

176. *Ulva intestinalis.*

*Gen. Fructific. in substant. membranacea longa tubulosa  
vel Campanulata larca diaphana.  
U. tubulosa simpliciter intestinorum figura  
in mari unelia.*

177. — *compressa.*

*U. tubulosa ramosa compressa  
in mari unelia.*

178. — *lactuca.*

*U. palmata prolifera membranacea ramentis infern. angustatis  
Fucus fronde imbricata frond. parvialibus erectiusculis sinua-  
to laciniatis obtusis lactuca similis. & planta.  
in mari unelia.*

179. — *linga.*

*U. fronde oblonga bullata  
in mari unelia.*



180. *Lycoperdon tuber.*  
in collibus liguriae.

181. *Aristolochia Clematidis*  
in arvis subhumidis, liguriae.

182. *Statice lappacea*  
St. caepodichotomo ancipiti fol. ovat. mucronat. floribus  
aggregatis.  
Limonium fol. lanceolat. mucronat. radicalibus caul. ancipiti  
dichotomo stipulis simplicibus.  
Occurrit circa Ventimiglia

183. *Artemisia camphorata*  
A. fol. petiol. palmato-multifidi, linearibus superveni simplicibus  
calycibus angulosis receptaculo hirsuto.  
Nascitur frequenter in Liguria locis superstitibus.

---

Cunei die 6. Junii 1800.

Hodie ab exercitu Caesareo in Cunei celebrata est deditio  
Genuae urbis principis Liguriae.

---







192. *Spizella socialis*  
in *Spizella socialis* *Spizella*

193. *Spizella socialis*  
in *Spizella socialis* *Spizella*  
in *Spizella socialis* *Spizella*  
in *Spizella socialis* *Spizella*  
in *Spizella socialis* *Spizella*

194. *Spizella socialis*  
in *Spizella socialis* *Spizella*  
in *Spizella socialis* *Spizella*  
in *Spizella socialis* *Spizella*

---

Cum die 6. Junii 1800

Hodie at tunc in *Spizella* *Spizella*  
in *Spizella socialis* *Spizella*

---








































1. Valerian  
/ sea  
Cipri

2. —  
V. ma  
in v  
Caul

3. —  
in G

4. —  
V.  
lan  
in  
Plant  
Jule  
Cune

5. — Co  
V. tan  
Dealt  
3

6. Galium /  
G. fol.  
Nico

(\*) Mo  
in /ua  
nat facit  
et alibi  
velicentur  
familiar  
vunt, et  
Cune eni  
parava ne  
propiciscar



# Plantæ Nicæenses, & Comitatus quarum locus in Flora ped. natalis designatur. (\*)

1. *Valeriana rubra*. V. pl. ligur.  
frequens ad muros. in vineis agri Nicæensis.  
Circa Jospello ubi appellatur *Cornaccin*.
2. ——— *Calcitrapa*.  
V. monostemon fol. im. integr. Superioribus pinnatis  
in umbrosis agri Nicæensis.  
Caulis cubitalis ramis folia humifusa corolla dilute rubra.
3. ——— *tuberosa*. V. pp. Monnegal.  
in Comitatu Nicæensi speciatim supra *Coua* vasa.
4. ——— *Jascanilis*  
V. fluv. triandri fol. subdentatis radicalibus ovatis caulinis linearilanceolatis.  
in montibus Nicæensibus.  
Planta nitida Caul. spitham. simpl. fol. subternis. ad ora obsolete pilosa radical. 4. caulin. 2. opposita ramis brachiatis. p. dunc. opposit. fl. panicul. corol. alba.
5. ——— *Coronata*  
V. caul. dichotomo fol. pinnatifidis sem. Corona radiatum dentata dentibus 6.  
In arvis Nicæensibus.
6. *Galium palustre*  
G. fol. quatern. obovat. inæq. caul. diffus.  
Nicæe semper pascuis & humida loca.

---

(\*) *Notandum*. Non omnes stirpes quas Celeber. Aldone in sua Flora pedemontana Comitatus Nicæensis indigenas facit hic recensentur. Nimirum enim liber increveret, et alibi descriptæ plantæ denuo enumerarentur. Ideoque relincentur Stirpes eæ, quæ in pedemontio frequentes, et familiares sunt, vel in hortis culta omnibus innotuerunt, & eæ quoque, quas ego vel in agro Patrio, vel Cuneensi jam observavi, quasque mihi velim comparare necesse non est, ut ad Comitatum Nicæensem propiscar.

---



17. *Galium uliginosum*

G. fol. dens lanceolatis retrorsum serratis aculeatis mucronatis rigidis co-  
rollis fructu majoribus.  
in agro Nicæensi ad vivat.

8. ——— *Jascale.*

G. fol. dens obovatis obtusis caul. ramoribus procumbenti-  
in edictoribus montibus S. Stephani Nicæensi.

9. ——— *purpureum*

G. fol. verticill. linearis-setaceis pedunc. capillar. fol. longior.  
abunde in agro Nicæensi.

10. ——— *lucidum*

G. fol. 6. rigidis obscure-vir. flor. e summo caul. prodeuntibus  
ubique in agro Nicæensi locis siccis.

11. ——— *cinerum*

G. caul. subligrolo fol. 6. long. elliptic. rigid. serrat. aculeat. var-  
chibus ovalis laevibus albescentibus  
in vineis Savorgii & Sorpittellii.

12. ——— *tenuifolium.*

G. fol. 6. subrigid. diffuse ramor. flor. alb. e summo caule prodeunt.  
in collium Nicæensium duneis.

13. ——— *murale*

G. fol. 6. linear. ram. simplicibus fructibus hispido-subscissilibus.  
therardia fol. florat. binis oppositis binis floribus.  
ad aridas rupes litorei Nicæensi.

14. ——— *maritimum*

G. fol. octon. lanceol. — linear. peduncul. 1. flor. caul. vespert. luto.  
In agro Nicæensi collectum est.

15. ——— *saccharatum*

G. sem. escaiperat. florul. lateral. mascul.  
Circa Nicæam provenit.  
Vera Varietas Valantia affavine L.

16. *Valantia muralis.* Coroll. 4. fida sem. tridentata.

fl. flor. mascul. trifidis hermaphroditi germine glabro  
insidentibus.  
in siccis & arenosis litorei Nicæensi.

Planta glabra caul. procumb. fol. verticill. 4. Corol. e viridi-  
pallens fruct. asperi.

17. *Rubia tinctorum*

ad muros & rupes ubique frequens in agro Nicæensi.







*N. C. L. ...*

*... in agro ...*

*... in agro ...*

*... in agro ...*

*... in agro ...*

*... in agro ...*

*... in agro ...*

*... in agro ...*

*... in agro ...*

*... in agro ...*

*... in agro ...*

*... in agro ...*

*... in agro ...*

*... in agro ...*

*... in agro ...*

*... in agro ...*



18. *Crucianella angustifolia*.  
Gen. Corol. 1. petal. intundibulifera. tub. filiformi limbo  
unguiculato 4. pido Calyc. 2. phyllis sem. 2. tenuia ob-  
longa flor. in spicis imbricatis folia verticillata.  
*C. erecta* fol. densis linearibus flor. spicatis.  
Secus arva Collium Nicaeensis.

19. ——— *monspeliaca*  
*C. procumbens* fol. acut. caul. 4. ovat. variet. linear. fl. spicat.  
Secus arva Coll. Nicaeni.

20. ——— *latifolia*  
*C. procumbens* fol. 4. lanceolat. fl. spicat.  
in Comitatu Nicaensi locis arenosis non infrequens.

21. *Salvia officinalis*  
in Coll. incultis agri Nicaensis

22. ——— *horminum*  
*S. fol. obtus. crenat. bract. sum. steril. majoribus coloratis.*  
Nicaentem agrum inhabitat secus arva nascitur.

23. ——— *verbenaca*  
*S. fol. serrat. sinuat. leeviuscul. Corol. Calyc. angustior.*  
in Comitatu Nicaensi non infrequens est.

24. ——— *aethiopis*  
*S. fol. oblong. eros. lanat. Verticil. lanat. bractes recurvatis*  
*sub spinosis.*  
Circa Sospello in Comitatu Nicaensi.

25. *Thymus vulgaris*  
in apricis salsis Collium Nicaenium sponte nascitur.

26. *Satureja juliana*  
*S. Verticill. fastigiatis fol. linear. lanceolat.*  
Circa Nicaeam in aridis

27. ——— *thymbra*  
*S. flor. Verticillat. hispida. fol. oblong. acut.*  
in salsis Collium Nicaenium

28. ——— *gracea*  
*S. pedunc. sub 3. flor. lateral. involucl. calyc. brevior.*  
in agro Nicaensi.

29. *Thymbra verticillata* Vide St. Monveg.  
*S. fl. verticillatis.*  
in agro Nicaensi.



30. *Lavandula stoechas*.  
L. fol. lanceol. lineav. integerr. spicis comosis  
in locis apvicis Maritimi agr. Nicæensi.

31. *Sideritis Romana* Vid. pl. figur.  
in avidis collibus Nicææ.

32. ——— *incana*  
*S. fruticosa tomentosa* fol. lanceolat. lineav. integerr. flor.  
bracteisq. dentat.  
inter *Laurene* & *Rocca Palliada*

33. ——— *hirsuta*  
*S. fol. lanceol. obtus. dent. pilos. bracteis dentat. spinos. caul.*  
*hirs. decumbentibus.*  
in Comitatu Nicæensi ad vappes & colles.

34. *Stachys brevifolia*  
*S. hirsuta* fol. obscure virentibus elliptico-lanceolatis, verticil.  
10. flor.  
in collibus Comitatu Nicæensi locis minime cultis.

35. ——— *hirta*  
*S. verticill. 6. flor. caul. prostratis corol. lab. superiori 2. fido*  
*divaricato reflexo.*  
Habitat in agro Nicæensi.

36. ——— *arvensis* V. p. figur.  
Nicææ secus arva.

37. *Phlomis lychnitis*  
Gen. Calyc. angulatus galea incumbens compressa hirta.  
Ph. fol. lanceol. tomentol. floral. ovat. involucri. setae. lanat.  
in Agro Nicæensi  
Caul. cubital. Corol. vix calyce major lutea.

38. *Molucella frutescens*  
Gen. Calyc. ampliat. campanulatus corol. latior spinosus.  
Ch. calyc. in profundis furcibus 5. fidi corol. calyc. longioribus.  
Fruticul. palmav. variet. foliis florigeris dichotomis foliis polluc.  
oval. breviter petiol. utrinque 2-3. profunde serrata pilis variis  
pubescentia nervosa venosa spin. areolaribus utrinque. Solit. ternat.  
ut in Berberide sed tant. ad florum verticillos. Cal. spinos. lo-  
bratus Corol. alb. galea lanata.  
Abunde inter Saorgio et Brelio et supra Ioppello & vespum  
filivis escit.

39. *Melissa cretica*  
Ch. racem. terminal. peduncul. solitar. breviss.  
Circum Lucerna.







*...fol. lanceolatis linearibus integris ...*  
*...in ...*

*...fol. lanceolatis linearibus integris ...*  
*...in ...*

*...fol. lanceolatis linearibus integris ...*  
*...in ...*

*...fol. lanceolatis linearibus integris ...*  
*...in ...*

*...fol. lanceolatis linearibus integris ...*  
*...in ...*

*...fol. lanceolatis linearibus integris ...*  
*...in ...*

*...fol. lanceolatis linearibus integris ...*  
*...in ...*

*...fol. lanceolatis linearibus integris ...*  
*...in ...*

*...fol. lanceolatis linearibus integris ...*  
*...in ...*

*...fol. lanceolatis linearibus integris ...*  
*...in ...*

*Circa lucerna.*



40. *Tenacium* *3ua*  
*T. fol. subtricuspidatis linear. flor. sessil. lateral. solitar.*  
*in Comitatu Nicæensi & Lincol. freq. ad Vias.*  
*Planta morchæm vedoleus Caut. lignos. humilis. hirs.*  
*subtus folia sessilia raro integerrima flor. magnus hirs.*  
*subtus purpureus Variat corolla lutea & alba.*
41. — *flavum.*  
*T. fol. cord. obtuse serrat. bract. integerr. concav. caud.*  
*pubicol. flor. tern. vacuol.*  
*in Agro Nicæensi ad rupes aridas sole percuttas.*  
*Nicæenses appellant Grasse Germanice.*  
*Caut. 2. ped. fol. petiol. bas. integerr. undulata dura utrinq.*  
*sub hirsuta supra atro virid. nitid. subt. canescent. floral.*  
*sessil. concav. Neutrilli. multi fl. utrinque 3. pedunculat.*  
*corol. ochroleuca vel alba raro purpur.*
42. — *Capitatum.*  
*T. capitul. pedunculatis fol. lanceol. crenat. tomentol.*  
*caule erecto.*  
*Minime varium in Comitatu Nicæensi & Liguria.*  
*Habitus majoranae sed tomentol. & angustifol. capitul. sub.*  
*volunda corol. alb. litur. Violæ.*
43. — *polium*  
*T. capitul. subrotund. fol. ovat. obtus. crenat. tomentol. sessil. caule*  
*prostrato.*  
*in sacris jugis Pyrenæis. Nicæam versus Cypri.*
44. *Lycopus* *variegata*  
*L. fol. repandis dentatis callos. caul. decumb. corol. cernuis.*  
*in agro Nicæensi.*
45. *Cinoglossum cheiripolium*  
*C. corol. calyc. duplo longior. fol. lanceol. incan. tomentol.*  
*Nicæa crescit Circa Lucerame Cauvini.*
46. *Lehim italicum*  
*L. caule erecto piloso spicis hirsutis coroll. subæqual. stamin.*  
*longiss.*  
*Locis siccis Nicæensibus Vulgare.*
47. — *lupitanicum*  
*L. caule simplicifol. caulic. lanceolat. serie. amplissimis*  
*flor. spicat. lateral. coroll. stamin. longior.*  
*in agro Nicæensi.*
48. — *plantagineum*  
*L. fol. radical. ovat. latit plantag. fol. simil. lineat petiol.*  
*utrinque ad Vias et loca sicca Nicæa.*



49. *Myosotis apula*  
in Comitatu Nicaeni.

50. *Orobancha purpurea*  
O. caule simplici bracteis tenuis calyce 5-fido.  
Locis apricis Satorii Comit. Nicaeni.

51. *Coris monspeliensis* V. pl. Monseg.  
in ha. bitat Colles sicco & rupes in agro Nicaeni.

52. *Euphrasia latifolia*  
E. fol. dentatis-palmatis flor. subcapitatis  
in agro Nicaeni.

53. ——— *viscosa*  
E. fol. linear calyce glutinoso hispido  
Orontites fol. viscidis varietes serratis.  
Locis aridis Collium Nicaenium.

54. *Bartia viscosa*  
B. fol. superioribus alternis serratis flor. distant. lateralibus.  
in agro Nicaeni.

55. *Bellardia triscago*.  
Gen. Calyce bilabiato labio superiore semibifido inferiore  
integro Galea vestita barba trifida segmento medio ampliore  
Fructus est Capsula ovata biloculari bivalvi polysperma.  
Bartia fol. oppositis linear. lanceolatis inciso-serratis spicis  
foliosis compactis  
Achiranthus calycibus hirsuto-tomentosis fol. oppositis obtuse  
serratis caule simplicissimo.  
Nascitur in litore Nicaeni prope Lanterna.  
Caul. palmat. fol. 3-uncial. angusta. Spica longa quadrata densa e  
cristatis foliis robustioribus Euphrasiae vulg. simil. compacta.  
Fl. magni purpureascentes barbati albi pallentibus Galea vestita  
Capsula nigra.

56. *Antirrhinum roseum*.  
A. fol. linear. conturb. inferne 4-nis calyce caput-angustibus  
in agro Nicaeni.  
Hoc differt ab A. monspeliense.

57. ——— *bipunctatum*.  
A. fol. linear. glabr. inferius 4-nis caule erecto paniculato  
fl. spicato capitatibus.  
in arvis Collium Nicaenium.



59. *Antirrhinum petraeanum*  
Hb. 2 caul. lineari-bus ad radic. oval. sessile breviss.

1/2 L. 2 Caul. lineariibus ad radic. oval. sessilibus;

601.

*L. latro*

in female  
d. 2 amplification  
by sperm.  
obst. sp. 1/2  
obst. oblique

devia e  
m h. u. c. n.  
a vostra

Wagner Library

Maniculah



49. Myosotis apula  
in Comitatu Nicæensi.

50

51

52

53

54

55

56

57

58. An

59. Juv

60

61

62

63

64. Planta

65

66

67

68

69



58. *Antirrhinum peltavianum*  
A. fol. ad caul. lineariibus ad radic. oval. saepe ternis  
A. fol. caul. n. altern. radical. lanceol. flor. corymbos.  
in agro Nicaensi

59. *Scrophularia auriculata*  
S. fol. cordato oblongis basi appendiculatis subus tomentosis  
racem. terminalibus.  
Ad fontem in agro Nicaensi.

60. ——— *Scrodonia*  
S. foliis cordatis duplicatis serratis racemo composito foliis intermixto  
Agri Nicaensis indigena planta est.

61. ——— *betonicefolia*  
S. fol. cordat. oblong. dentatis dentibus integerrimis  
in agro Nicaensi

62. ——— *peregrina*  
S. fol. cordat. lineatis lucid. peduncul. axill. 2. flor. caule 6. angul.  
in umbrasis Agri Nicaensis.

63. ——— *lucida*  
S. fol. interior. bipinnatis subcarnos. glaberr. racem. 2. parvis.  
in agro Nicaensi locis saluberr.

64. *Plantago altissima*  
p. fol. lanceol. 5. nerv. dentat. glabr. scap. subangulat. spica  
oblonga cylindrica  
Jecur le Pailhon prope Nicaeam.

65. ——— *lagopus*  
p. foliis lanceolatis subdentulatis spica ovata hirsuta scap. teret.  
in Comit. Nic. copiose

66. ——— *albicans*  
p. fol. lanceol. obliq. villis. spic. cylindr. erect. scap. teret.  
in agro Nicaensi

67. ——— *Bellardi*  
p. pubescens fol. sublineariibus plan. petiol. scap. teret. fol.  
longitud. spicis cylindricis  
invenit. Cl. Bellardi locis sterilibus Villafranca Nica-  
ensis prope lanternam

68. ——— *maritima*  
p. fol. semicylindricis integerr. basi lanat. scap. teret.  
in agro Nicaeae ad rupes.

69. ——— *Coronopus*  
frequens in agro Nicaensi.



70. *Plumbago Europaea.*

Calyx 5. dentatus persistens Coroll. infundibuliformis limbo 5. fido Stamina  
squamis basium corollae claudentibus inserta. Stigma 5. fidum Capsula 1. lo-  
culari 5. valv. 1. sperma.

Pl. fol. ovato-lanceol. amplexic. ciliat. scabr. fl. congest. Stam. erectis.  
in Agro Nicaensi.

Caul. 2. cubit purpurascens fol. quandoque obverse ovata atrovirentia florum  
congesti Calycis erecti tuberculis basi 5. praeter glandulas petiolatos Coroll.  
purpurascens.

71. *Anagallis Verticillata.*

A. erecta fol. caul. Verticill. axill. folior. in caule & ramis florigeris  
flora corulea.  
in Com. Nicaensi. Pl. Bellardi.

72. *Cressa cretica. pentandria Digynia.*

Calyx 5. phyll. persistens. Coroll. tubulos. infern. ventricos. limbo 5. fido  
filamenta tubo insidentia Capsula 1. locut. 2. valv. 1. sperma.

C. Coroll. imberbibus capsula 1. sperma.

Nicaensi agri indigena est prope hospitium di. Villafranca.

Planta parva humifusa incana. caulicul. dodrant. ramori subrubent.  
fol. minim. altern. sessil. ovat. vel lanceolata florum inter folia pusilli  
sic albo purpurasc.

73. *Junculus Valerandi* Vis. Pl. Valder.

ad fontium scaturigines in agro Nicaensi.

74. *Lysimachia linum Stellatum*

L. calyc. coroll. superantibus caule erecto ramorissimo fol. subulatis  
oppositis peduncul. folio brevioribus.  
in agro Nicaensi frequent.

75. ——— *tenella.* V. Pl. Liguor.

locis aquosis secus flumen le Var & ad oppidum Lucerame.

76. *Gentiana spicata*

G. coroll. 5. fidi infundibuliform. fol. altern. sessil.

in pratis & herbiculis locis Nicaensibus

Caulis plerumque unicus subdodrant fol. opposit. hypervic. mac. fl. termin.  
albi vel rubelli.

77. *Nerium Oleander*

in montibus Nicaea & collibus Sympitellensibus

78. *Pervinca major*

frequent in Comit. Nicaensi.

79. *Aclepias nigra*

A. fol. basi barbati caule superne subvolubili fol. cordati undi  
flor. confert. patentissimi petalis nigri.

in collibus Nicaensibus & unctis locis sacris & editionibus nascitur.



lila / ramina  
capitata lila  
st. / ram. erecta  
ramina florum  
violaceo caeruleo  
diffusis  
limb. 5-fido  
eterna  
la / pancia  
subtubent  
filia parilli  
fol. subulati  
m. lucerna  
st. / ter min.  
cordat. nudis  
d. / kon / turkasi / tur

al. fib. ment. flor. fol. cum flor. erump. albis. plant. paull. ultra 2. 1/2



*[Faint, mostly illegible handwritten text in a cursive script, likely Latin or Greek, covering the majority of the page.]*

flor. confert. parenthemi, pterali, nigri.  
 in collibus et caenibus & unctis locis, siccis, & editionibus nascitur

80. Cyn...  
 81. Hyocy...  
 82. —  
 83. Ver...  
 84. —  
 85. Conu...  
 86. —  
 87. C. fol...  
 2. flori...  
 inap...  
 Tota flo...  
 obtusior...  
 88. C. compl...  
 folio...  
 in syl...  
 89. —  
 C. cau...  
 agra...  
 90. Colchic...  
 C. flore...  
 in m...  
 afflu...



80. *Cynanchum monipeliacum*.

Gen. Nectario cylindraceo ovifloro 5-dentato ab *Asclepiade* differt.  
C. caul. volubili herbaceo fol. reniformi - cordati acuti  
Rarum in agro Nicaeensi nascitur.

Radice & caul. lactiferi fol. opposita longe petiolata integerr. crassa  
subalbida fl. axill. fascicul. parvi alb.

81. *Hypocyanus albus* V. pl. lig.  
in comit. Nicaeensi.

82. ——— aureus V. pl. lig.  
ad muros Nicaeae.

83. *Verbascum phlomoides* V. pl. lig.  
in aridiss. & siccis comit. Nicaeensi. abunde circa Villafrauca

84. ——— sinuatum V. pl. ligur.  
saepe occurrit in comit. Nicaeensi.

85. *Convolvulus althaeoides* V. pl. ligur.  
in comit. Nicaeensi ad super. mar. proximam, & ad muros in vineis.

86. ——— lineatus  
C. fol. lanceol. sericei lineat. petiolat. pedunc. biflor. calyc. sericei  
subfoliacei.  
Nicaea nascitur.

Caulis prostratus flexuosus semiped. inanis fol. remota basi angustiora acuta  
ad apiculum acutum lineata Hyact. 2. Calyce longiore Corol. dilute  
purpurea. exst. hirta.

87. ——— *Cantabrica*  
C. fol. lineari-lanceolat. acut. caule erectiusc. ramoso: pedunc. sub.  
2-flori calycibus pilosis.  
in agro Nicaeensi abunde

Tota planta villosa caul. sesquiped. fol. altern. integerr. radical. petiolat.  
obtusiora; caulic. sessil. pedunc. axill. cor. carnea c. lineis rubris.

88. *Campylosiphium medium* C.  
Capsul. 5-locul. obtechi; caule indiviso erecto  
foliis floribus erectis.  
in sylvis apricis & locis incultis agri Nicaeensi.

89. ——— *erinus*  
C. caul. dichotomo fol. sessil. superior. oppos. induratis  
agri Nicaeensi indigena.

90. *Colchicum montanum*.  
C. flore cum foliis conjuncto petal. linear.  
in praet. macilentis montanis non tamen editioribus circa Ispello.  
altiss. mont. flor. fol. cum flor. erump. altiss. plant. paull. ultra 2. digit.



91. *Agave americana*. Mercur. monogyn.  
 Gen. coroll. erecta superba & spica effusiva. Stamina corollae lon-  
 giora capsula 3. locul. 3. valv. polypetala. fol. Carnos. succulent.  
 A. fol. dentato spinos. scap. amolo.  
 Ad vases in agro Nicaeensi inter Villafranca & Monaco.  
 Fol. applanatissima succulenta hinc convexa inde plana mucrone duriusculo  
 firmi peroram spinis obita. Scap. erect. etiam 30. pedat. simplici  
 spica ramosa fl. innumeri petala sublatea.

92. *Erica scoparia* V. sp. Lig.  
 C. Bellardi in montibus di. Cathon in Comitatu Nicaeensi.

93. *Cotyledon umbilicus* Vener. V. pl. Ligur.  
 in Comit. Nicaeensi. ad muros & vases

94. *Zarminum fruticosum*  
 ad sepes in agro Nicaeensi

95. ——— officinale  
 frequens in comitatu Nicaeensi

96. *Olea europaea*  
 in Comitatu Nicaeensi

97. *Phillyrea media* Diandria monogyn. L.  
 Calyx 4. dent. corolla 4. fida rotata Baccæ monosperma.  
 sp. fol. ovato-lanceolatis subintegerrimis  
 locis siccis & vasis in agro Nicaeensi

98. ——— angustifolia  
 sp. fol. lineari lanceol. integerr.  
 iidem crescit locis

99. ——— latifolia  
 sp. fol. ovato cordatis serratis.  
 Orietur ad vases Montalbani in agro Nicaeensi.

100. *Momordica elaterium*.  
 in agro Nicaeensi provenit.

101. *Nitescens agnus catharticus* V. sp. Ligur.  
 inter Mentan. & Monaco.

102. *Lycium europaeum* Pentandria monogyn.  
 Cal. 5. dentatus coroll. in fundiculis. Stamina non defluunt. Bacc.  
 villosa. baccæ 2. locul. polisperma.  
 fol. obliq. ramul. flexuos. teret.  
 Fructus ramor. fol. 5. 5. etc. ead. gemma prodeunt fl. singulis gemmis  
 25. cal. 5. fida coroll. fauce pallid. nigro. int. int. purpurea.



liggenwij  
v.a/c.

44.

U.

my.



91. *Agave americana*. *Hermodactylus*

92

93

94

95

96

97

98

99

100

101

102

Fruktus v. m. fol. 5. 6. etc. ead. gemma prodeunt fl. e. lingulis gemmis  
2. 5. cal. 5. fidus corol. fauce pallid. nigro. int. int. purpureo.



103. *Viburnum hirsutum*  
 4. fol. ~~integerrimis~~ ovatis ramificationibus Ucnarum subny  
 Villoso glandulosis.  
 in Sylvis opacis Nicaeensibus.

104. *Thymatea lauroleuca*  
*Daphne* racemis axillaribus 5. flor. fol. lanceol. glabris  
 Tota planta flexilis supra medium ramosa ramis apice foliis pubescent. 2. pedat.  
 erectis cinereo-fuscis. Fol. digit. longa laevia & similitudine alterna breviter.  
 petiol. perennant. crass. integerr. supra laevia. virid. subt. dilutiora.  
 Racemi nutantes bracteis concavis alternis Fl. frutres.  
 in Sylvis umbris non humidis Mont. Comit. Nicaeni. sparsim.

105. *L. lanamunda.*  
*D. thymatea* flor. sessil. axillar. fol. lanceol. caul. simpliciss.  
 Caul. plant. palm. fl. sparii non racemosi corol. ex albo hutea.  
 Circa Nicaeam.

106. *—— tarton-raila.*  
*D. fl. sessil. aggregatis axillar. fol. ovatis utrinque pubesc. nervos.*  
 in agro Nicaeni.  
 Fruticulus fol. mollia candicans. & velut. Sericea fl. albo pall.

107. *—— quidium*  
*D. panicula terminali fol. linear-lanceol. -acuminat.*  
 Frequens in Comitatu Nicaeni.  
 Caulis humilis fruticulus. Cubit. & ultra ramos. fol. Viridis. annua glabra.  
 fl. congest. alb. supra rubri cinct. fol. suaveolent. fruct. matur. rubri  
 succosi Bis quotannis florent.

108. *Arbutus Unedo* Decand. Monogyn. 4.  
 Cal. brev. 5. fid. petiost. Coroll. campaniform. globos. 5. fid. Ovar.  
 intra florem. fructus 5. locust. polysperm.  
 A. caule arboreo fol. glabr. serratis baccis polyspermis.  
 in agro Nicaeni abunde. Arbor.  
 Frutere magnis. fol. laevib. altern. petiol. oblong. duris late  
 viridia nitida corol. herbido rubra Balca magna sphaerica. rubi-  
 cunda hirsuta. sem. exasperata.

109. *Styracis officinale.* Decand. monogyn. Cal. 5. dent. hypocrater.  
 Corol. infundibuliform. in 5. vel 6. profund. lacin. divisa fructus est  
 drupa 1. locust. 2. sperma.  
 In agro Nicaeni. abunde in rup. marit. prope Joet.  
 Arbore. fol. altern. petiol. ovatis supra virid. nitent. subt. tomentol. mal.  
 Cotton. simil. sed minor. pedunc. saepe 2. flor. Cor. alba.



110. *Scabiosa leucantha*  
f. corollul. 4. fid. subgymn. squam. calyc. ovat. obtus. fol. pinnatifid.  
in collibus aridis Nicaeni. abunde.

111. — *integrifolia*  
f. corollul. 4. fid. fol. indivis. radic. ovat. serr. ramis lanceol. caul. herbac.  
abunde secus arva in agro Nicaeni.

112. — *grammifolia*  
f. corollul. 5. fid. calyc. breviss. fol. caul. bipinnat. filiformibus.  
Nicaeni in agro indigena est. *Scabiosa stellata*. addenda.

113. *Globularia alyssum*. V. pl. figur.  
in sylvis et incultis agro Nicaeni.

114. *Ambrosia maritima*. Monoclea pentandria.  
chalc. cal. comm. 1. phyll. 2-8 fid. corol. inmundibul. 3. fid.  
recept. nud.

Form. cal. 2. phyll. integ. ventr. 5. dentat. 1. fl. corol. o. nuda  
e calyce indurata monosperma.

A. foliis multifidis spicis solitariis pilosis sessilibus.  
in arenosis maritimis Nicaenibus.

Caul. subped. ramosis puber. sicut fol. & racemi folia saepe  
opposita petiolata mollissima albescens Racemi densi fl. subressi.  
conferti corol. vivid. lutea.

115. *Xanthium spinosum* V. pl. figur.  
ad maris latus occurrit.

116. *Echinops ritro*.

E. foliis supra glaberrimis subtus tomentosis pinnatifid. capitul.  
globos. caul. multiflor. corymbos. flor. caeruleis.  
abunde in montibus Nicaenibus.

117. *Carduus leucographus*

C. fol. decurrentibus dentatis spinosis peduncul. nud.  
longissimis uniflor. calycibus spinosis inclinator.  
Cirr. maculis argenteis notat.  
in olivis & rudibus circa Nicaeam.

118. *Cirsium oleraceum*

Cirsium fol. pinnatif. farinatis nud. amplexic. inerm. bracteis  
subcoloratis integris concavis flore flavicante  
in humoribus Nicaeni agro.



pl. spiralis

rostr. caud. hinc

l. p. m. b. u. s.

addenda

ia.

3. p. d.

o. n. u. r.

l. u. r.

l. a. l. o. p. e.

l. l. r. u. b. r. i. f. i.

cap. h. u. l.

al. n. u. d.

al. h.

Wach. i.

C. l. u. c. e. r. a. m. e.

o. l. i. n. e. l. l. u. m.

o. f.

o. f.

pro. 100.0 atant.  
in montan. circa Lucetame.



110.

111.

112.

113

114

11

11

1

119. (a)

120. -

at

121. C

G

n

P

C

lycl

6

122. C

123. C

Gen

em

col

in

124. C

125. C

126. C

127. C

128. C

129. C

130. C

131. C

132. C

133. C

134. C

135. C

136. C

137. C

138. C

139. C

140. C

141. C

142. C

143. C

144. C

145. C

146. C

147. C

148. C

149. C

150. C

151. C

152. C

153. C

154. C

155. C

156. C

157. C

158. C

159. C

160. C

161. C

162. C

163. C

164. C

165. C

166. C



119. *Cirsium tuberosum*.  
C. radicibus tuberosis fol. semipinn. pinn. angulos. ciliat. fl. nud.  
Ubique in agro Nicaensi.

120. — *stellatum*.  
Carduus fol. integr. sessil. inerm. subt. tomentos. pinn. ramos.  
aristat. flor. sessil. lateral.  
in agro Nicaensi ad vias secus arva.

121. *Acaena cancellata*. *Attractyl.* Linn.  
Gen. Calyce circumdat. bract. foliac. corollat. hermaphrod. b. fid.  
receptac. paleac. paleis in 2. aut 3. aristas abeuntibus in nostra  
planta sem. Vittor. papp. plumos. coronat.  
Attract. involucri cancellati. Ventricosi linearibus dentatis ca-  
lycibus ovatis floribus flocculosis  
ad Rupes Montalbani in agro Nicaensi et ejusdem Comit.  
loc. Therit. praecalis.

122. *Carthamus tinctorius*.  
Ubique in ar. di. collibus Nicaensibus.

123. *Carduncellus mompelienis*. *Carthamus* L.  
Gen. Calyce imbricatus squamis appendice foliacea auctis floccul.  
omn. hermaphrod. limbo b. fido receptac. paleac. sem. papp. simpl.  
coron. Carth. fol. caul. linear. pinnat. longitud. plantae.  
in montibus interpositis oppido Lucevame & Lantolen.

124. *Cnicus ferox*. V. M. Tard.  
Cl. Dana in collibus d. Braun.

125. — *acarna*  
C. fol. decurr. lanceol. indiv. calyc. pinnat. spinos.  
in Comit. Nicaens. freq. loc. Therit. Sarcot. & ad vias.

126. *Carlina lanata*  
C. caul. b. fido calyc. sanguin. terminal. primo aristati sessil.  
ad marit. litus locis siccis.

127. — *achanthifolia*  
C. acant. fol. tomentos. semipinn. plicatis pinn. semipilob. pinn.  
gentibus.  
In montibus Braun alique Sorpella praecaris frequent.

— Cent. panicul. Var. flor. albo ad rupes in agro Nicaensi.

128. *Centauria cineraria*  
C. calyc. ciliat. terminali sessil. lobis fol. tomentos. bipinn.  
lobis lobis acutis.  
in montibus circa Lucevame.



129. *Centaurea sonchifolia*.

C. calyc. palmatis spinosis fol. decurrentibus spinulosis  
veppando-dentatis  
loc. 7 aridis collium Nicæensium.

130.

seridis

C. calyc. palmatis spinosis fol. decur. toment. orbiculis infim.  
sinuato-dentatis.  
in agro Nicæensi

131.

ilnardi

C. calyc. palmatis spinosis fol. lyvato-dentatis hispidi;  
subamplexicaulis flor. sessil. terminat.  
in vineis Nicæensibus & arenosis maritimis.

132.

aspera

C. calyc. palmatis spinosis fol. lanceolatis dentatis.  
locis sterilib. Comit. Nicæens.

133.

Nicæensis

C. calycibus ciliatis spinosis fol. virid. asper. ovatis. rad.  
cal. petiol. dentatis. caul. n. amplexicaul. decurrent.  
Cl. Bellardi circa Nicæam legit.

134.

galactites V. pl. ligur.

ubique abunde in Comit. Nicæens.

135.

*Tanacetum annuum*.

T. fol. bipinnatifid. linear. acut. corymbis tomentosis.  
Nicææ, et in arvis Nicæens. secus flumen le Varrou. infrequens.

136.

*Anacyclus Valentini*

Gen. Calyx imbricat. hæmiphor. placent. paleac. flor. disc. 5-fid.  
hermaphrod. sem. compress. nuda in amb. h. form. paucae corol.  
lingul. brev. integerr. sem. alata apic. emarginata.  
A. fol. decompositis linearibus lacin. div. ferebianscul. acut.  
flor. flosculosis.  
in Agro Nicæensi

137.

*Artemisia Gerulescens*

A. fol. caul. lanceol. integr. tomentosis. radical mul-  
tiplo. flor. long. spicat. tenuis.  
Nicæense litus inhabit.







137. *Conium maculatum*  
Herb. perenne, foliis bipinnatis, semine  
negundo, semine  
Herb. perenne, foliis bipinnatis, semine

138. *Conium maculatum*  
Herb. perenne, foliis bipinnatis, semine  
negundo, semine  
Herb. perenne, foliis bipinnatis, semine

139. *Conium maculatum*  
Herb. perenne, foliis bipinnatis, semine  
negundo, semine  
Herb. perenne, foliis bipinnatis, semine

140. *Conium maculatum*  
Herb. perenne, foliis bipinnatis, semine  
negundo, semine  
Herb. perenne, foliis bipinnatis, semine

141. *Conium maculatum*  
Herb. perenne, foliis bipinnatis, semine  
negundo, semine  
Herb. perenne, foliis bipinnatis, semine

142. *Conium maculatum*  
Herb. perenne, foliis bipinnatis, semine  
negundo, semine  
Herb. perenne, foliis bipinnatis, semine

143. *Conium maculatum*  
Herb. perenne, foliis bipinnatis, semine  
negundo, semine  
Herb. perenne, foliis bipinnatis, semine

144. *Conium maculatum*  
Herb. perenne, foliis bipinnatis, semine  
negundo, semine  
Herb. perenne, foliis bipinnatis, semine

145. *Conium maculatum*  
Herb. perenne, foliis bipinnatis, semine  
negundo, semine  
Herb. perenne, foliis bipinnatis, semine

146. *Conium maculatum*  
Herb. perenne, foliis bipinnatis, semine  
negundo, semine  
Herb. perenne, foliis bipinnatis, semine

138. *Microglossa*  
Calyx  
5. den  
penn  
chr. c  
M

139. *Filago*  
F. flor  
in  
Aure can  
Mori fane  
Lana. Fl.

140. *Guaiacum*

141. *Conium maculatum*

142. *Stachys*  
Calyx  
phrod  
J.  
Naf

143. *Conium maculatum*  
Caul. pe  
iter. fan  
Calyx. f

144. *Conium maculatum*  
C. fo  
acu  
re

145. *Conium maculatum*  
ad

146. *Conium maculatum*  
n. 6.



138. *Micropus supinus*.

Calyce. 5-phill. calyculatus placenta nuda flor. discipnascendi corol.  
5. dentata styli rudim. habentes & abostientes in ambitu flores  
feminei nudi sem. nuda squam. Calyc. inclusa.  
Chr. caul. procumb. fol. gemin. Rachis muriculi  
Nicaea ad mari latus.

139. *Filago acaulis*.

foliis  
F. flor. acaulibus sessil. floral. majoribus  
in stagnis exsiccatis Nicaeensis

Raro caulescit id accidit locis fertilibus & tunc. caul. superius va-  
mosi ram. parul. in rostrum - fol. imbricata linearia tomentosa - in-  
cana. Fl. alii majores alii minores Viridescens.

140. *Guaphalium Stachas*.

G. pubesc. fol. linear. var. virgat. Cymis. composito.  
in aridis collibus Nicaeensis abunde.

141.

luteo album

G. herbaceum fol. semiamplexice. ensiform. repand. obtus.  
utrinque pubescent. flor. conglomerat.

Abunde secus rivulos & fluentes aquas in agro Nicaensi.

142. *Stachelina Dubia* L.

Calyce. oblongus imbricatus ex squamis coloratis flor. herma-  
phrod. 5-phill. placenta paleacea semina. papp. plumos.

J. fol. linear. denticulatis squamis. calycinis lanceolatis

pappo calyc. duplo longiore.

in Agro Nicaensi.

Caul. ped- cubit incano-tomentosi. Rami pl. erecti circ. med. 1-2 pe-  
dit. ramosi. fl. 2-3. conglomeratis fol. plur. inordinat. sessil. fl. oblongi.  
Calyce. squamulis purpurascens

143. *Corypha Sarcobatis*

C. fol. linear. subdent. peduncul. longit. uniflor. capitulo  
acuto & angust.  
frequens ad muros in Comit. Nicaensi.

144.

Sordida. V. pl. Ligur.

ad Rupes & Vetus muros in Com. Nicaensi.

145.

Buphtalum. spinos. V. pl. Ligur.

ubique ad Rupes & vias in agro Nicaensi.



146. *Buphtalmum agrarium*  
*B.* calyc. obtuse foliis. sessil. arcu. fol. altern. oblong. obtus. caul. herbac.  
 in agro Nicaeensi ad font. scaturigines.
147. ——— *maritimum*  
*B.* calyc. obtuse foliis. peduncul. fol. altern. spatul. caul. herbac.  
 in agro Nicaeensi prope marit. litus.
148. *Achillea ageratum*  
*A.* fol. lanceol. obtus. acut. serrat. flor. umbell. latis.  
 in Comit. Nicaeensi. Frequens circa la Turbia et secus flum.  
 spailion in arenosis.
149. *Chamaemelum maritimum*  
*Anthem.* fol. pinnat. dentat. Carnos. nudis pinn.  
 stalis caul. prostrato calyc. subtomentos.  
 in agro Nicaeensi.
150. ——— *tomentosum*  
*A.* fol. pinnatif. obtus. plan. pedunc. hirsut. foliis. calyc.  
 tomentos.  
 in arenosis maritimi agri Nicaeensi.
151. ——— *mixtum*  
*A.* fol. simplic. dentat. laciniat. coronop. simil.  
 in Coll. Nicaeensi. mar. prociis.
152. ——— *nobile*  
*A.* fol. lanuginos. pinnat-composit. linear. acut.  
 in Comitatu Nicaeensi.
153. ——— *valentinum*  
*A.* caul. ramor. fol. pubesc. 3. pinnat. setac. calyc. villos.  
 peduncul.  
 in tabulor. flum. spailion in agro Nicaeensi.
154. *Calendula arvensis*  
 Jecur arva in agro Nicaeensi.
155. *Cysanthemum discordium*  
*Cohula* fol. altern. lanceol. dentat. caul. erect. fl. solit.  
 ubique locis aridis in Comit. Nicaeensi.
156. ——— *inodorum*  
*C.* fol. pinnat. multifid. caule ramoso diffuso.  
 inter segetes in agro Nicaeensi.
157. ——— *montanum*  
*C.* fol. imis spatulat. lanceol. serrat. sum. linear.  
 in monte Syran aridis & siccior locis nascitur.  
 Vix a *C. leuchanthemo* differt.



47.

9.

1/111.  
1/.

ymb.

in montibus circa Zuerame li. Richardi prope larnou etc.



146.

147.

148

149

150

151

152

153

154

155

156

Vix a C. leucanthemo differt.



lanche

647.

sup.

plan.

pl.

orgmb.

in montibus circa Lucerne Cl. Bellard proprietarius C.C.



146

147

148

149

150

151

1

152

153

154

1

Vix a C. Leuchanthemo differt.

158 Chrysanth

159

160

Caul. 2. h.  
vel integr.  
futei pale

161 Belli  
A.  
in

162 Aster

C.  
J.

163

164

165. Evig

166

167 Senecio

168 Ciner  
Abunde a  
supra 1/2

169. Lappan  
Lappana Ca  
in me



158. *Chrysanthemum Segetum*  
C. fol. amplexic. superm. lacin. inferm. dentato-serratis.  
in salubriori fl. & Var. et ad marit. l. hui.
159. ——— *myconis*  
C. fol. lingulat. obtus. serrat. Calyc. squam. aequalibus.  
in agro Nicaeensi.
160. ——— *coronarium*  
C. fol. pinnat. incis. echinorum latioribus  
Cum superiori respicit. est a D. Grudice.  
Caul. 2. l. pedat. Ramor. sulcat. folios. fol. sessil. pinnis vel semipinn.  
vel integr. pinnat. Extrem. major. 2. fid. moll. virent. flor. ampl.  
lutei pallescentes.
161. *Bellis annua*  
fl. minima praeterea caule folioso.  
in gl. hui. quamvis Nicaeensibus sed rara.
162. *Aster Hippolium*  
C. fol. lanceol. integerr. carnos. glabr. ram. inaequat. flor. corymb.  
ad mar. Nicaeensi. l. hui. loc. paludos.  
Caul. cubit. Ramor. striat. Ruffere. pedunc. nud. l. flor. Calyc.  
squam. obtus. coroll. rad. coroll. discus luteus.
163. ——— *acris* V. pl. liguv.  
in Comit. Nicaensi.
164. ——— *bifrons* V. pl. liguv.  
in Com. Nicaensi. non in freq.
165. *Erigeron Viteorum*  
C. peduncul. l. flor. lateral. fol. lanceol. denticul. Gali refle-  
xos. Calyc. squam. coroll. radiat.  
in Comit. Nicaensi.
166. ——— *graveolens* V. pl. liguv.  
loc. & aridis agris Nicaeensibus.
167. *Senecio Sylvaticus* V. pl. liguv.  
abunde in C. Nicaensi.
168. *Cineraria maritima* V. pl. liguv.  
Abunde ad marit. Nicaeensi. l. hui. nec non ad v. sup. Roque Tagliade  
supra l. l. carene.
169. *Lappana fatida* Calyc. ex squam. pat. l. l. unius ordinis. Sem.  
nudum.  
Lappana caule nudo indiviso fol. semipinnat. pinn. retraversi. dentat.  
in montibus circa Lucerame et Bellavide prope Carthou Nic.



170. *Lappula stellata*

l. calyc. fructus undiq. patentibus rad. subul. fol. caul. lanceol.  
indivisi.  
Nicæa observata est.

171. *Catananche cerulea* V. M. figur.

abund. in sterilibus Comit. Nicæens.

172. — *lutea*

C. squam. calycin. inferior. lanceol.  
prope Breglio.

173. *Scolymus maculatus*.

Calyc. imbric. sem. coronat. avert. 2. aut 3. facile decidui.  
J. floribus solitariis  
Anetia & Nicæa ad maris litus.

174. — *hispanicus*.

J. floribus congestis.  
in agro Nicæensi.

175. *Wewha barbata*.

Cal. ad basin calycul. ordine squam. linear. sem. tam disci quam  
peripherie avert. 2. instruct.  
Crepit involucri. Calyc. longior squam. setac. spurs.  
in agro Nicæensi ad limites agrorum.

176. *Leontodon bulbosum*

l. fol. oblong. ovat. subdent. glabr. calyc. lœv. scapp. superh.  
hirsut.  
Nicæensi & agri indigena planta est.

177. *Picris tuberosa* V. M. figur.

in pratis Nicæensibus frequens est.

178. *Hieracium Capillaceum*

H. caul. tabund. uniflor. fol. nervos. gravim. integerr.  
hectum circa Sorpello.

179. *Crepis buvisifolia*

C. fol. pinnatifidis crenatis scapo nudo paniculato.  
Agri Nicæensi indigena planta est.











180. *Crepis albidus*  
*C. fol. runcinato-dent. incan. squam. calyc. simpl. membranac.*  
*fl. albid.*  
 locis Juncorum montanis redhibetur apertis in Com. Nicæensi.
181. — *vulva*  
*C. fol. amplexic. lyrat-runcinat. f. h. fl. suaverubenti.*  
 in Com. nicæensi. secus arva inter Sospello & Leucarene
182. — *pusilla* v. *fol. ligur.*  
 in vineis & oliveis Nicææ.
183. — *nemantensis*  
*Hieracium sanctum Jacq. nud. multiflor. fol. lyrat-obtus. dentat.*  
 in sterilibus circa Quavasa in Com. Nicæensi.
184. — *pyrenæica*  
*H. caul. ramos. folios. fol. semiamplexicaul. oblong. repand. hisp.*  
 ad limites agrorum in Comit. Nicæensi.
185. — *echinodes*  
*Picris perianthiis exterior. 5. phyll. inter. aristato majoribus*  
*frequent in agro Nicæensi.*
186. *Juncus maritimus*  
*J. fol. omn. integr. denticul. spic. ram. 1. flor. semiflorul.*  
*5. dentat.*  
*J. peduncul. nud. fol. lanceol. amplexic. indiv. vel 2. argut.*  
*dentat.*  
 in humidis & ad fluent. ag. in agro Nicæensi. nascitur.
187. — *picroides*  
*Scorzonera fol. superior. amplexicaul. integer. infer. runcin. pe-*  
*dunc. squamos.*  
 in Com. Nicæensi.
188. *Lactuca saligna*  
*L. fol. im. semipinn. super. amplexic. hamat. nerv. spinos.*  
 in Com. Nicæ. secus arva copiose evasit.
189. *Prenanthes viminea* v. *St. Jend.*  
 in Com. Nic. prope le Scarene Sospello.
190. — *variorum*  
*Lactuca variorum. pene ligurica fol. radical. pinnatif. caul. lanceol.*  
 in salubris ad maris Nicæensi. libet.



191. *Alhagadiolus hedyotidis*. Hyoseris vel Lappiana L.  
Gen. demum a radic. Sapium nuda Gen. Valia pappo aut ignam.  
Coronata.  
Hyoser. fruct. subglobosis glabris caule ramoso  
in vineis et arvis incultis agri Nicæens. nascitur et in arenis.  
Jecur fl. pallidum.
192. ——— creticum  
H. fruct. ovat. scab. caule ramoso.  
Jecur arva in agro Nicæensi.
193. ——— scaber  
H. scapp. simplicissimo l. floro fol. scab. sem. pappos.  
in olivet. Villa franca montis alban: ad mania urb. Nicæ.
194. ——— *Racintia*  
Lappiana calyc. fruct. torulosis depress. obtus. Jecur.  
loca inculta mari Nicæensi: proxima inhabit.
195. ——— radiatum  
H. scapp. uniflor. nud. fol. glabr. runcinat. angulis dentatis.  
in collibus maritimis Nicæensibus lectus est.
196. ——— *kavarencoides*  
H. fol. sinuato dentatis sem. central. papp. plumos. pilis folior.  
juventibus radice fibrosa  
in rupibus circa Lucervam legit D. Cavvini.
197. *Scorzonera hispanica*  
J. caule ramoso fol. amplexic. integr. serrulatis.  
in pratis comit. Nicæens.
198. ——— laciniata N. Pl. Vall. Macra.  
in com. Nicæens. freq. est.
199. *Tragopogon Dalechampii*  
J. calyc. l. phyllis. corol. brevior. inerm. fol. runcinat.  
in vineis & arv. Nicæens.
200. ——— *picroides* V. Pl. Lignv.  
in olivetis Nicæensibus.



ficulat.

ab iguam

in culat.

ipm.  
ulst-licia

erit.  
t. hat.

Denk

gip/ist.

ulahi.

incinat.



191

19

193

194

195

196

197

198

199

200

201. G

202. T

203. H

204. F

om  
J.

205. A

C

206. —

207. Cn

g  
3

F  
16

208. Tr

T

209. Bu

Ca  
br  
ia



201. *Geropogon glabrum*.  
Cal. 1. phyll. profund. 8. fid. Sem. d. rei pappi plumos. Rad. ar. denticulat.  
coronata.  
G. fol. glabris gramineis flore dilute incarnato.  
in agro Nicaeni.

202. *hirsutum*.  
G. fol. pilosis caule hirs. calyc. corol. brevior. peduncul.  
superne incrassat. fl. luteo.  
in herbidiis collibus Nicaenibus.

203. *Hypochaeris radicata*  
H. fol. vuncinatis obtusis scabris caule ramoso undato laevi  
pedunc. squamol.  
in agro Nicaeni: secus arva frequens.

204. *Seriola urens*. Gen. Calyc. excl. ord. squamarum. Sem.  
omnium pappi plumosus.  
S. caul. ramoso foliolo fol. dentat. calyc. hirsid.  
ad vias et loca sterilia Comit. Nicaeni.

205. *Andriata integrifolia*  
Calyc. ex unico ordine squamarum pappi simplici.  
A. fol. inferior. vuncinat. superior. ovatis oblong. tomentis.  
in sterilibus agris Nicaeni.

206. *sinuata*.  
A. fol. vuncinatis  
secus flumen le Var in agro Nicaeni.  
Cl. Vilmor. ad praecedentem refert.

207. *Cneorum tricoccon*.  
Gen. Calyc. 3. dent. persistent. petal. 3. aequal. Stam. 3. Stigma  
3-fid. Capsula tricoeca.  
Rupes agri Nicaeni inhabitat.  
Fruticulis fol. sessil. oblong. angust. integerrima peduncul.  
solitar. 1. flor. corol. lutea Bacca rubra dein fusca

208. *Triglochin maritimum*.  
T. capsul. 6-locularibus ovatis  
locis palustribus maritimis mari proximis.

209. *Batonia tenuifolia* Tetrandria Digyn.  
Calyc. 4. phyll. persistent. foliol. altera. brevior. petala 4. calyce  
breviora angusta stigmata 2. Capsula 1-locul. 2. valv. 2. 1. pedun.  
in pascuis sterilibus Comit. Nicaeni.



210. *Alytrum maritimum* V. *Pl. Ligurica*.  
interstibus agri Nicaeensis mari proximi.

211. — *Sativum*  
Vern. Raviston sponte in agro Nicaeensi.

212. *Linaria Clypeata*  
*Alytrum* caule erecto herbaceo silicul. setis oval. compresso.  
plant. petal. acuminat. linear.  
in agro Nicaeensi.

213. — *halmiphila* V. *Pl. Vall. Macra*.  
neglecti in Comit. Nicaeensi prope Sorpello.

214. *Fossetlinia jonthlapi* silicula 1-locular. orbicul. plan. 1-sperm.  
*Clypeola* silicul. orbicul. 1-locular. 1-sperm.  
in agro Nicaeensi.

caulicul. plant. sub palm. ex rubore obscuro incana fol. sessili: altera  
exserta incana fl. terminat. racemosi pauci petal. luteola.

215. *Biscutella auriculata* Vid. *Pl. Tenu*.  
inter seget. Comit. Nicaeensi circa Sorpello.

216. *Thlaspi hirtum*  
J. silicul. sub. turdis pilosis fol. caulic. sagittat. villis  
lineatis & nicae.

217. *3beris sempervirens*  
3. frutescent fol. linear acut. integerr. fl. alto.  
in sabulosis flum. spallion Nicae.

218. — *umbellata* A  
3 herbacea fol. acuminat. inferior. serrat. superior integerr.  
in Comit. Nicaeensi.

219. *Lepidium procumbens*  
l. fol. pinnatifid. extremo major. scap. nud. caul. prostr. racemifer.  
in aridit agri Nicaeensis oditur.

220. — *spetuum*  
l. fol. pinnat. integerr. petal. emargin. calyc. minor.  
ad super locis aridis in Comit. Nicaeensi abunde.

221. *Cochlearia officinalis*  
in sparsis lapideis alpinis di. (autotica) sponte nascitur.

222. — *diva* ba  
C. fol. lanceol. dent. amplexic. incana.  
loc. incult. & sec. arv. in agro Nicaeensi.



in agro Nicaeni prope Villanica.



210. *Alytrum maritimum* V. M. L. L. L.

211.

212.

213.

214.

215.

216.

217.

218.

219.

220.

221.

222.

23. *Cram.*  
Fl.  
1-6  
Caul.  
petal.

34. *Linop.*

35. —

36. *Bra.*

37. —

38. —

39. —

40. *Hepp.*

41. —

42. *Che.*

43. —

44. —

45. —



223. *Crambe maritima*  
 Fruct. subglob. non delicens deciduus. Baccaam siccam interdum referens  
 1. locularis aut 2. locular. sed. 1. sperm. alter. locul. abortiente.  
 Caut. 3. pedat. fol. magna sinuato-pinnatifid. carnos. succulent. glauca.  
 petal. majuscula alba.  
 ad litus Nicaeense.
234. *Sinapis alba.*  
 Siliq. hispida. cort. obliq. longit. erisperm. sem. albo & ruffo.  
 loc. pingvibus in agro Nicaeensi.
235. — *maritima* V. pl. Tend.  
 in montibus Versus Nicaeam.
236. *Brassica orientalis*  
 B. fol. cord. amplexic. glob. radical. scabr. integerr. Siliq. 4. gonis.  
 in Cons. Nicaeensi. Lecta est. flor. albam prodest.
237. — *arvensis* <sup>amplexicaul. cordatis</sup>  
 B. fol. spatul. repandis sum. ~~linear. lanceol.~~ integerrimis.  
 B. campestris pinnatifida fl. purpureo foliis puberula.  
 Ad vias & muros hortorum circa Ventimiglia.
238. — *alpina*  
 B. fol. cantin. cord-sagittatis amplexic. radic. ovat. petal. erectis.  
 loci umbrosi & abrupti Cons. Nicaeensis circa la Giandola
239. — *eruca*  
 Nicaea in incultis reperitur aruga Vern. nobis
240. *Herpestis verua*  
 H. caul. erect. ramos. fol. cord. amplexicaul. serrat. villor.  
 Habitat in agro Nicaeensi
241. — *laciniata* Vid. pl. Tend.  
 Circa Soppello.
242. *Cheiranthus Borone*  
 inter Villafranca & Menton. locis aridissimis.
243. — *maritimus*  
 C. fol. lanceol. acutiusculis caul. ramos. diffus. anther. eminent.  
 Habit. in agro Nicaeensi.
244. — *litoreus*  
 Ch. fol. lanceol. subdentat. subfomentol. subcarnos. petal. emar-  
 ginat. siliq. tomentol.  
 Hae litoreis Nicaeensis indigena planta est.
245. — *sinuatus*  
 C. fol. tomentol. obtus. subinnat. ram. integerr. Siliq. mucicat.  
 in agro Nicaeensi prope Villafranca.



246. *Lythrum cheiranthoides*.  
*L. fol. lanceol. integerrimis; litiguit parulis;*  
*Mareitur in agro Nicaeensi.*
247. *Silymbrium murale*  
*S. subcaul. fol. lanceol. sinuato serrat. laeviusculi; Scappi subscabri;*  
*adscendentibus.*  
*Secus arva frequent in agro Nicaeensi.*
248. — *Vimineum*  
*S. acaul. fol. lyrat. laev. Scapp. adscendentibus fl. minut.*  
*Circa Sospitellum Onoratus Cauvini legit.*
249. — *Barvelleri*  
*S. caul. subnud. ramor. fol. radical. vuncin. dentat. hispida.*  
*Secus arva in agro Nicaeensi.*
250. *Ruta graveolens*  
*in Comit. Nicaeensi.*
251. *Euphorbia chamaerice*  
*E. dichotom. fol. crenulat. subrotund. glabr. fl. solit. axillar.*  
*caul. prostrumb.*  
*Ubique locis aridis agri Nicaeensis.*
252. — *peplis*  
*E. dichot. fol. integerr. semicordat. flor. solit. axill. caul. prostrumb.*  
*ad litus maris Nicaeensis.*
253. — *exigua*  
*E. umbella 3-fida dichot. involucl. lanceolat. fol. lineav.*  
*in agro Nicaeensi frequent.*
254. — *spinosa V. pl. figur.*  
*Ubique ad Russas in Com. Nicaeensi.*
255. — *epithymoides V. pl. figur.*  
*locis incultis & ad oras sylvarum Com. Nicaeensi.*
256. — *Nicaeensis*  
*E. glabra caul. subsimplici umbella 5-fida 2-fida fol. ovat.*  
*capitul. laevibus non sulcatis*  
*in Com. Nicaeensi inter Cimie & la Trinità.*
257. — *navalia*  
*E. umbella sub-5-fida bipida involucl. cordat. reniformibus*  
*fol. surr. imbricat.*  
*ad maris Nicaeensis litus & in sabulosis.*



long. folia

rub.

hipp.

lat. ov. folia

coral. folia

linear.

st.

da folia

da.

ren. folia

... ..  
... ..  
... ..  
... ..

... ..  
... ..  
... ..  
... ..

... ..  
... ..  
... ..  
... ..

... ..  
... ..  
... ..  
... ..

... ..  
... ..  
... ..  
... ..

... ..  
... ..  
... ..  
... ..

... ..  
... ..  
... ..  
... ..

... ..  
... ..  
... ..  
... ..

... ..  
... ..  
... ..  
... ..

... ..  
... ..  
... ..  
... ..



fol. juv. imbric. ad marii Nicaguentis limus & iſa ſabuloſit.



258. *Cappadocia pinea*  
*C. umbell. 5. fid. dichot. involucl. cordat. fol. linear.*  
*acuminat. confert. caput. leviusculi.*  
*Agri Nicæensis indigena planta est.*
259. ——— *segetalis*  
*C. umbell. 5. fid. dichotom. involucl. cordat. acut. fol.*  
*linear-lanceol. super. latioribus.*  
*in agro Nicæensi.*
260. ——— *serrata*  
*C. umbell. 5. fid. 2. fid. dichotom. involucl. diphyll. re-*  
*niform. fol. amplexicaul. cordat. serratis.*  
*ubique locis incultis agri Nicæensis.*
261. ——— *spilota*  
*C. umbell. 5. fid. 3. fid. 2. fid. involucl. ovat. petal. integr.*  
*fol. lanceol. subspilota. apice serrulatis.*  
*in Com. Nic. loc. umbrat. & montanis frequens est.*
262. ——— *myrsinites.*  
*C. umbell. sub 8. fid. 2. fid. involucl. <sup>sub</sup> ovatis fol. lanceolatis.*  
~~*sub 8. fid. 2. fid. involucl. ovatis fol. lanceolatis.*~~  
*scabris. spatulatis patentibus carnos. mucronat. margine*  
*scabris.*  
*Agri Nicæensis indigena planta est.*
263. ——— *dendroidea V. pl. Ligur.*  
*Circa Nicæam in scopulis frequens est.*
264. ——— *characias*  
*C. involucl. persiliat. emarginat. petal. obtus. fol. integerr. hirsut.*  
*umbell. multifida bifida caule pubescente.*  
*Tum in convallibus tum locis siccis agri Nicæensis ubiq. copiose.*
265. *Glaucium phœniceum.*  
*Chelidonium pedunc. 1. flor. fol. sessil. pinnatifid. caul. hirsuto.*  
*in agro nicæensi inter segetes.*
266. *Capparis spinosa*  
*Spontanea Nicæa.*
267. *Clematis erecta*  
*C. fol. pinnat. foliol. ovatis lanceol. integerr. caul. erect. flor.*  
*5. aut. 4. petalis.*  
*in collibus agri Nicæensis abunde*



268. *Clematis maritima*  
C. fol. pinnat. linearibus cauli simplic. hercagori;  
in Sylvis & locis incultis agri Nicæensis.

269. *Fumaria Capnoides*  
F. siliquis linearibus 4-gonis cauli diffus. acutangulis.  
in Montibus edioribus Comit. Nicæens.

270. ——— *spicata*  
F. parvicarpis 1.1 permis fol. lobis capillaribus.  
Jecur arva Nicæensia.

271. *Polygala monspeliaca*  
P. flor. cistatis racemos. cauli erect. fol. lanceolat-linear. acut.  
in pascuis siccis nicæensibus  
Varietatem P. Vulgaris Credit hanc Cl. Aldrov.

272. *Tritileum subterraneum*  
T. capitul. villos. 5-flor. coma central. reflexa rigida  
pact. obsolet.  
in agro Nicæensi non varum.

273. ——— *Chevleri*  
T. capitul. villos. globos. terminal. solitar. calyc. omn.  
fertil. cauli procumb. fol. obcordatis.  
Habitat in agro Nicæensi.

274. ——— *Lappaceum*  
T. spic. globos. terminal. calyc. set. hisp. rigidi; caule  
paulo foliol. ovalibus  
in Com. Nic. Jec. prata & agror. marg. & in torrent. tabulor.

275. ——— *angustifolium*  
T. spic. villos. conico oblongis dent. calycin. setaceis  
subequal. foliol. linear.  
Frequentissimum in agro Nicæensi.

276. ——— *stellatum*  
T. spic. pilos. ovar. calyc. patent. cauli diffus. foliol. obcordat.  
locis siccis Jecur arva Nicæensia frequenter.

277. ——— *scabrum*  
T. capitul. sessilibus lateral. ovar. calycibus laciniis inæq.  
rigidis recurvis  
locis siccis agri Nicæensii.



770 *Tritolium glomeratum.*

*Lemnicheae v. Vigid. sessil. calyc. striat. patul.*

mb.

col.

hem.



269. Clematis maritima  
C. fol. pinnat. linearibus cauli simplic. hexagonis,  
in sulcis & locis humilis & arboribus

269

270

271

272

273

274

275

276

277

278 Tri

279

280

281

282

283

T. spic.

284

Mel.  
Tripo  
caule  
30

285

Loh

286

287

Loh

288



- 278 *Trifolium glomeratum*.  
*T. capitul. hemisphaer. rigid. sessil. calyc. striat. patul.*  
*equal.*  
*Locis macilentis & siccis agri Nicaeensis nascitur.*
279. — *resupinatum*.  
*T. spicis subovatis. calyc. inflat. dorso gibbis. caul. prostrat.*  
*in prat. & arv. macilentis, tum siccis vias Nicaeae abunde.*
280. — *tomentosum*  
*T. spic. sessil. globos. tomentos. calyc. inflatis obtusis.*  
*etiam in agro Nicaeensi nascitur haec Trifolii species.*
281. — *fragiferum*.  
*T. spicis subrotundis. calycibus inflatis 2. dentatis reflexis*  
*caul. repentibus.*  
*in agro Nicaeensi. ad prat. marginibus sabulosa torrentium.*
282. — *procumbens*  
*T. spicis ovalibus imbricatis. versic. deflex. persistentibus. caulibus procumbentibus.*  
*Nicaeensis agri siccus est.*
283. — *filiforme*  
*T. spic. subimbric. versic. deflex. persistent. calyc. pedicell. caul. procumb.*  
*in pratibus Nicaeensibus.*
284. *Melilotus italica*.  
*Trifol. leguminibus racemosis nudis dispersis rugosis obtusis*  
*caule erecto foliol. integris.*  
*In Com. Nic.*
285. *Lobus maritimus*  
*L. legum. solitar. membranac. 4. angul. fol. glabr. bract. lanceol.*  
*in Com. Nic.*
286. — *tetragonolobus*  
*L. legum. solitar. membr. 4. angul. bracteis ovalibus*  
*inhabitat agrum Nicaeensem.*
287. ~~Lobus~~ *edulis* v. *prol. ligur.*  
*sec. arva macilentia & ad loca inculta in agro Nic.*
288. — *angustissimus*  
*L. legum. subbinatis linear. strict. erect. caul. erect. peduncul. altern.*  
*In agro Nicaeensi.*



289. *Lohet ornithopodioides*  
l. legum. subternatis acutatis compressis caulibus diffusis.  
secus arva Nicæensis abunde.

290. — *Dioicoridis*  
l. caul. erect. ramol. pedunc. subflor. legum. teret. ovalis foratis  
ad maris ripas in arenosis herbidi; prope Hospitium sic dictum  
Villafranca repertus est.

291. — *hirtula*  
l. capitul. subrotundis caul. erecto hirsuto legum. ovalis  
horis incultis & in sylvis apparet Nicæens. vulgaris.

292. — *recta*  
l. capitul. subglobos. caul. erect. laevi legum. rect. glabris.  
Nicæa ad sepes & pratorum margines.

293. — *cythroides*  
l. capitul. dimidiatis caule diffuso ramorissimo fol. tomentosis.  
in arid. Nicæensibus mari proximis.

294. *Dorychnium monspeliense*  
l. capit. asphyllis fol. sessil. 5 ual. h.  
abunde in Com. Nic.

295. *Anthyllis tetraphylla*  
A. herbacea fol. quaternopinnat. flor. lateral.  
secus arva macilenta Nicæentia.

296. — *barbajovis*  
A. pilicosa fol. pinnatis aequalibus tomentosis flor. capitulis  
ad abruptis rupes mari Nicæensi proximis.

297. *Buceva polyceraton. Trigonella?*  
Folia terrena Vereill. coroll. alis subaequale cavina brevissima  
fructus est siliqua compressa falcata.  
J. legum. subrotundis congest. erect. suberect. longil. linear. pe-  
dunc. mult.  
in agro Nicæensi.

298. — *corniculatum*  
J. legum. pedunc. congestis declinat. subfalcat. pedunculo long.  
subspinoso caule erecto  
in Agro Nicæensi.



dispositio  
hij  
die dicitur

vari  
gari

globis

fol. hinc

capituli

brevissima

linearis

edunculo long.

Prunus pennsylvanica  
L. fructu. Cerasus americana  
L. fructu. Prunella americana  
L. fructu. Prunella americana

Prunus pennsylvanica  
L. fructu. Cerasus americana  
L. fructu. Prunella americana  
L. fructu. Prunella americana

Prunus pennsylvanica  
L. fructu. Cerasus americana  
L. fructu. Prunella americana  
L. fructu. Prunella americana

Prunus pennsylvanica  
L. fructu. Cerasus americana  
L. fructu. Prunella americana  
L. fructu. Prunella americana

Prunus pennsylvanica  
L. fructu. Cerasus americana  
L. fructu. Prunella americana  
L. fructu. Prunella americana

Prunus pennsylvanica  
L. fructu. Cerasus americana  
L. fructu. Prunella americana  
L. fructu. Prunella americana

Prunus pennsylvanica  
L. fructu. Cerasus americana  
L. fructu. Prunella americana  
L. fructu. Prunella americana

Prunus pennsylvanica  
L. fructu. Cerasus americana  
L. fructu. Prunella americana  
L. fructu. Prunella americana

Prunus pennsylvanica  
L. fructu. Cerasus americana  
L. fructu. Prunella americana  
L. fructu. Prunella americana

Prunus pennsylvanica  
L. fructu. Cerasus americana  
L. fructu. Prunella americana  
L. fructu. Prunella americana

Prunus pennsylvanica  
L. fructu. Cerasus americana  
L. fructu. Prunella americana  
L. fructu. Prunella americana



229. *Leptocarpus* ...  
F. legum. ...  
...

230. *Didymopanax* ...  
F. caul. ...  
...

231. *Didymopanax* ...  
F. caul. ...  
...

232. *Didymopanax* ...  
F. caul. ...  
...

233. *Didymopanax* ...  
F. caul. ...  
...

234. *Didymopanax* ...  
F. caul. ...  
...

235. *Didymopanax* ...  
F. caul. ...  
...

236. *Didymopanax* ...  
F. caul. ...  
...

237. *Buceca* ...  
F. caul. ...  
...

238. *Buceca* ...  
F. caul. ...  
...



299. *Bucera monipeliacum*

J. leg. sessil. congest. arcuatis divaricatis inclinatis bre.  
vibus peduncul. mucronato inermi.  
in sterilibus agris Nicaeni.

300. ——— *forium graecum.*

J. legum. sessil. strict. erectiscent. subsulcatis acuminat. caule  
bas. ramoso stolonifero  
ad agrorum margines in Com. Nic.

301. *Medicago radiata*

Ch. leguminibus veniformibus marginedentatis fol. ternatis.  
in agro Nicaeni.

302. ——— *orbicularis*

Ch. legum. solitar. cochleatis depressis planis stipul. ciliatis. caud. diffus.  
Copiose sedes agris in Com. Nicaeni.

303. ——— *scutellata.*

Ch. cochleata major dicarpos capsula rotunda globosa scutellata.  
Frequens in Com. Nicaeni. inter seget. & secus arva.

304. ——— *interversa.*

Ch. cochl. major. spinosa dicarpos capsulae spinis longioribus sursum  
& deorsum tendentibus  
Haec etiam Nicaeni agro indigena est.

305. ——— *arabica*

Ch. cochl. minor polycarpos annua capsula majore alba folio  
cordato macula fusca.  
In agro Nicaeni inter segetes.

306. ——— *lurbinata*

Ch. legum. cochl. inerm. subcylindr. eschreum. complanatis.  
inter segetes in agro Nicaeni.

307. ——— *ciliaris*

Ch. legumin. congest. globos. hispida. stipul. ciliatis.  
Cum praecedentibus in eodem locis.

308. ——— *virgula*

Ch. 3-phyll. legum. cochl. spinosa foliol. inferior. cuneiform. rem.  
superioribus subrotundis.  
In agro Nicaeni.

309. ——— *muricata*

Ch. Cochl. dicarpos capsula spinosa rotunda minore  
etiam in agro Nic. nascitur.



310. *Medicago laciniosa*  
*Med. legum. coroll. spinos. stipul. contort. fol. oblong. duplicat. serratis.*  
 In agro Nicaeensi.
311. ——— *marina*. V. pl. figur.  
 in sterilibus litor. maris proximis.
312. *Ononis antiquorum*.  
*O. flor. solitar. foliol. major. fol. inferior. ternat. ram. laeviuscul.*  
*spinol.*  
 in agro Nicaeensi.
313. ——— *Cherleri*  
*O. pedunc. 1. flor. avistatis fol. ternat. stipulis serratis.*  
 In Jussulor. & ad Rupe. Villae Francae Nicaeensi.
314. ——— *Vitcola*  
*O. pedunc. 1. flor. avistatis fol. simplic. infim. ternat.*  
 locis aridis circa Menton et inter Roche Tallade & Biot.
315. ——— *pinguis*  
*O. pedunc. 1. flor. avistatis fol. ternat. lanceolat. stipul. integerr.*  
 in collibus Nicaeensibus.
316. *Cytisus lupinus* V. pl. Tenu.  
 inter Jussello & Lantoca.
317. ——— *gracilis*  
*C. fol. simplic. lanceolat. lineari. ramis angulatis.*  
 Nicae exemplar lectum habet Cl. Allione.
318. *Genista sagittalis*  
*G. ramis arispidibus membranac. articulatis fol. ovato-lanceolatis.*  
 inter Jussello & Lantoca.
319. ——— *hispanica*  
*G. spinis decompositis ram. flor. fer. inerm. fol. lineari pilol.*  
 in aridis apricis Com. etc.  
 Dubitat. Cl. Allione num vere a Germanica differat.
320. ——— *candicans*  
*G. fol. ternat. subt. villor. peduncul. lateral. sub. 5. flor. foliol. legumin. hirsutis.*  
 in montibus Nicaeensibus calidioribus.
321. *Hedysarum spinosissimum* V. pl. figur.  
 in agro Nicaeensi loc. Sabator. mari prope. & securaria.



322. *Orobrychi*, *Sarcobiti* N. pl. figur.  
in *Larv. T. p. 100*. Com. nicaens.

Var. latifolia

af.

1/2001.

Trade & B

1/2 post-integ

1911.

fol. 65a

total linear vol.

2:45 PM

*foliat. e.*

Accuracy



310. *Medicago laciniata*

M. legum. corb. minor. Rhinul. can. fol. oblong. duplicat. 1/2 v. 1

311. *Ononis asperifolia*

3. fol. latior. corb. major. fol. asperifolia. can. 1/2 v. 1

312. *Ononis asperifolia*

3. fol. latior. corb. major. fol. asperifolia. can. 1/2 v. 1

313. *Ononis asperifolia*

3. fol. latior. corb. major. fol. asperifolia. can. 1/2 v. 1

314. *Ononis asperifolia*

3. fol. latior. corb. major. fol. asperifolia. can. 1/2 v. 1

315. *Ononis asperifolia*

3. fol. latior. corb. major. fol. asperifolia. can. 1/2 v. 1

316. *Ononis asperifolia*

3. fol. latior. corb. major. fol. asperifolia. can. 1/2 v. 1

317. *Ononis asperifolia*

3. fol. latior. corb. major. fol. asperifolia. can. 1/2 v. 1

318. *Ononis asperifolia*

3. fol. latior. corb. major. fol. asperifolia. can. 1/2 v. 1

319. *Ononis asperifolia*

3. fol. latior. corb. major. fol. asperifolia. can. 1/2 v. 1

320. *Ononis asperifolia*

3. fol. latior. corb. major. fol. asperifolia. can. 1/2 v. 1



322. *Onobrychis sarcocolla* V. pl. figur.  
in agro Nicæensi communis.

323. — caput galli  
*Medicago*. fol. pinnatis legum. monosperm. ciliat. aculeat.  
pet. equal. caul. diffus.  
ad Colles aridos & sabulosa torrentium in agro Nicæensi.

324. *Vicia Buthynica*  
V. legum. pedunc. solit. erect. foliol. quaternis ovato-lanceol.  
hiput. dentatis.  
Nicæensi agri indigena planta est.

325. — *Lathyrus*  
V. leg. sessil. solitar. erect. glabr. foliol. den. inferior. obcor-  
dat.  
in agro Nicæensi.

326. — *peregrina*  
V. leg. subsessil. pendul. glabr. 4. sperm. foliol. linear. emarginat.  
inter seget. agri Nicæensi.

327. *Ervum Monanthos*  
l. pedunc. uniflor.  
agr. Nic. indigena planta est.

328. — *erilia*  
l. germin. undat plicat. fol. impar. pinnat.  
inter seget. Agr. Nic. occurrat.

329. *Lathyrus Cicera*  
l. pedunc. uniflor. circ. 2. phyll. legum. ovat. compress.  
obso. canaliculatis  
in agro Nicæensi. spontan.

330. — *laticus*  
l. pedunc. 1. flor. circ. 2. phyll. 4. phyll. legum. ovat. compress.  
obso. bimariginatis.  
etiam Nic. agri indigenus.

331. — *setifolius*  
l. pedunc. 1. flor. circ. 2. phyll. foliol. setac-linear.

332. — *angulatus*  
l. ped. 1. flor. aristat. circ. 2. phyll. simpliciss. foliol. linear.  
Vulgaris in agro Nicæensi.

333. — *heterophyllus*  
l. pedunc. multiflor. circ. 2. phyll. 4. phyll. foliol. lanceolat. internod.  
membranaceis.  
in umbris di lucevane.



334. *Pisum maritimum*  
*P. petiol. supra planiusculis caul. angulat. stipul. sagit.*  
*tal. peduncul. 1. flor.*  
*In agro Nicaensi nascitur.*
335. — *ochrus*  
*P. petiol. decurrentibus membranac. diphyllis pedunc. 1. flor.*  
*inter segetes Nicaenses.*
336. *Cotonilla juncea*  
*C. fruticosa fol. pinnat ternatisq. lineav-lanceolat. subcartis.*  
*obtus.*  
*in agro Nicaensi*
337. — *Valentina*  
*C. fruticosa fol. subovatis, stipul. suborbiculatis.*  
*in colle di Montalbano.*
338. — *glauca*  
*C. fruticosa foliol. suborbiculatis. stipul. ovatis. pedunc. complanatis.*  
*in agro Nicaensi ad super.*
339. — *securidata*  
*C. herbacea legum. falcato-glabriatis foliol. plurimis.*  
*in agro Nicaensi vulgariter est.*
340. *Ornithopodium compressum.*  
*O. fol. pinnat. legum. recurvat. compress. rugos. bract. pinnat.*  
*in agro Nicaensi.*
341. — *scorpioides* *N. pl. figur.*  
*Securaria macilentia agri Nicaensis.*
342. *Scorpiurus vermiculata*  
*Gen. Calice. Semi-5-fidus dentibus unguis minus divitis Vaseillum*  
*reflexum subrotundum emarginatum patens ala oblonga laeva Carina*  
*recta gibba ungue diviso Siliq. teres in se contort. articulat. rugos. vel hispida.*  
*J. peduncul. 1. flor. legum. techis axillig. squam. obtus.*  
*inter segetes & ad vias in agro Nicaensi.*
343. — *muricata*  
*J. pedunc. 2. flor. legum. erectissimum obtuse aculeatis*  
*similibus locis prioris nascitur.*
344. — *subvillota*  
*J. pedunc. sub 4. flor legum. erectiss. spin. confertis erectis.*  
*Nicaentem agrum incolit.*



*Alipat*

*ne. l. f. h.*

*h. h. f. h.*

*comp. h.*

*mit.*

*black. pinn.*

*Vanillan  
laser. Catina  
1911. vel. h. h.  
h. h.*

*len. h.*

*ethi. v. h.*



*[Faint, illegible handwriting, likely bleed-through from the reverse side of the page]*



Handwritten text at the top of the page, possibly a title or introductory paragraph.

Second block of handwritten text, continuing the narrative or list.

Third block of handwritten text, appearing as a distinct entry or section.

Fourth block of handwritten text, continuing the sequence.

Fifth block of handwritten text, showing further detail or description.

Sixth block of handwritten text, possibly a conclusion or summary for a section.

Seventh block of handwritten text, continuing the main body of the document.

Eighth block of handwritten text, showing a transition or new point.

Ninth block of handwritten text, continuing the flow of information.

Tenth block of handwritten text, possibly a final note or signature area.

Eleventh block of handwritten text at the bottom of the page.



345. Hipp  
Gen.  
inter  
intra  
H.

346 —

347. Spont

348 —

349. / M  
Gen.  
margin  
long  
Trib  
in

350. Ceter

351. Ana  
Gen. P  
Ala ol  
omni d  
Ci  
poffente  
legere  
utile. m  
volut.

352. Alva  
d. can  
obcord



345. *Hippocrepis unilicquola*  
 Gen. Essentialis. Character consistit in siliqua compressa longa incurva tubera  
 inferiore ad medium usq. inter sem. sinuata solea & quinque ad instar se-  
 mina incurva.  
 H. legum. sessil. solitar. vact.  
 Abunde in Com. Nicæensi.

346. ——— *multilicquola*  
 H. legum. pedunculatis confertis <sup>circularibus</sup> ~~areolatis~~ marg. <sup>altero lobali</sup> ~~retusis~~ <sup>reflexis</sup>.  
 Etiam haec sed rarior in agro Nic. occurrit.

347. *Spartium purgans*  
 J. ram. teret. striat. fol. lanceol. sessil. pubesc.  
 freq. est. in mont. calidior. Com. Nicæensi.

348. ——— *spinosum*  
 J. fol. ternat. ram. angulat. spinos.  
 in Com. Nic. uberrime crevit.

349. *Psoralea bituminosa*  
 Gen. Calyce. 1. phyll. 5. fid. irregul. tubercul. asperus. Vercill. subrotund.  
 emarginat. alturgens. alae oblongae parvae. Carina dipetalata lunulata siliqua  
 breviter calyce contenta semen 1. aut 2. contin.  
*Trifolium bitumen redolens*  
 in Comit. Nicæensi ubique.

350. *Cercis siliquastrum*  
 C. fol. cord. orbiculatis glabris  
 frequent in sylvis et ad rupes in agro Nicæensi.

351. *Anagyris foetida*.  
 Gen. Calyce. campanul. 5. dentat. irregul. Vercill. breviss. vact. ann.  
 Alae oblongae Vercillo longiores Carina 2. petal. longissima Hamina  
 omnia distincta siliqua teretiuscula magna semina reniformia.  
 Circa Nicæam ad rupes  
 Jussente fol. altern. petiol. ternata foliol. sessil. oblong. acuminat.  
 laeviss. supra virid. subt. albicans. odor. grav. & fetid. pedunc.  
 arbor. multifi. binis - ternis - quinis oblongis Cal. 5. fid. glauc.  
 & velut argenteae Vercill. aureum coctus. petala pallidiora.

352. *Alvagalus hamolus*.  
 A. caulescens procumbens legum. tubulat. recurvat. glabris foliol.  
 obcordat. subtus villosis  
 Locis siccioribus Collium Nicæensium.



353. *Bilevula pelecini* L.  
 Gen. Callyx monophyllus semib. fid. irregulari. Vexillum majus late-  
 ribus reflexis alae oblongae vexillo breviores Cavina oblonga siliqua com-  
 pressa 2-loculari septo valvi contrario semina reniformia.  
 Nicotia inter segetes non rara.  
 Caul. herbac. ramos. fol. altern. imparipinn. foliol. mult. opposita  
 subseriat. obcordat. integerr. flor. axill. solitarii peduncul. dimidio  
 capituli rubelli.

354. *Eragium maritimum* V. pl. lign. v.  
 in sabulo ad mare litor.

355. *Echinophora spinosa* L.  
 Gen. involucri pance. folior. involucri. 1-phylli. multif. flos. radiat.  
 pet. inaequal. sem. sterilia coronata in involucrio indurata.  
 E. foliol. subulato-spinosis integerrimis.  
 frequent ad litor. Maris Nicotensis.  
 Caul. er. b. ramos. striatus subullos. fol. altern. 2-pinnatif. subullosa  
 lacin. linear. apice spinos. pedunc. later. subullos. pet. alb. sub. pur.  
 purascentia.

356. *Pastinaca oppositifolia* V. pl. lign. v.  
 in agro Nicotensi locis umbratis & montibus superstitibus.

357. *Ferula communis*. Gen. Semen. magn. oblong-ovate 3-striat. cinch.  
 ala magna solida non membranacea.  
 F. foliol. linear. longior. simplicibus.  
 In agro Nicotensi nascitur.  
 Albidus humana radice crassa lactescens fol. alterna. amplectis.  
 2-pinnat. umbell. robust. petal. parva reflexa lutea.

358. — nodiflora V. pl. lign. v.  
 in Monte Joet Comit. Nicotensis.

359. *Pencedanum minus* V. pl. Tenu.  
 inter la Ghiaudola & la Scavena.

360. *Selinum carvifolia* Chabraz  
 1-fol. multiplic. radical. circa cortem decussatis  
 in pratis Nicotensibus



361. *Thapsia villosa* Gen. Flor. aequal. petal. integri lanceolati; invo-  
lucrum nullum. Sem. oblong. 3-lobatum magna ala circumscriptum



353. *Bilevula pelecini* L.

Gen. *Callix* monophyllus semib. p. d. irregulari; *Urcillum* majus late.

35

3

35

35

35

35

36



361. *Thapsia villosa* Gen. Flor. opposit. petal. integris lanceolatis involu-  
crum nullum. Sem. oblong. 3-lobatum magna ala circumscripta su-  
pra & infra ampliter truncata est & emarginata.  
T. foliol. dentat. Villis bar. coadunatis

in Agro Nicotenti.  
Caul. ca. bit. & ultra brachiatus fol. crassiusculis. & aspera in ala  
oppositis subjecta sylvant. longior. semunciat. umb. lat. pet. lutea.

362. *Lactopitium Gallicum* V. pl. Vall. Mur.  
inter Tenda & Eleavena.

363. *Ligusticum spelopponeticum*.  
l. fol. multiplicato pinnatis foliol. pinnatis incisis.  
in com. Nicotenti. circa Soppello & La Ghianola.  
invol. unis. 5-6. phyll. parial. 2. aut 3. phyll. petal. loddid. pallid.  
Sem. glabr. oblong. non rostrat. profund. sulcat. live luteat.

364. *Coriandrum testiculatum*. V. pl. lignv.  
in agr. com. Nicotenti.

365. *Pimpinella peregrina*  
p. fol. radica. pinnatis crenat. sum. cuneiform. incisis.  
l. Belhard: in sterilibus Nicotenti. bus ad prohem d. pimpia.

366. *Seseli annuum*  
l. petiol. ramis. membranac. ventricol. emarginat.  
in collibus com. Nicotenti. frequens est.

367. — *glaucum*  
l. petiol. ramis. membranac. oblong. integris foliol.  
singulis binatisq. canaliculatis. latius petiole longior.  
in collibus aridis agr. Nicotenti.

368. — *tormentosum*  
l. caul. alto rigido foliol. linear. fasciculat.  
inter Lucerne & La Scarenna ad vupes.

*Seseli genus* flores hab. florulas. pedicels involucri. parial. poly-  
phyll. universale vix ullum aut o. petala cordata reflexa fructus  
subrotundus aut ovatus. Sem. ovat. acuminat. intus planum gibbum  
h. corbatur.

369. *Oenanthe globulosa* Vid. Gen. ad pl. lignv.  
o. fructibus globosis  
in herb. d. & sub hum. agr. Nicotenti.

370. — *pimpinellodes* V. pl. lignv.  
par. loc. sub hum. d. et graminos. in com. Nicotenti.



371. *Supplementum volundifol.*  
in agro Nicaeensi collium.

372. — *Jemiconoposium*  
p. caule herbaceo fol. lanceol. umbell. acillar. terminalibus.  
sem. scabris.  
Circa Villa Parva in sterilibus vasis Saint Olyce.

373. — *juncum* v. p. l. ligur.  
Habitat in agro & Com. Nicaensi.

374. — *temissimum*  
p. fol. angustissim. umbell. simp. altern. 5-phyl. sub 3-flor.  
in agro Nicaensi.

375. *Carum Bunium*  
Gen. involucri. utriusq. deest petal. cord. aeq. incis. sem. oval. profund.  
striata intus plana.  
Cum bell. & p. d. fol. radical. bipinnat. linear. cauliu. sub erectis.  
in Vineis Savorgii & Circa Lucerne.

376. *Sison amomum*. Involucri. p. n. c. fol. sub 4-phyl. petal. cordat.  
aequal. p. oval. oblong. parva sem. sub bil. ter striata.  
*Carum Bunium* & *Sison* ad finem sunt gen. ut agere disting. possint.  
p. fol. pinnat. umbell. erect.  
in Com. Nicaensi.

377. *Cistium maritimum* v. p. l. ligur.  
in muris v. v. h. Com. Nicaensi.

378. *Sium falcaria*.  
Gen. fl. aeq. f. v. h. petal. aequal. cordat. p. u. c. oblong-oval.  
sem. oval. leuiter striata intus plan. calyc. minim. coronata.  
p. foliolinear. decurrent. cennatis.  
in agro Nicaensi.

379. — *nodiflorum*  
p. fol. pinnat. umbell. acillar. sessil.  
Cl. Bellard: prope Sappella ad muros sublimidos.

380. *Myrrhis australis*  
p. sem. tubulat. bipid. flor. radical. caul. h. v. b. u.  
in agro collium Nicaeentium.

381. *Daucus mauritanicus*  
p. sem. bipid. flor. c. v. h. ster. carnol. receptacul.  
Comm. hemisphaerico  
in p. v. h. agro Nicaeentium.



382. *Donner leippanent.*  
*et leippanentis cen wal stetit: nullo cane lucido.*



371. *Supplementum volucrium.*  
in aq. *Nicotianum* collum.



382. *Daucus hispanicus*.  
D. sem. aculeatis centrat. sterit. nullo caule lucido.  
in agro Nicasensi.

383. *Caucalis latifolia*  
Involucr. exc. foliol. non pinnatifid. sed integ. v. paucifol. quod  
interdum deest sem. striat. aut costat. fect. aculeis aut setis virgidis  
uncinatis.  
C. umbel. unives. 3-fida partial. 5-sperm. fol. pinn. serratis.  
in agro collium Nicasensium.

384. ——— *maritima*  
C. umbel. 2-fid. umbellat. 2-sperm. involucr. semine  
brevior. linear.  
in agro Nicasensi.

385. ——— *nodiflora*  
*Tardylum* umbell. simplic. setit. sem. exterior. h. 7 sp. 2:1.  
ad limit. agror. & in palust. lacor. C. Nicasens. frequens.

386. ——— *nodosa*. V. sp. l. qu.  
in vineis & sterit. limit. Nicasens.

387. *Geranium ciconium*  
G. pedunc. multiflor. 5-andr. fol. pinnat. pinnatif.  
obtus.  
Nicasa ad semitas.

388. ——— *moschatum*.  
G. ped. multif. flor. pentandr. fol. pinnat. incis. cotyledon.  
pinnatifidit.  
Nicasa reperitur

389. ——— *malacoides*  
G. ped. multiflor. flor. pent. fol. cordat. sublobat.  
Circa Nicasam videtur.

390. ——— *maritimum*  
G. pedunc. sub 3. flor. flor. pentandr. fol. cordat. incis.  
crenatis scabris caul. procumbentibus.  
Secus vias Nicasae. ad loca maris proxima.

391. ——— *lucidum*  
G. ped. 2-flor. calyc. pyramidal. angulat. elevato rugosis  
fol. 5-lobis robustioribus  
in agro Nicasensi.



392. *Malva parviflora*  
Calyce exterior 3phyllus interior semi 5. fidus.  
Ch. caule patulo fol. angulat. flor. axillar. sessil. glo-  
meratis. Calyc. glabri patentibus.  
in Comit. Nicæensi.

393. ——— *Nicaeensis.*  
Ch. caul. pilos. decumbent. herbar. fol. subcordat. 5. lobis acutis.  
pedunc. erectis.  
Cl. Bellardi in agro Nicæensi prope portum di. Rimpia.

394. *Lavatera arborea.*  
Differt. Genus hoc. a *Malva* calyc. exterior. 3. fido  
non 3phyllus.  
l. caule arboreo fol. 7. angulat. tomentos. plicatis.  
pedunc. confert. 1. flor. axillaribus.  
Reperitur ad rupes litus Nicæensis.

395. ——— *olbia*  
l. caul. pilos. fol. 5. lob. dentat. pedunc. solitari.  
Nicæensem agrum inhabitat.

396. ——— *thuringiaca*  
l. caule herbaceo fruct. denudatis calyc. incisis.  
in Comit. Nicæensi.

397. ——— *trimestris*  
l. fol. glabris caule scabro herbaceo pedunc. 1. flor.  
fruct. orbicul. sect.  
in olivis Villafranca Nicæensi reperitur.

398. ——— *maxima.*  
l. caule pilosissimo lanato fol. cordat. orbiculat. 5. lob. obtus. crenat.  
pedunc. 1. flor. stipulis setaceis deciduis.  
ad rupes litus & agri Nicæensis abunde.

399. *Alcea Rosea*. Calycem exterior. habet. semi 6. fidum.  
A. fol. sinuatis angulosis  
in Comitum Nicæensi.



on *Althea hirsuta* Cal. erect. plumbeous 6. fid.  
*A. hirsuta* Cal. erect. plumbeous 6. fid.

Ind.

Nov. Jupit. glo.

nt. 5-106/ acub

no 2: *limp*ia.

interior. 3. hido

*mentes plicat.*

1917.

предмет. 10/1/7

ly calyc. inc./

danceal. 1. flor.

*[Faint handwritten text at bottom left]*

*6 pul. crenat.*

*G. fidum.*



392. *Malva parviflora*  
C. ... .. h. ...

39

39

39

396

397

398

399



400. *Althaea hirsuta* Cal. erect. pl. usquam 5. fid.  
A. fol. inter. luteo-rosea. in c. 70. crenat. ceter. 3. fid.  
inhabitat super & Colles Com. Nicaeensis.

401. *Hypericum tomentosum*.  
H. fl. trigynis; calyc. serratis - glandulosis. fol. semi am-  
plexicaulis. flexuos. tomentos. caul. prostratis.  
ad frutices rivulos & aquaeductus in agro Nicaeensi.

402. *Croton tinctorium*. Monoezia monadelphica.  
Gen. Masc. calyc. cylindr. 5. dentat. coroll. 5. petal. Stam.  
10-15.  
Fem. Cal. polyphyll. Coroll. 0. Styl. 3. bifid. Capsul. 3. locul.  
sem. 1.  
C. fol. rhombis repandis capsul. pendul. caul. herbac.  
in Agro Nicaeensi frequens.  
Planta tenella pulverulenta caul. cubital. ramis. fol. altern.  
long. petiol. sinuata nervosa fl. axillar. congesti pedunculat.  
calyc. 5. phyll. petal. lanceol.

403. *Ranunculus muricatus*  
R. sem. aculeatis fol. simpl. lobat. obtus. glabr. caul. diffus.  
in agro Nicaeensi.

404. ——— *parviflorus*. V. fl. ligur.  
in arvis subhumid. Com. Nicaeensis.

405. *Corrigiola litoralis* V. fl. lig.  
locis arenosis mar. proximis.

406. *Delphinium peregrinum*  
D. nectar. diphyllis flor. solitar. fol. multipartit. foliol.  
linear. acuminatis.  
Circa Nicaeam nascitur.

407. ——— *Staphysagria*  
D. nectar. 4. phyll. petal. brevior. fol. palmat. lob. obtus.  
Reperitur Nicaea.

408. *Nigella damascena*  
Agri Nicaeensis familiaris planta.



409. *Hebeborus niger*.  
H. scapo subbifloro subundo fol. pedat. / flore albo  
in frigidioribus Nicodemi montibus.

410. *Sarcopraga granulata*  
S. fol. caul. n. reniform. lobat. caul. ramor. radic. granulata.  
Locis ruperibus Com. Nicodemi.

411. *Saponaria Vaccaria*  
S. calyc. pyramidalis 5. angulat. fol. ovat. acumin. sessil.  
in Com. Nicodemi. inter leges.

412. *Velexia rigida*. Calyc. filiform. cylindr. 5. dent. Corolla parva  
5. petala Stam. 5. vel 6. Styl. 2. Caput. 1. locut. cylindr. polyperm.  
Stam. in simplic. serbe.  
in herbis ad limites agrorum Nicodemi.

Caul. desquissulcat. Vannosill. duri. fol. connat. tubulat. integ.  
pilosa. fl. axillar. solitar. sessil. Corol. limbus minimus.

413. *Silene anglica*  
S. hirsuta petalis emarginatis flor. erect. fruct. 6. reflex. peduncul.  
altern.  
Secus arva Nicodemi.

414. ——— *ceratoides*  
S. hirs. pet. emarg. fruct. 6. erect. calyc. subpilos. sessil.  
Medic. Canvini in agro Nicodemi legib.

415. ——— *S. Vulnera*  
S. petal. integ. subrotund. fruct. erect. altern.  
Secus arva Nicodemi.

416. ——— *nocturna*  
S. flor. spicat. altern. secund. sessil. pet. bifid.  
in agro Nicodemi. Oliveb. partem occupat.

417. ——— *conoides*.  
S. calyc. fruct. globos. acuminat. Styl. 30. fol. glab. petal. integ.  
Nativus in agro Nicodemi.

418. ——— *noctiflora*  
S. calyc. 10. angul. dent. tub. aequant. caul. dichot. petal. bifid.  
in Com. Nicodemi.

419. ——— *Nicodemi*  
S. v. hor. & v. hor. fol. linear. obtus. petal. renib. fruct. ovat. calyc. 10. fl.  
in agro Nicodemi. Secus le vas frequenter.



olla

annulata.

Junil.

Corolla  
cylindric. ped

annulab. int  
unipinnatif.

reflex. ped

Junil.

loba petal. 5

petal. 6

loba ovat. caly 10  
ob.

prima...



400. *Silene acaulis*  
Caulis prostratus in arenis & peltis. Petala  
in brevibus cymis cernuis.

401. *Silene acaulis*  
Caulis prostratus in arenis & peltis. Petala  
in brevibus cymis cernuis.

402. *Silene acaulis*  
Caulis prostratus in arenis & peltis. Petala  
in brevibus cymis cernuis.

403. *Valeriana rigida*. Calyx filiformis. Lilla de 5 dent. Petala  
in brevibus cymis cernuis. Stylis 2. Capsula bilobata. Lilla  
in brevibus cymis cernuis.

404. *Valeriana rigida*. Calyx filiformis. Lilla de 5 dent. Petala  
in brevibus cymis cernuis. Stylis 2. Capsula bilobata. Lilla  
in brevibus cymis cernuis.

405. *Silene acaulis*  
Caulis prostratus in arenis & peltis. Petala  
in brevibus cymis cernuis.

406. *Silene acaulis*  
Caulis prostratus in arenis & peltis. Petala  
in brevibus cymis cernuis.

407. *Silene acaulis*  
Caulis prostratus in arenis & peltis. Petala  
in brevibus cymis cernuis.

408. *Silene acaulis*  
Caulis prostratus in arenis & peltis. Petala  
in brevibus cymis cernuis.

409. *Silene acaulis*  
Caulis prostratus in arenis & peltis. Petala  
in brevibus cymis cernuis.

410. *Silene acaulis*  
Caulis prostratus in arenis & peltis. Petala  
in brevibus cymis cernuis.

411. *Silene acaulis*  
Caulis prostratus in arenis & peltis. Petala  
in brevibus cymis cernuis.

400. *Silene acaulis*  
Caulis prostratus in arenis & peltis. Petala  
in brevibus cymis cernuis.

401. *Silene acaulis*  
Caulis prostratus in arenis & peltis. Petala  
in brevibus cymis cernuis.

402. *Silene acaulis*  
Caulis prostratus in arenis & peltis. Petala  
in brevibus cymis cernuis.

403. *Valeriana rigida*. Calyx filiformis. Lilla de 5 dent. Petala  
in brevibus cymis cernuis. Stylis 2. Capsula bilobata. Lilla  
in brevibus cymis cernuis.

404. *Valeriana rigida*. Calyx filiformis. Lilla de 5 dent. Petala  
in brevibus cymis cernuis. Stylis 2. Capsula bilobata. Lilla  
in brevibus cymis cernuis.

405. *Silene acaulis*  
Caulis prostratus in arenis & peltis. Petala  
in brevibus cymis cernuis.

406. *Silene acaulis*  
Caulis prostratus in arenis & peltis. Petala  
in brevibus cymis cernuis.

407. *Silene acaulis*  
Caulis prostratus in arenis & peltis. Petala  
in brevibus cymis cernuis.

408. *Silene acaulis*  
Caulis prostratus in arenis & peltis. Petala  
in brevibus cymis cernuis.

409. *Silene acaulis*  
Caulis prostratus in arenis & peltis. Petala  
in brevibus cymis cernuis.

410. *Silene acaulis*  
Caulis prostratus in arenis & peltis. Petala  
in brevibus cymis cernuis.

411. *Silene acaulis*  
Caulis prostratus in arenis & peltis. Petala  
in brevibus cymis cernuis.



420. *Silene inappetita*.  
l. caul. dichotom. panicul. calyc. laev. petal. breviss. emargin. fol.  
glabr. lanceol.  
Haec etiam Nicæensis agri indigena est.

421. *Agrostemma ciliolata*  
A. glabra fol. linear. lanceol. petal. emargin. coronat.  
in Comitatu Nicæensi.

422. *Tamariacus gallicus*  
T. flor. pentandr. fol. viridib. tenuibus  
in agro Nicæensi.

423. *Statice limonium*  
l. scapo paniculato breviter fol. laev. succ. sub mucronat.  
agrum Nicæensem ad maris litus inhabitat.

424. — *cordata*  
l. scapo panicul. fol. spatul. velut.  
ad vases maritimos frequens.

425. *Frankenia pulverulenta*.  
calyc. subcylindricus 5-dent. persistent. l. am. 6. petal. 5. styl. 1.  
stigmat. bifido caput. 1. locul. 2. im. adum. in 3. valvul. dehisc.  
polyperma.  
f. f. obovat. velut. sub pulverat.  
in agris Nicæens. arenos.  
caul. humifus. palmat. ramor. rament. fol. oppos. vel. quatern. fascicul. erigui  
amæthistini.

426. *Rereda alba*  
a. fol. pinnat. foliis tetragynis calyc. serapantib.  
in C. Nicæens. locis sterilibus & arenosis.

427. *Tribulus terrestris* V. pl. ligur.  
Nicæa seu arva abunde.

428. *Halmala multifida*. *Pegannum* L.  
cal. 5. phyll. coroll. 5. petal. aequal. l. am. 15. styl. 3. bif. caput. 3. locul.  
polyperma.  
p. fol. multifidi.  
in agro Nicæensi nascitur.  
caul. cubital. ramor. /wizhu. fol. alterna sessil. succulenta flor.  
oppositifol. solitar. pedunculati fol. Calycin. linear. 3. longior.  
petal. luteo virid.



429. **Cistus**. monspeliensis.

C. arboreus, erectus. fol. lineari-lanceol. sessil. utrinq. villor.  
3-nerv.  
Freg. in arid. incult. et Jaceo. circa Villanueva Copposill.  
circa Berra in Com. Nicæni.

430

— **salvifolius**.  
C. arbor. erectus. fol. ovato. petiolat. utrinq. hirsutis.  
in agro Nicæni ad rupes aridas.

431

— **incanul.**  
C. arb. erectus. fol. spatul. toment. rugor. inferior. basi va-  
ginant. connat.  
in Com. Nicæni. non in freg.

432

— **medius**  
C. fraxinea, folio Salviae flore ochrea colore  
in agro Nicæni.  
Caul. rubent. villos. pedunc. solitari. axillar. & terminal.  
fol. petiol. multum rugor. virid. ovato-lanceol. in ambiguentibus

433

— **albidus** V. pl. liquor.  
in agro Nicæni ad rupes aridas.

434

— **crispus**  
C. arbor. erectus. fol. lanceol. pubesc. 3-nerv. undul.  
in Comit. Nicæni.

435

— **laevipes**.  
C. suffrutescens, adscendens, erectus. fol. altern. falcatis.  
filiform. glabr. pedunc. racemos.  
in agro Nicæni frequent.

436

— **canus**  
C. suffrutescens. erectus. procumbens. fol. opposit. obovat.  
villor. subt. tomentor. fl. subumbellat.  
Habitat in C. Nicæni.

437

— **italicus**.  
C. suffrut. erectus. fol. oppos. hirsut. inferior. ovato. superior.  
lanceolat. v. patent.  
in rupibus circa Lucerame.



38 Cithus tuberosaria  
C. eschschol. perenn. fol. radic. ovat. 3. nerv. toment. caul. glabr. lanceol. lumen. alterni.

ultra. v. l. l.

dist. ab.

non. b. p. v.

terminal.  
in ambiguitate

pal.

non. fascicul.

non. ovat.

non. superior.

at.

ill.

lin.

lyper.



429. *Cistus*. *monipellensis*.

(*C. arboreus*) *exhispul.* fol. lineari-lanceol. sessil. utrinq. villis.

430

431

432

433

434

435

436

437



438. *Cistus tuberosus*  
C. erectus. perenn. fol. rad. ov. 3. nerv. toment. caul. glabr. lanceol. lumen. alterni.  
in Com. Nicaeni. Sterilib. & ruppert. frequenter  
circa Nicaenam.

439. ——— *guttatus*. V. pl. lign. in palen. sicis Sterilib. Nicaeni. Comit.

440. ——— *ledifolius*  
C. herbac. glaber stipulat. flor. solitar. suberit. fol. ternat. opposit.  
in Com. Nicaeni.

441. ——— *laticifolius*  
C. herbac. palat. villos. stipulat. flor. racemor. erect. pedunc. horizont.  
in Com. Nicaeni.

442. ——— *thymifolius*  
C. suffrutic. stipul. procumb. fol. oval. linear. oppos. breviss. congest.  
in Collibus maritimis frequenter.

443. ——— *roseus*  
in Com. Nicaeni.

444. ——— *rosmarinifolius*  
C. hirsut. suffruticosus stipul. fol. ov. calyc. hirsut. lin. in ar. di. Circa Villafancam Nicaeni.

445. ——— *glutinosus*  
C. suffruticosus stipulat. fol. linear. oppos. alternis que peduncul. villos. glutinos.  
in rupibus agri Nicaeni non infrequenter.

446. *Gavideella nigellastrum*. Decandria 3-gynia  
Cal. 5-phyl. coloratus petala 5. aequal. 2. lab. nectariifera caps. 3. connatis. polysper.  
G. fol. tenuiss. divisi.  
Nasitur in agro Nicaenibus.



447. *Telephium imperati*  
 Calyce. 5. phyll. petal. 5. aequal. Stam. 5. Styl. 3. pist. Capitul. 1. locul. 3. valv.  
 Polysp. J. fol. alterni;  
 frequent in Comit. Nicæensi  
 Caulis procumbens terrib. foliis fol. obverse ovata glauca nervi; in caulem  
 decurrentibus ima robustiora fl. terminal. umbell. alb.

448. *Linum perenne*  
 l. calycibus capitul. 9. obtusis fol. altern. lanceol. integerr.  
 in montibus circa Joissello & Lucerame.

449. ——— *hirsutum*. V. pl. lig.  
 in Com. nicæensi. passim occurrit.

450. ——— *navbonense*.  
 l. calyc. acuminat. fol. lanceolat sparsi. Hirt. scabr. acuminat  
 caul. terrib. basi ramos.  
 frequent in Jascosii inter Lucerame & Tonet C. Bellardi in mon.  
 tib. a Castiglione ad Menton & a l'Escavene ad Breglio.

451. ——— *hirtum*  
 l. calyc. lobul. fol. lanceolat. Hirt. mucronat marg. scabris.  
 locis arid. in agro Nicæensi uberrime.

452. ——— *campanulatum*  
 l. fol. bas. utrinque punctat. glandulosa.  
 inter l'Escavene & Drapp.

453. ——— *flavum*.  
 l. calyc. suberrato scabris lanceol. suberectis. panicul. ram. dichot.  
 in collibus maritimis Nicæensibus.

454. ——— *maritimum*  
 l. calyc. ovat. acut. n. n. fol. lanceolat. inferior. opposit.  
 in humidis & stagnantibus agr. nicæensis & praecipue l'Escav.  
 fl. le Var.

455. *Polycarpon tetraphyllum*  
 Cal. 5 phyll. petal. 5. persistens. cal. minor. Stam. 3. Styl. 3. Capr. uniloc. 3. valv.  
 Nicæensis agr. locis cultis sponte nascitur.  
 Caulis semipalmat. glaber prostratus ramosus fol. elliptica obtusa  
 integerr. bina in caule a ramos. originem quaternam flosc. albi.



1. loc. 3m

456. *Arenaria marina*

et *L. lina* - *laevigatum reticulatum*

in caulis

st. crumena

Mar. in m.

ad Bregia

arg. scabij.

in al. fam. dia

oppositi

igne / ecy

3. Gyn. uniloc.

thica obliqua

flor. albi.



447. *Telephium imperati*  
Calyc. 6. sepall. Petal. 4. equal. Stam. 5. Styl. 3. pist. Capsul. 1. locul. 3 valv.

44

44

45

45

45

45

45

45



11/10/71  
Arenaria marina

at El Llanero - Laguna de San Rafael - Isla de San Rafael



447. *Telephium imperati*

Stem 6. Hyl. 2. h. d. Capit. 1. locut. 3 val. v.

4.

4.

4

4

4.

4.

4.

45

Arno  
Ch. fol.  
prope

Secun  
Ch. fol.  
h. v. d.

Cor  
Chac.  
Fam. l.  
C.  
ad  
Caul. p.  
foliol.  
form

Ha

M  
Ca  
lo. ant  
For. p.  
temor

Mon  
p. f.  
i.



456. *Arenaria marina*  
A. fol. linear. longitudine internodiorum  
prope portum Villafrancae uiceni.

457. *Sedum Nicaeense*.  
S. fol. subulatis parvis basi solutis alternis cyma bifida flor. long. pedunculatis  
proxima planta nascitur in scopulosis prope Savorgio.

458. *Coriaria myrthifolia*. Dioecia decandria  
Male. Cal. 5. phyll. corol. 5. petal. calyc. immixtum connexa anther. 2. stam.  
Fem. Cal. 5. Corol. maj. styl. 5. sem. 5. petal. succulent. baccalis becca.  
C. fol. ovata oblongis  
ad sepes in Com. Nicaeensi Vulgaris planta est.  
Caul. pubescent. 2. cubit. 4. gon. cort. cinereo maculat. fol. pinnat.  
foliol. 3-nerv. bractae ovatae axemilares cor. parva viridis.  
tam. gland. anth. steril.

459. *Rhamnus alaternus*  
R. inerm. fl. dioic. stigmat. 3. foliis fol. serratis.  
in agro & collibus Nicaeensi abunde.

460. — *Rhamnus*  
R. aculeis geminis inaequal. fol. binervi. subternat. peduncul.  
solitar. fruct. marg. membranaceo clypeiformibus.  
incolit Comit. Nicaensem.

461. — *Zizyphus*  
R. aculeis geminatis altero recurva flor. digyn. fol. ovata oblong.  
glabris.  
Habitat in Comit. Nicaensi.

462. *Melia Azedarach*  
Calyc. 5. dent. persist. in med. flor. nectar. cylindric. multidentat.  
10. anther. cartilagineis breviss. pedicelatis styl. 1. fruct. drupaceo  
Circa Nicaeam Collectum habuit Cl. Alcedonius.  
Fol. parcinervis altern. petiol. foliol. petiol. integr. fl. axillar. ra-  
cemosi longe peduncul. parvi purpurei.

463. *Prunus Mahaleb*.  
P. flor. terminal. corymb. fol. ovata.  
in montibus Nicaens. Comitatus.



464. *Myrtus communis* V. *Styr. ligur.*  
in sylv. aprie. Nicaeni. maxime vulgaris. hujus Variet. Circa  
Montalbani. Nicaeni. crescit.

465. *Punica granatum*  
Frequent in Com. Nicaeni.

466. *Myrtus azarolus*  
Caelestis fol. obtus. subtrifid. dentatis.  
Comit. Nicaeni. indigena planta est.

467. *Orchis arachnites* V. *pl. ligur.*  
in coll. calidior. Nicaeni.

468. *Epipactis nidulans*  
l. aplylla flor. inveni labello bicolori  
in sylv. umbrat. Com. Nicaeni. Circa Sorpello.

469. ——— *Vulva*  
l. caul. pauciflor. label. lanceolat. lineat. lineis undulatis.  
in sylv. umbrat. montum Com. Nicaeni. minime rara.  
Nativitatem circa Blum.

470. *Pancratium maritimum* V. *pl. ligur.*  
ad ripas prope litus maris, ubique frequens.

471. *Narcissus tazetta*  
N. spatha multiflora nectar. campanul. truncat. brevior  
petalis fol. planis.  
Nicaenium collium indigena.

472. *Allium robustum*.  
A. caul. planifol. umbellifero stamin. brevissimis um-  
bella subglobosa flor. lateral. nutant.  
in Com. Nicaeni.

473. ——— *oreum* V. *pl. ligur.*  
in olivetis & arvis, Com. Nicaeni. rarum.

474. ——— *salicifolius*.  
A. caul. subteretifol. umbellifero flor. pendul. truncat.  
stamin. simpl. coroll. aequantibus.  
in colle. Albani prope Nicaeni.



75. *Lilium pomponium*.  
f. fol. spars. tubulatis, flor. reflexis, corol. revolut.



462. Myrtus communis N. Steg. Lign. in hyl. aprie. Nicaeni. maxime vulgaris. hujus Valiet. Circa  
in hyl. aprie.

46

46

46

46

46

46

46

47

47

47

Lilium  
L. fol.  
in l.

Juncus

App.  
Calyce  
petali  
Bipidus  
in po  
Rad. re  
Dial. g

Arpa

Lithra

Ane

83. Ca

84. Lyp

Ha. S. i.  
milan  
petra



475. *Lilium pomponium*.  
l. fol. spars. tubulatis, flor. reflexis, corol. revolut.  
in Comit. Nicæens. frequent.

476. *Scilla maritima*  
ad maris litus arenosa & rupes inhabitat.

477. — *italica*  
l. corymb. conferto subconico.  
loc. lapidos. non macilentis & umbratis Com. Nicæens.

478. *Aphyllanthes monspeliensis*.  
Calyx etc squamis imbricatis compositus tubulatus corolla caryophyllæa  
petalis in tubum superius limbo patente instructum dispositis stigma  
3-fidum capsula tripartita 3-angulatis 3-locularis sem. ovata.  
in pratis siccis agri Nicæensis.  
Rad. repens culm. pl. 2-palm. simp. cinct. basi vaginis utriusque foliar  
dical. graminea flor. terminal solistr. cævat.

479. *Asparagus acutifolius* V. pl. ligur.  
liberrime in Com. Nicæens.

480. *Lythrum hyssopifolia*.  
l. fol. altern. linear. flor. hercandr.  
in tabulis hum. torventis pailon frequentissima.

481. — *thymifolia*.  
l. fol. altern. linear. flor. ferrapetalis  
in tabulis hum. pailon.

482. *Anemone hortensis*  
et fol. digitalis  
Nicæa occurrat.

483. *Cactus* ~~lentus~~ *opuntia*.  
ad rupes circa Solpello.

484. *Ephedra distachya*.  
l. pedunc. opposit. ament. gemin.  
Habit. fruticosa Caul. 3-ped. Vam. fol. equiseti fol. & vami si-  
milares sub spicis stipul. 2. sicca lanceol. Vaginae cylindricæ.  
perforatae biconvex.  
in Agro Nicæensi.



485. *Pinus Pinæa*  
*p. fol. geminis primordial. solitar. ciliat.*  
 in mont. Comit. Nicaeni.
486. *Juniperus* *Oxycedrus* V. *p. Ligur.*  
 ubique in collibus & sylvis affici Nicaeni.
487. ——— *Lycia*  
*j. fol. tern. undique imbricati ovati obtusi.*  
 inter s. Dalmanzo & Soppello ad vapes.
488. ——— *phænicea* V. *p. Ligur.*  
 frequens in Comit. Nicaeni.
489. *Quercus ilex.*  
*Q. fol. ovat. oblong. indivis. serratisque subt. incan. cortice integro.*  
 Sylvarum agri Nicaeni indigena.
490. ——— *uber*  
*Q. fol. ovat. oblong. indivis. serrat. subt. tomentol. cortice vi-*  
*molo fungoso.*  
 in montanis Nicae habitat.
491. ——— *coccifera*  
*Q. fol. ovat. indivis. spinos. dentat. glabr.*  
 in sylv. Nicaeni. variorum quam præcedens.
492. *Pistacia terebinthus*  
*p. fol. impar. pinnat. foliol. ovat. lanceolat.*  
 in agro & collibus Nicaenibus ubique.  
*Foliot. 3juga inbreitalia dura fl. axill. corymbos. peduncul. ramul. vacuol.*  
*alterni fol. breviores.*
493. ——— *lentiscus*  
*p. fol. abrupte pinnatis foliol. lanceolat.*  
 in aridioribus et salsol. coll. Nicaeni. prope Mare.  
*Frutice parvus foliol. sæpius 5. 6. juga varia integerr. perfoli. mar-*  
*ginatis fl. axillar. vacuol. amenta breviter geminata sessilia dura*  
*divergentia. Fructus Nicae magnitud. primum rubelli dein nigri.*
494. *Salicornia herbacea.* Calyc. integer ventricos. Stam. 1. Sem. 1.  
 ad marit. Nicaeni (lit.).  
*Frutice sæpe cubito altior erectus. Articuli cylindrici apice clas-*  
*siove spiræ articulata. Floretes terni approximati parvi lutei.*
495. *Beta maritima*  
*B. flor. geminis in agro Nicaeni.*



Play.

15

Page - 60

No. 6844

1000000. (1000000)

have.

...helfen, was?

*benilia* *sub*

high

1

100/100

10

*apical*

1/2 hr.



178. *Chenopodium* ...  
 179. *Chenopodium* ...  
 180. *Chenopodium* ...  
 181. *Chenopodium* ...  
 182. *Chenopodium* ...  
 183. *Chenopodium* ...  
 184. *Chenopodium* ...  
 185. *Chenopodium* ...  
 186. *Chenopodium* ...  
 187. *Chenopodium* ...  
 188. *Chenopodium* ...  
 189. *Chenopodium* ...  
 190. *Chenopodium* ...  
 191. *Chenopodium* ...  
 192. *Chenopodium* ...  
 193. *Chenopodium* ...  
 194. *Chenopodium* ...  
 195. *Chenopodium* ...  
 196. *Chenopodium* ...  
 197. *Chenopodium* ...  
 198. *Chenopodium* ...  
 199. *Chenopodium* ...  
 200. *Chenopodium* ...

*B. flor. gemini* in agro Mica'enti.

*Chenopodium*  
*C.*  
 in  
 m  
*C. fol.*  
 in a  
 w  
*C. fol.*  
 in lo  
 Atvisple  
 Hemaphro  
 nym  
 From cal. 2 p  
 A. cau  
 in com  
 Caul. erect. sub  
 per. fertile fr  
 C  
 C  
 Frutic. rham  
 terminalis rac  
 Lo  
 A  
 eli  
 Vanta tota rec  
 vint virgat. f  
 hermaphrod.  
 compertus 5.  
 Urtica pi  
 la. fol. oppos  
 Comitab



496. *Chenopodium Scoparia*.  
C. fol. linear-lanceol. plan. integerrimis.  
in Cam. Nicæni. frequent.

497. ——— *maritimum*.  
C. fol. subulatis semi cylindricis.  
in agro Nicæni.

498. ——— *fruticosum*  
Saltola erecta fruticosa fol. filiform. obtusiuscul.  
in solo argillaceo ad mare libe, & secus flumen de Var.

499. *Atriplex rosea*  
Hermaphrod. Cal. 5-phyll. corol. 0. Stam. 5. Styl. 2-part. Sem. 1.  
depress.  
Fam. Cal. 2-phyll. coroll. 0. Stam. 0. Styl. 2-part. Sem. 1. compress. obliq.  
A. caule herbaceo fol. incanis serratis fruct. 4 angular. dentat.  
in Cam. Nicæni frequent.  
Caul. erect. sub 2-ped. ramoliss. ram. diffus. albus, flor. axillares glo-  
mer. sessiles fruct. compress. roseus.

500. ——— *Halimus*  
A. caule fruticoso fol. dentatoidibus integris.  
ad litus Nicæni.

Frutice vhamno simil. ramos. fol. inordinata caruosa alba flores  
terminales racemosi parvi purpurei.

501. ——— *laciniata*.  
A. cauli herbac. fol. dentatoid. dentat. subt. argenteis.  
Etiam hæc agri Nicæni, indigena planta est.

Planta tota tecta quasi epidermide seredente Caul. erect. teres  
nudus virgat. flavescens fol. except. ultim. alterna spicæ terminal.  
hermaphrod. anth. rubris fl. femin. axillares gemini Cal. fructus  
compressus 5-dent. Dente intermedio majore.

502. *Urtica pilulifera*.  
U. fol. opposit. ovat. serrat. ament. fructifer. globos.  
Comitatum Nicæni incolit.



503. *Rumex maritimus*  
*R. flor. hermaphrod. valvat. dentat. granifer. fol. lineav.*  
*in agro Nicaeni.*
504. — *bucephalosphorus*  
*R. fl. hermaphrod. valvat. dentat. nudis pedicell. plan. reflex.*  
*in arvis et sabulosis in agro Nicaeni.*
505. — *tuberosus*  
*R. flor. dioicis fol. lanceolato oblong. sagittat. hamis*  
*patentibus radice tuberosa.*  
*in prah? agri Nicaeni.*
506. — *multifidus*  
*R. flor. dioic. fol. hastat. articul. hastat.*  
*Circa Nicaeam & Villafranca ad vupes.*
507. *Polygonum maritimum.*  
*sp. flor. octandr. hyginis axillaribus fol. oval. lanceolat.*  
*semperviventibus caule protercente*  
*in arvis & sabulosis agri Nicaeni.*
508. *Mecebrum paronychia*  
*Cal. 5. phyllis cartilagin. Stam. 5. Styl. 1. capsula 1. sperma.*  
*3 flor. bract. nigr. argenteis, obvallatis caul. prosumb. fol. laevibus*  
*in sterilibus Com. Nicaeni.*
509. — *capitatum.*  
*3 floribus bract. nigr. occultant. capitul. terminali. caul.*  
*erectum. subt. villos.*  
*in agro Nicaeni.*
510. *Camphorosma monspeliaca*  
*Cal. 4. fid. legum. altern. major. reflexe. Stam. 4. Styl. bifid. capsul. 1. sperm.*  
*C. fol. hirsutis linearibus.*  
*In arenosis & aridis agri Nicaeniis praeterhumiditate Villafranca*  
*& la Cantera*  
*Caul. pedat. fruticos. subvillos. albesc. fol. altern. serrit. odor. Champhor.*
511. *Glauc Maritima.* *Cal. 5. fid. colorat. Stam. 5. Styl. 1. capsula*  
*1. locul. 5. valv. 5. sperm.*  
*Rad. geniculata caul. erect. digital. ram. oppositis fol. serrit. lineat.*  
*lanceol. obtusiusc. integerr. subcarnea late viridia glabra inter. opposit.*  
*floral. altern. fl. axill. serrit. solitar. etc albid. - dilut. purpure.*



2. *Salvia Kali* v. *fol. ligur.*



50 Rumex maritimus . . . . . l. . . . . dentat. graniter. fol. lineari.

50

50

50

50

50

50

51

51

Salola  
Cithul  
Monogyn  
B. l. l. l.  
Parale  
Caul.  
imbricat  
mali pun  
interio  
Circu  
alberet  
juncu  
Theligo  
Caul. diff  
poriti div  
alterna o  
gine / ca  
len. vol  
Cervat  
pruc  
lobia  
Folia alt  
orbiculata  
ity.  
Rannich  
in  
Caul. tenu  
apillar. i  
orientem  
interperor



512. *Salsola Kali* V. fl. ligur.  
Circa Nicæam.

513. ——— *Soda*.  
J. herbac. patula fol. inermibus  
in agro Nicæensi.

514. *Cithnus hypocithis*.  
Monogyna Cal. tubul-campanul. 4. fid. colorat. super. Anther.  
16. sessil. Bacca 8 locul. polysperm.  
Paralibica Cithi habet orobancher  
Caul. 3. 4 pollic. caud. purpur. aut sublut. fol. aut squam  
imbricatis sessilibus obliquis. Flor. termin. foliol. color. calyc.  
mali punicea sordid. luteus flor. subiacent. squam. 2. opposit.  
in densis purpureiscentes.  
Circa Bessa in Com. Nicæensi et Circa Nicæam Lica  
arbores et Cith. radic. inracent.

515. *Juncus acutus* V. fl. ligur.  
in locis maritimis et vocat.

516. *Theligonum Cynocrambe Monoclea Heptandra*  
ad muror & sepes Nicææ.  
Caul. diffusi flexuosi teretes succulenti sum. arcuati opp.  
positi divaricati fol. oleracea petiolata inter. oppos. super.  
alterna ovata obtusiuscula nervosa sublineata lobis mar-  
gine scabra. stipul. breves 3. dentata lata membranac.  
sem. volum.

517. *Ceratonia Siliqua* Vern. *Carrobo*  
fruct. siliqua magna compressa pulpa repleta  
ubique circa Nicæam Arbor  
Folia alterna abrupt. pinnat. foliola b. iuga subrecta sub-  
orbiculata integerr. crassa firma. fl. axillar. racemosi sex-  
lites.

518. *Zannichellia pumila*  
in Com. Nicæensi ad rivulos.  
Caul. tenues ramosi. crispiter densiorefformant fol. sessil. simpl.  
capillar. integerr. ima alterna summa opposita vagina lata  
orientem ramum excipit fl. axillar. solitar. sessil. capsul.  
subarbores.



519. *Ruppia maritima*

A. foliis acutissimis gramineis longior. fruct. fere umbell.  
ad lit. Mar. Nicaeni.  
Caul. demers. herbac. articulat. fol. simplic. altern. sessil li-  
nearia flor. axillar. solitar. pedunculati.

520. *Xoiteva Oceanica.*

? pericarpis, pedicellatis oliviformibus.  
ad mari, Nicaeni (h. h.)  
Differt a sequenti radice in pilum complicatis &  
fruct. pedunculatis.

521. ——— *marina*

? pericarpis sessilibus.  
ad lit. mari Nicaeni.

Planta aquis marinis innatans, flexuosa radice fasciculos folior.  
angustorum emittit. fascicul. caulini, culmi graminei ad inter  
singuli & singulis radicalibus ramulis prodeunt, valde infirmi  
laece geniculati flexiles natantes. Fol. altern. vaginant li-  
nearia verticaliter obtusa glabra mucosa sapore viridia.  
in apice ramulorum 2-3-4. folia unica vagina inclusa  
radices semiunciales linea paullo crassiores.

522. *Blitum Capitatum Monandra Monogynia*

Caul. 3-fidus corol. o. sem. 1. Calyc. - baccato fl. plur. in glomer.  
colliguntur stabilius Chenopodii  
fl. capitellis spicatis terminalibus  
Non infrequens in Comit. Nicaeni. Sicut alia.  
Caul. pedat. simplic. fol. longe petiol. hastata semihiloba quan-  
doque integra dentata. global. florales sessiles corollae.

523. *Smilax aspera.* Vern. *Gramon d' Montagna*

A. caule aculeato fol. dentat. aculeatis cordatis 9-nerviis.  
Vulgaris est in Comitatu Nicaeni.

Caul. scandens cirrhis ad bat. petiol. fol. altern. brev. petiol. sicca  
virida acuta quandoque auriculata. Fl. axillar. glomerato-  
spicati corol. alba odora baccae rubra vel nigra.

524. *Ruscus hypoglossum*

in sacris, asperis, incultis, sterilibus Nicae & Lobelii.



5. *Olyvil alba.*

in Comitatu Nicaeni in vallis ad Villam Sanctam de 1715



51 *Amphibia maritima*

5.

5

5

5

52

36 Olyvil  
in Com  
mural  
Frabre  
11. perma  
lutan  
38 Nicun  
27 Arun  
Radice  
salmar  
28. Clav  
Phleu  
ph. spic.  
in Ag  
Phalaris  
V.  
ph. spic.  
param  
Allope  
Pha  
2.2  
C/



525. *Osyris alba*.  
in Comitatu Nicaeni praesentem ad Villa Franca depre-  
munt vulgari.

Frutic. 2. 3. pedalis striatus fol. altern. serr. acut. integerr.  
fl. terminal. conferti subracemosi pedunculati Calyce viridi  
luteo fruct. erecto umbilicatus suber.

526. *Nicum album* V. pl. ligur.  
locis maritimis frequentior quam in praedemonio.

527. *Arum aviculare*. V. pl. ligur.  
in Sylvis Nicaenibus copiose.  
Radice globosa verrucosa fol. atro virentia rugosa petiol.  
palmar. maculati flor. pallide atro purpur.

528. *Nardus arvensis* V. pl. ligur.  
in agro Nicaeni

529. *Phleum Schoenoides*  
ph. spic. ovalis obsoletis fol. breviss. mucronat. impleticant.  
in Agro Nicaeni

530. *Phalaris Canariensis*  
ph. panicula subovata spic. termi. glumis carinatis  
in Comit. Nicaeni.

531. ——— *paradosca*  
ph. spica cylindrica floralis mucronatis neutris plurimis infimis  
praemortis.  
in Comit. Nicaeni.

532. *Alopecurus moissetianus*.  
A. panicula pubescentia calyc. scabr. coroll. aristatis  
Ad limit. agrorum in Comit. Nicaeni.

533. ——— *ciliatus*  
A. culm. spicab. erect. glumis ciliatis.  
C. Bellard: in praesentibus editionibus d. Breglio.

534. ——— *litoreus*  
phalar. calyce divaricato plano ovato ciliato glum. floral. interior.  
2. dentata.  
C. Bellard: ad litora maris a Clisara ad flum. le Var.



535. *Milium leudigerum*.

Ch. panicul. subspicat. floribus aristatis.  
in Comit. Nicaeni.

536.

— *paradoxum*

Ch. flor. panicul. aristat.

in agro Nicaeni: Cl. Bellardi: ad ruper mari proximam

537.

*Agrostis miliacea*

A. spet. arboribus arist. termin. veta Mich. medior.

Circa Nicaenam legit Cl. Bellardi.

538

— *pinguis*

A. panicula compacta multa foliis involutis; rigidi pinn.  
genuibus superior. oblique oppositi.

Legit Cl. Bellardi locis aridis Maris Nicaeni: prope.

539.

— *aculeata*

Schœnus culmi tereti ramulo capitul. terminal. involucri  
3 phyllo brevissimo, rigido patente.  
in agro Nicaeni.

540.

— *minima*

A. panicula multa foliis pinn.  
in agro Nicaeni.

541.

*Rotbolla incurva* V. pl. Ligur.

Nicaea in agro mari proximo.

542.

*Lagurus ovatus*.

Gen. Calycis glumae aristatae aristae villosae. Folliculus biglumis  
valvula majori terminata duplici aristae cum tercia ad dorsum ac-  
cedente valvula altera minori acuminata Semen oblongum folli-  
culo arcte inclusum aristatum.

h. spica ovata aristata

in agro Nicaeni: frequens.

543.

— *cylindrica*

h. spica cylindrica multa

etiam agri Nicaeni: indigena planta est.

544.

*Cynurus cuneus*.

C. panicul. spic. sterilibus pendul. ternat. flor. aristat.

in agro Nicaeni. loc. steril. mari proximo.



45. *Lolium tenue*  
h. *spica multiflora* *tereti spiculit trifloris*



5. *Milium leudigerum.*

5

52

51

53

54

51

5

5.

5

*Silene tenuis*  
L. sp.  
in Co.

*Don. rigida*  
ad Gr.

*Virga Vir*  
B.  
in a

— m  
B.  
pro

— m  
B.  
in

*Stomus*  
B.  
in

—  
h

—  
C

— m  
B.  
92  
20

—  
B.  
in

*Ferrea*  
*Traceno*  
in pr



545. *Lolium tenuis*  
h. spica multiflora tereti spiculis trifloris.  
in Com. Nicaeensi, praesens & durne h. fasciis.
546. *Poa rigida* V. M. L. g. v.  
ad limites agrorum secus vias locis aridis in Com. Nicaeensi abunde.
547. *Phlox vivens*  
h. spiculis ovalis calyce florulis (7) aequali.  
in agro Nicaeensi.
548. ——— *maxima*.  
h. spiculis cordatis florulis 17.  
frequens in agris Comit. Nicaeensi.
549. ——— *monispermatana*.  
h. spica nutante simplici spicul. altern. pedunculatis  
subsistatis ovalis calyce 5-floris.  
in agro Nicaeensi.
550. *Bromus rubens*.  
h. panic. fascicul. spicul. suberectis. villis aristis erectis.  
in agro Nicaeensi.
551. ——— *inermis*  
h. panicula erecta spicul. suberectis, tubulatis nudis  
subnullis.  
(C. Bellardi circa Torpello.
552. ——— *plukenetii*  
h. fol. convolut. pungent. locis binis terminalibus.  
C. Bellardi legit locis aridis Nicaeae circa Villafanca.
553. ——— *madritensis*  
h. panicul. varior. paucis - erect. spicul. linear. internod.  
geminis pedicellis superne incrassatis.  
locis siccis circa Tashion in Com. Nicaeensi.
554. ——— *distachyos*.  
h. spicis duabus erectis affinis.  
in Com. Nicaeensi. ad vias & agrorum limites.
555. *Festuca pharicoides*.  
F. racemo indiviso spicul. altern. suberectis. tereti fol. involut.  
mucronatis pungent.  
in praetis & ad sepes in Com. Nicaeensi.



556. *Triticum junceum*.

T. calyc. truncat. 5. fl. fol. involutis.  
ad Vias & loca sicca in agro Nicaeensi.

557. ——— *tenellum*

T. calyc. subq. fl. fl. mult. acut. fol. setaceis.  
in Comitatu Nicaeensi.

558. ——— *unilaterale*.

T. calyc. unilateral. alternis mult.?  
Nascitur ad litus maris Nicaeensis.

559. ~~*Pharus*~~ *avenarius*.

P. spica erecta arcta calyc. tomentis. fl. longioribus.  
in arena ad maris litus.

560. *Hordeum geniculatum*

H. culm. decumbent. foliis obrecto summo spatheo  
spica subjecto supremo nodo incurvato.  
C. Bellardi legit ad litus maris Nicaeensis prope  
portum di Limpia.

561. ——— *secalinum*.

H. fl. lateral. mascul. aristis involucri setaceis  
scabris  
Habitat in agro Nicaeensi.

562. *Andropogon distachyon*

A. spic. 2. terminali. culmo indiviso.  
Ubique ad vasa et in sacris Comit. Nicaeensis.

563. *Aegilops ovata*.

Gen. locusta 3. vel 4. fl. fl. perfectis laterales & masculis in  
terminis continens calyc. biglumis glum. ovata aristat. foliis  
2. valv. valvula exterior ovata triplici aristata terminata interiora  
multa sem. oblongum valvulae foliis interiori aequalem.  
A. spica ovata calyc. omnibus 3. aristatis.  
Loci siccis ubique in Comit. Nicaeensi.

564. ——— *vincialis*.

A. spica aristata calyc. inferioribus biaristatis.  
in agro Nicaeensi.



in foliis

foliis longis

in foliis

foliis

foliis

foliis

foliis

foliis

foliis

foliis



556. *Pharusia* *maritima*  
F. caulis breviss. foliis lanceolatis  
acutis & longioribus apice subulatis

557. *Pharusia* *maritima*  
F. caulis breviss. foliis lanceolatis  
acutis & longioribus apice subulatis

558. *Pharusia* *maritima*  
F. caulis breviss. foliis lanceolatis  
acutis & longioribus apice subulatis

559. *Pharusia* *maritima*  
F. caulis breviss. foliis lanceolatis  
acutis & longioribus apice subulatis

560. *Pharusia* *maritima*  
F. caulis breviss. foliis lanceolatis  
acutis & longioribus apice subulatis

561. *Pharusia* *maritima*  
F. caulis breviss. foliis lanceolatis  
acutis & longioribus apice subulatis

562. *Pharusia* *maritima*  
F. caulis breviss. foliis lanceolatis  
acutis & longioribus apice subulatis

563. *Pharusia* *maritima*  
F. caulis breviss. foliis lanceolatis  
acutis & longioribus apice subulatis

564. *Pharusia* *maritima*  
F. caulis breviss. foliis lanceolatis  
acutis & longioribus apice subulatis



*[Faint, illegible handwriting throughout the page, possibly bleed-through from the reverse side.]*



565

566

567

568

569

570

571

572

573

574

575



565. *Cenchrus capitatus*  
C. spica ovata simplici  
in Monte Syra & circa Ioppello C. Bellardi.

566. *Cornu lachryma jobi*  
in agro Nicaeni legit D. Guidice.

567. *Cavere hirta*  
C. caule foliis, capitulisq. hirsutis.  
in agro Nicaeni locis sylvestribus vulgaris.

568. *Scirpus mucronatus*  
S. culmo 3. angulo modo acuminato spicis conglomeratis  
sessilibus lateralibus.  
Nascentur in Maritimis Nicae.

569. ——— *maritimus*  
S. culmo trigonito panicula conglomerata foliacea spicularum  
ignavis 3. fidi intermedia subulata.  
in com. Nicaeni frequent.

570. *Schœnus mucronatus* V. pl. Lignv.  
ad litora maris a Monaco ad Menton.

571. *Lichen Rocella*  
L. punctulatus solidus asphyllus lobramosus tuberculis alternis  
ad Rupes maris Nicaeni.

572. *Fucus serratus* V. pl. Lignv.  
in Mari Nicaeni.

573. ——— *vesiculatus* V. ut supra  
ibidem.

574. ——— *canaliculatus*  
F. proinde plana dichotoma integerrima canaliculata  
linearibus multiplicationibus tuberculatis in partibus obtusis  
ad Rupes litoris Nicaeni.

575. ——— *nodosus* V. pl. Lignv.  
ad mare Nicaeni.



576. *Fucus siliquosus*  
*F. fronde compressa ramosa fol. distich. alternis integerrimis* / *fructificationibus pedunculatis oblongis mucronatis.*  
*ad rupes littoris Nicæensis.*
577. — *selaginoides* V. *sp. lign.*  
*ad litus Nicæense*
578. — *fœniculaceus* V. *sp. lign.*  
*in rupibus littoris Nicæensis.*
579. — *abrotanoides* V. *sp. lign.*  
*in mari Nicæensi*
580. — *natus*  
*F. caud. filiformi ramoso fol. lanceol. serratis fructificat. glo-*  
*bos. pedunculatis.*  
*in mari Nicæensi*
581. — *acinarius* V. *sp. lign.*  
*in mari Nicæensi*
582. — *linearis*  
*F. dichotomus planus linearis acutus vertic. ovatis spicatis*  
*Ad rupes littoris Nicæensis.*
583. — *fastigiatus*  
*F. fronde filiformi dichotoma ramosissima fastigiata*  
*obtus.*  
*ad rupes littoris Nicæensis.*
584. — *robustus* V. *sp. lign.*  
*in mari Nicæensi.*
585. — *ceranoides*  
*F. frond. plana dichot. integer. punct. lanceol. fructific.*  
*hiberculat. bifid. terminal.*  
*in mari Nicæensi.*
586. — *dentatus*  
*F. frond. membranac. nerv. alternatim pinnatis. lacin. apice*  
*evolutis*  
*in mari Nicæensi.*



repin

201 Thoms. Mendocino

2. 1. 1887

Wash

Wash

Wash

Wash

Wash



576. *Ficus siligmaria*  
*F. fronde compressa, moia fol. distich. alternatim integerrimis* Wuch.

57

57

57

58

58

58

58

58

58

58



587. *Tucus splendens* videri  
F. parvus caulis. teret. summis. membranac. dilat. & lacerat.  
Ad ruper maris Nicaeensis.

588. — *Ramificatus*  
F. caulis. teret. ramulosis. fol. altern. subulat. veteiculis.  
Volund. iunior. sens. lib. & ciliatibus  
Ad ruper maris Nicaeensis.

589. — *mucoides* N. sp. lignu.  
in mari Nicaeensi.

590. — *helminthochortus*  
Conspersa helminthoch. dichotoma uncinatis & periculis  
ramis horizontalibus  
Lechus est in rupibus maris Nicaeensis.

591. — *veluticulus*  
F. filiformis. teret. ramulis. ramis simplicibus verticatis  
Volund. lateralibus  
ad ruper maris Nicaeensis.

592. — *longissimus*  
F. teret. rubens. minus. ramulis in longum proterentibus  
in mari Nicaeensi.

593. — *parvus*  
F. filiformis. teret. ramulosis. ramis alternis ramulosis  
confertis. fructibus fructibus globosis.  
ad mare Nicaense

594. — *plicatus*  
F. capillaris. uniformis. ramulosis. implicatus sub  
diaphanum  
ad ruper maris Nicaeensis.

595. — *corneus*  
F. cartilagineus. caule teretibus. compresso ramulosis.  
limbo. ramis subplanatis. laciniis acutis. fructibus  
ad ruper maris Nicaeensis.



596. *Fucus confervoides* V. pl. ligur.  
in mari Nicaeensi.

597. — *plumosus* V. pl. lig.  
*ibidem*.

598. — *pinnatifidus*  
F. frond. plan. vamo. vami dentato pinnatifidi  
marginibus callosis  
ad super mare Nicaeensi.

599. — *cartilagineus*.  
F. fronde cartilaginea compressa supradecomposita  
pinnata laciniis linearibus  
Ad super mare Nicae.

600. — *volubilis*  
F. fronde plana spirali pectol. repand. dentata.  
ad mare Nicaeense.

601. — *spiralis*  
F. frond. plan. dichot. integerr. punctata inferne  
lineari canaliculata tuberculat. tubercula b. geminis.  
ad mare Nicaeense.

602. — *palmatus*  
F. fronde palmata plana.  
ad super mare Nicaeensi.

603. *Ulva pavonia*  
U. plana veniformis serrilis decussatum striata  
in mari Nicaea.

604. — *umbilicalis*  
U. plana orbiculari serrili peltata coriacea  
Habitat ad super mare.

605. — *intestinalis* V. pl. ligur.  
in mari Nicaea.



606. *Ulva compressa* V. pl. fig.

planifolia

supradu

repandens

chala infera  
scula genit

stadium / miala

ta rosacea

ad vasser (hnt) viderunt



and

5

5

5

60

60

60

60

6.

60

606

60

60

605

610.

611.

612.

613.

614-

615.

16.



606. *Ulva compressa* V. pl. lig.  
Habitat in mari Nicaea.

607. — *Lactuca* v. ut lupan  
idem

608. — *Linga*  
idem.

609. *Conferva fasciculacea*.  
Gen. Tubercula inaequalia in fibris capillaribus longissim.  
C. filament. aequal. ramisiff. ramis ramulif. longiss.  
Habitat in rupibus maris Nicaeensis.

610. — *tomentosa*  
C. filament. aequal. ramisiff. ram. simpl. confert. fuscis.  
ad litus maris Nicaeensis.

611. — *litorealis*  
C. filam. aequal. ramisiff. elongat. scabrisculis.  
Habitat in rupibus litus Nicaeensis.

612. — *ovulginosa*  
Plant. marinis litus Nicaeensis innascitur.  
C. filament. ramis. moll. digit. brevior. viridiss.

613. — *Scoparia*  
C. fil. proliferis fasciatis lithis  
Habitat in mari Nicaeensis.

614. — *Corallina*  
C. filis geniculatis dichotomis  
in litore maris Nicaeensis.

615. — *polymorpha*  
C. filamentis geniculatis ramis fasciculatis  
ad litus maris Nicaeensis.

616. — *rupetris*  
C. filamentis geniculatis ramisiff. virid.  
ad rupes litus Nicaeensis.



617. *Conserua elongata*  
C. filam. genicul. ramisill. Ramuli longiss. distant. acut.  
Habitat in litore Nicaeensi.

618. ——— *fusca*  
C. filam. geniculat. ramisillim. Ramuli. altern.  
simplicibus  
in litore Nicaeensi.

619. *Aristolochia rotunda*  
A. fol. cordat. subressit. obtus. caul. inferm. flor. solitar.  
in sylvis collinis umbratis Nicaeensibus.

620. ——— *longa*  
A. fol. cordat. petiol. integerr. obtusiuscul. caul. inferm.  
fl. solitarii  
Sponte nascitur in vineis Nicaeensibus.

621. *Chamaerops humilis*  
Ch. non palmat. plicat. stipit. spinos.  
Palma humilis  
in agro Nicaeensi.  
Auctantii

623. *Valeriana cornucopiae*  
V. flor. diandr. ringent. fol. ovat. sessil.  
Nascitur in Comit. Nicaeensi.

624. *Antirrhinum origanifolium*. V. pl. Tende  
supra l' Escavena.

625. *Scabiosa hybrida*  
S. corollat. 4fid. radiant. fol. infer. pinnatis pinnul. obtus. dent. Ramis integris.  
in arvis et pascuis agris Solypitellensibus.

626. *Senecio squallidus*  
S. coroll. radiant. calyc. longior. integ. fol. pinnatis f. lacin. linear. distantibus.  
Collis elicaeensis inhabitat et frequens est in Olivetis & locis lapideis  
praesertim prope Saorgio.



Dist. 1-20

Dist. 1-20

Dist. 1-20

Dist. 1-20

Dist. 1-20

Dist. 1-20

Dist. 1-20

1840

1840

1840

1840

1840

1840

1840

1840

1840

1840



617 *Conspicua elongata*  
C. p. elongata, thallus linearis,  
habitat in littore. (Linn.)

618 *Conspicua elongata*  
C. p. elongata, thallus linearis,  
habitat in littore. (Linn.)

619 *Conspicua elongata*  
C. p. elongata, thallus linearis,  
habitat in littore. (Linn.)

620 *Conspicua elongata*  
C. p. elongata, thallus linearis,  
habitat in littore. (Linn.)

621 *Conspicua elongata*  
C. p. elongata, thallus linearis,  
habitat in littore. (Linn.)

622 *Conspicua elongata*  
C. p. elongata, thallus linearis,  
habitat in littore. (Linn.)

623 *Conspicua elongata*  
C. p. elongata, thallus linearis,  
habitat in littore. (Linn.)

624 *Conspicua elongata*  
C. p. elongata, thallus linearis,  
habitat in littore. (Linn.)

625 *Conspicua elongata*  
C. p. elongata, thallus linearis,  
habitat in littore. (Linn.)



627. *Hieracium praenotum*.  
in pratis editioribus di. Soppello.
628. *3bevis ciliata*  
3 fol. lingulat. spatulat, acutis, ciliatis, fruct. corymbois.  
Inter la Scavona & Mizza loci dactylis et in spatis ad ora  
sylvarum.
629. — *linifolia*  
3 herbacea fol. linear. integerr. caul. dextratis, caul. paniculat.  
corymb. hemisphaericis.  
In arenosis collibus agri Nicacensis versus flumen de Var.
630. *Brassica oleracea*  
A radice caulescente tereti carnosa.  
Sponte nascitur inter Villafranca & Monaco in rupibus mari  
proximis.
631. *Cheiranthus incanus*  
C. fol. lanceol. integerr. obtus. incanis, siliguis apice  
truncatis compressis, caule suffruticosa  
Inhabitat rupes mari proximae prope Villafranca  
atque spontaneus est.
632. — *micropsidatus*  
C. siliguis apice 3-dentatis, fol. lyvatis.  
in Sabulosis ad Muri Nicacensis litus.
633. *Euphorbia Taurinensis*  
in vineis Nicacensibus frequentissima.
634. — *Pappaver dubium*  
Vulgare in com. Nicacens, ibique variat. fl. albo.
635. *Medicago circinnata*  
M. legum. reniform. marg. dentat. fol. pinnat.  
in agro Villafrancae Nicacens. prope mare versus Monaco
636. — *laciniata*  
Frequent in arvis circa Soppello & Breglio.



637. *Hedysarum Cristagalli*.  
H. fol. pinnat. Legum. 1. specim. aculeatis crist. lacin. lanceol. denticul.  
ad mare Nicaeni: latus in Sabulosis.

638 *Cachrys Moriloni*.  
Gen. Umbella magna involucri univers. & partial. polyphyllum.  
flores aequales petala integra angusta; semina oblonga gibba  
crassa ad latus non explanata substantia crassa fungosa.  
C. foliol. linear. exsertim binis aut ternis punctibus semiovatis  
lanceolatis utrinque rotundatis  
Supra Breglio loco dicto Mauciana

639 *Smyrniolum olusatrum*.  
Non infrequent in Comit. Nicaeni.

640 *Bupleurum Gerardi*  
Frequentissime occurrit in Comit. Nicaeni.

641. *Bubon macedonicum*.  
G. Umbella magna involucri univers. & partial. polyphyllum.  
petala oblonga sem. ovata striata villosa.  
B. foliol. rhombico-ovatis crenatis umbell. numerosissimis.  
in pascuis siccis prope Nicaeam.

642. *Geranium malacoides*.  
Non modo circa Nicaeam videtur sed frequent est in Comitatu  
Nicaeni.

643 *Lavatera punctata*  
L. caul. herbac. fol. subrotund. 5. angular. deinde hastatis; peduncul.  
1. floris axillar. flor. campanulatis.  
in agro Nicaeni: frequens  
caulis rubet & minutis punctis albis aspersis & pedatis.

644. *Ranunculus agrarius*  
R. radic. fibros. ann. fol. hirsut. semi-3-lob. petiol. acute serratis  
calyc. hirsut. retroflexis  
in agro Collinis Comit. Nicaeni in praesentem in quibus  
agrum aliquandiu perficit.



Real. de...

Real. de...  
Real. de...  
Real. de...

Real. de...

Real. de...

Real. de...

Real. de...

Real. de...

Real. de...

Real. de...



637. *Hedysarum Cristagalli*.  
H. fol. pinnat. Legum. 1. 1 per m. aculeatis crist. lac. lanceol. denticul.  
1. 1 1 2 1. 6. 10. 5

£

6

6

E

45. H.

248 Ga

547. A

2649 —  
A. b.  
1. a. b.

649. Fri

3650. A.  
A.c.  
22/1/19

661. Ag  
in C

852. Sky

653. —  
St. avit  
Fregu  
Villap



645. *Statice speciosa*  
limonium fol. lanceol. mucronat. radicalibus caul. ancipit di-  
chotomo stipit. simplicibus.  
Occurrit Circa Ventimiglia.

646. *Galanthus autumnalis*.  
leucos. spatha multiflora stylo filiformi.  
in rupibus prope Villafanca Nicæentium.

647. *Allium magicum*.  
A. caul. planifol. umbellifer. ramulo bulbifero stamin.  
simplicibus  
Com. Nicæent. indigena planta est.

648. ——— *monspellanum*.  
A. caulis tereti solido umbell. fastigiata. petal. linear. fol. lanceol. stamin.  
tubulatis  
in Olivetis Villafanca Nicæent.

649. *Fritill. involucr.* N. sp. Monvey.  
in Com. Nicæent. supra Breglio

650. *Atriplex hastata*.  
A. caul. herbaceo fol. 3. angular. calyc. valvat. femineis magnis  
deltoidibus sinuatis 3. angular. subaspreis  
in arena litoreis Nicæentis.

651. *Agrostis miliacea*  
in Com. Nicæent. frequens est secus torrentes locis siccior.

652. *Stipa juncea*  
St. arist. nud. rect. calyc. sem. longior. fol. intus (arvibus).  
locis siccioris calidior. in agro Nicæent. prope Villafanca.

653. ——— *aristella*.  
St. aristis nudis rectis calyce duplo longioribus germin. lanatis.  
Frequens locis rupestribus & calidioribus C. Nicæ. abunde circa  
Villafanca.



654. *Brixa minor*  
Frequentest secus arva locis calidioribus agri Nicaeensis.

655. *Bromus plukenetii*  
Frequentissimus est in collibus calidioribus & siccis Com. Nicaeensis.

656. *Melica Benthii*  
M. spica ramosa unilateralis fertilibus floribus petalo subciliato  
provenit locis rupestribus aridis in Com. Nicaeensis.

657. *Avena fama*  
Frequentissima est secus agros & in pascuis collium Calidionum  
Nicaeensium.

658. — *bromoides*  
A. subspicata spiculis binatis altera pedunculata aristata  
varicatis calyc. 8-floris.

659. *Holcus arenarius*  
H. spica conica tomentosa lucida glumis bifloris hermaphrodito sessili masculo pedunculato  
in arena ad mar. Nicaeensis litus.

660. *Agilops truncialis*  
provenit praesertim secus agros inter Sospello & Carigliana

661. *Cyperus distachyos*  
C. culmo Wignebo nudo spicis sessilibus lateralibus  
radice perenni.  
locis spatiosis inter Nicam & le Var.

662. — *globosus*  
C. rad. perenn. caespitosus planifol. culm. Wignebo. fol.  
brevior. capit. globos. terminal. involucri. 4-phylla.  
in pratis spongiosis secus le Var.



11. 1. 2. 3. 4. 5. 6. 7. 8. 9. 10. 11. 12. 13. 14. 15. 16. 17. 18. 19. 20. 21. 22. 23. 24. 25. 26. 27. 28. 29. 30. 31. 32. 33. 34. 35. 36. 37. 38. 39. 40. 41. 42. 43. 44. 45. 46. 47. 48. 49. 50. 51. 52. 53. 54. 55. 56. 57. 58. 59. 60. 61. 62. 63. 64. 65. 66. 67. 68. 69. 70. 71. 72. 73. 74. 75. 76. 77. 78. 79. 80. 81. 82. 83. 84. 85. 86. 87. 88. 89. 90. 91. 92. 93. 94. 95. 96. 97. 98. 99. 100. 101. 102. 103. 104. 105. 106. 107. 108. 109. 110. 111. 112. 113. 114. 115. 116. 117. 118. 119. 120. 121. 122. 123. 124. 125. 126. 127. 128. 129. 130. 131. 132. 133. 134. 135. 136. 137. 138. 139. 140. 141. 142. 143. 144. 145. 146. 147. 148. 149. 150. 151. 152. 153. 154. 155. 156. 157. 158. 159. 160. 161. 162. 163. 164. 165. 166. 167. 168. 169. 170. 171. 172. 173. 174. 175. 176. 177. 178. 179. 180. 181. 182. 183. 184. 185. 186. 187. 188. 189. 190. 191. 192. 193. 194. 195. 196. 197. 198. 199. 200. 201. 202. 203. 204. 205. 206. 207. 208. 209. 210. 211. 212. 213. 214. 215. 216. 217. 218. 219. 220. 221. 222. 223. 224. 225. 226. 227. 228. 229. 230. 231. 232. 233. 234. 235. 236. 237. 238. 239. 240. 241. 242. 243. 244. 245. 246. 247. 248. 249. 250. 251. 252. 253. 254. 255. 256. 257. 258. 259. 260. 261. 262. 263. 264. 265. 266. 267. 268. 269. 270. 271. 272. 273. 274. 275. 276. 277. 278. 279. 280. 281. 282. 283. 284. 285. 286. 287. 288. 289. 290. 291. 292. 293. 294. 295. 296. 297. 298. 299. 300. 301. 302. 303. 304. 305. 306. 307. 308. 309. 310. 311. 312. 313. 314. 315. 316. 317. 318. 319. 320. 321. 322. 323. 324. 325. 326. 327. 328. 329. 330. 331. 332. 333. 334. 335. 336. 337. 338. 339. 340. 341. 342. 343. 344. 345. 346. 347. 348. 349. 350. 351. 352. 353. 354. 355. 356. 357. 358. 359. 360. 361. 362. 363. 364. 365. 366. 367. 368. 369. 370. 371. 372. 373. 374. 375. 376. 377. 378. 379. 380. 381. 382. 383. 384. 385. 386. 387. 388. 389. 390. 391. 392. 393. 394. 395. 396. 397. 398. 399. 400. 401. 402. 403. 404. 405. 406. 407. 408. 409. 410. 411. 412. 413. 414. 415. 416. 417. 418. 419. 420. 421. 422. 423. 424. 425. 426. 427. 428. 429. 430. 431. 432. 433. 434. 435. 436. 437. 438. 439. 440. 441. 442. 443. 444. 445. 446. 447. 448. 449. 450. 451. 452. 453. 454. 455. 456. 457. 458. 459. 460. 461. 462. 463. 464. 465. 466. 467. 468. 469. 470. 471. 472. 473. 474. 475. 476. 477. 478. 479. 480. 481. 482. 483. 484. 485. 486. 487. 488. 489. 490. 491. 492. 493. 494. 495. 496. 497. 498. 499. 500. 501. 502. 503. 504. 505. 506. 507. 508. 509. 510. 511. 512. 513. 514. 515. 516. 517. 518. 519. 520. 521. 522. 523. 524. 525. 526. 527. 528. 529. 530. 531. 532. 533. 534. 535. 536. 537. 538. 539. 540. 541. 542. 543. 544. 545. 546. 547. 548. 549. 550. 551. 552. 553. 554. 555. 556. 557. 558. 559. 560. 561. 562. 563. 564. 565. 566. 567. 568. 569. 570. 571. 572. 573. 574. 575. 576. 577. 578. 579. 580. 581. 582. 583. 584. 585. 586. 587. 588. 589. 590. 591. 592. 593. 594. 595. 596. 597. 598. 599. 600. 601. 602. 603. 604. 605. 606. 607. 608. 609. 610. 611. 612. 613. 614. 615. 616. 617. 618. 619. 620. 621. 622. 623. 624. 625. 626. 627. 628. 629. 630. 631. 632. 633. 634. 635. 636. 637. 638. 639. 640. 641. 642. 643. 644. 645. 646. 647. 648. 649. 650. 651. 652. 653. 654. 655. 656. 657. 658. 659. 660. 661. 662. 663. 664. 665. 666. 667. 668. 669. 670. 671. 672. 673. 674. 675. 676. 677. 678. 679. 680. 681. 682. 683. 684. 685. 686. 687. 688. 689. 690. 691. 692. 693. 694. 695. 696. 697. 698. 699. 700. 701. 702. 703. 704. 705. 706. 707. 708. 709. 710. 711. 712. 713. 714. 715. 716. 717. 718. 719. 720. 721. 722. 723. 724. 725. 726. 727. 728. 729. 730. 731. 732. 733. 734. 735. 736. 737. 738. 739. 740. 741. 742. 743. 744. 745. 746. 747. 748. 749. 750. 751. 752. 753. 754. 755. 756. 757. 758. 759. 760. 761. 762. 763. 764. 765. 766. 767. 768. 769. 770. 771. 772. 773. 774. 775. 776. 777. 778. 779. 780. 781. 782. 783. 784. 785. 786. 787. 788. 789. 790. 791. 792. 793. 794. 795. 796. 797. 798. 799. 800. 801. 802. 803. 804. 805. 806. 807. 808. 809. 810. 811. 812. 813. 814. 815. 816. 817. 818. 819. 820. 821. 822. 823. 824. 825. 826. 827. 828. 829. 830. 831. 832. 833. 834. 835. 836. 837. 838. 839. 840

Town - Akron

1.  $\frac{1}{2} \log \frac{1}{2}$

*Californicus*

1/2 0.001

*Epilobium* *her-*

Ms. B. Castiglione

*Lupinus (ibid)*

✓

W. g. n. v. fol.  
v. 4. phylla.



654. *Arctostaphylos*

655.

656.

657

658

659

660

661

662

653. *Lichen*  
h.  
m.  
in

654. *Arctostaphylos*

655. *Bulbo*  
Cord. int.  
super.  
ad radicem  
in C.

656. *Tulipa*  
T. fl. 2.  
in

657. *Anemone*  
A. fol. v.  
Mars.  
Flor

658. *Penstemon*  
P. fol. c.  
Calyc.  
D. pro  
in

659. *Genipa*  
G. fol.  
Jaki



663. *Lichen diffusus*  
h. pyxidatus endiv. (virg. fol. in fundibul. explanat)  
unguent  
in pascuis sterilibus & dumetis Com. Nicæentis.  
Appendicis.

664. *Anchusa officinalis*  
A. fol. lanceol. spic. imbricatis secundis  
Nascitur in Com. Nicæent. ad Vitis.

665. *Bulbocodium Vernum*.  
Corol. in fundibul. form. 6. petal. ungu. angust. staminifer. Corol.  
supra. Genus Colchico a quo differt petalis divisis, uique  
ad radicem <sup>unguibus</sup> petalis staminiferis.  
in Com. Nicæensi reperit Marchio De-Suffren.

666. *Tulipa geneviana*.  
T. fl. erecto fol. ovat. lanceolatis.  
in collibus locisque incultis circa Nicæam.

667. *Anemone Coronaria*  
A. fol. radical. ternato decompositis involucri folioso.  
Nascitur in montibus Nicææ locis incultis.  
Flor mire ludit color. violac. subalbido rubro & stramineo.

668. *Jeneverium tomentosum*  
J. fol. crenat. tomentos. flavicantibus floribus aureis  
calycibus & foliis floralibus intente flavis distinguitur a  
J. folio.  
in Com. Nicæenti.

669. *Genista Scoparia*  
G. fol. simplicibus hirsutis ramis (o. angulatis cinereis)  
satis frequent in Com. Nicæent. loc. montan. calidior.



670. *Tragopogon porrifolium*.

*T. calyc. coroll. radio longiori longioribus fol. integris strichis pedunc.  
superne incrassatis.*

*Nascitur in Com. nicolensi Obsev. D. Galli.*

*Flor. purpureo carat. prodest.*

671. *Acer opulifolium*

*A. fol. lobis subrotundis obtuse dentatis fructibus cymosis.*

*in Com. Nicolensi. Sponte nascitur in rupibus a Fontana ad  
Saurgium.*


672. *Pteris Cretica*

*pp. frond. pinnat. pinn. opposit. lanceol. serrat. bas. angustat. in pinn. sub-*

*tripartitis*

*in Vallecula umbrosa agri Nicolensis decescit. Cl. Desuffren.  
etiam in sylvaticis umbrosis Tenda.*

Gunei Die 11.  
Junii 1800





intelligit

67.

in hoc (ymeri)  
apud a Fontem

an angustit. in fin.

in fin. t. 17. 18.

nam

Die 11.

1800





67

671

672







6

67

67









































































































— 1 —